


# nPDF studies in pPb collisions with the CMS detector

Andre Ståhl  
on behalf of the CMS Collaboration

Laboratoire Leprince-Ringuet, École Polytechnique, France

ICHEP 2018  
39th International Conference on High Energy Physics





# W bosons and top quarks in pPb

$$W \rightarrow \mu + \nu_{\mu}$$

$$t\bar{t} \rightarrow (e \nu_e b) (\mu \nu_{\mu} b)$$


CMS Experiment at the LHC, CERN  
 Data recorded: 2016-Nov-19 01:31:00.355584 GMT  
 Run / Event / LS: 285517 / 956100501 / 652

CMS Experiment at the LHC, CERN  
 Data recorded: 2016-Nov-19 06:44:18.053352 GMT  
 Run / Event / LS: 285517 / 2067670785 / 1459



- Proton-lead asymmetric collisions
- Center-of-mass frame rapidity boost  $\approx 0.465$
- Define forward-backward ratios:

$$R_{FB} = \frac{N(\eta_{CM} > 0)}{N(\eta_{CM} < 0)} = \frac{N(p\text{-going})}{N(Pb\text{-going})}$$






# Top quarks in pPb collisions

Dominant production modes:  
**gluon  $\rightarrow t\bar{t}$**

- Top quarks decay before interacting with the medium

- Measurement of top quark and dijet production probe modifications to gluon PDF

- Top quarks constrain the high  $x$  ( $x > 0.05$ ) and high  $Q$  ( $Q^2 \sim m_{\text{top}}^2$ ) region





# W bosons in pPb collisions


Dominant production modes:

$$u\bar{d} \rightarrow W^+, d\bar{u} \rightarrow W^-$$

- Sensitive to PDFs of **valence and sea quarks**
- Quarks behave differently inside bound nucleons compared to free-nucleons  $\rightarrow$  **nuclear PDF effects.**
- W production in pPb @ LHC probe the quark nPDFs in  $10^{-3} < x < 10^{-1}$  @ **high  $Q^2$**





W in pPb @ 8.16 TeV



# W bosons in pPb collisions

- Measurement of **W production @ 5.02 TeV** by CMS helped to improve nPDF sets (CMS data included in EPPS16) and provided a hint of nuclear modifications with poor significance




Phys. Lett. B750 (2015)

- Small deviation observed in **charge asymmetry @ backward region** : possible different modification of up and down quark PDFs in the nucleus?

- W boson yield increased by ~10x** in **2016** data w.r.t. 2013, due to increase of **~2x** in **xsec** and **~5x** in **lumi**

Year	$\sqrt{s_{NN}}$	Luminosity (CMS)
2013	5.02 TeV	35 nb <sup>-1</sup>
2016	8.16 TeV	174 nb <sup>-1</sup>

# Outline



- **Top quark production in pPb at 8.16 TeV**

- Phys. Rev. Lett. 119 (2017), 242001



- **W boson production in pPb at 8.16 TeV**

- CMS-PAS-HIN-17-007



# Top quarks in pPb at 8.16 TeV

# Top quarks in pPb

Phys. Rev. Lett. 119 (2017), 242001



## 0 b-tagged jets




## 1 b-tagged jet

## >=2 b-tagged jets


pPb ( $174 \text{ nb}^{-1}$ ,  $\sqrt{s_{\text{NN}}} = 8.16 \text{ TeV}$ )

pPb ( $174 \text{ nb}^{-1}$ ,  $\sqrt{s_{\text{NN}}} = 8.16 \text{ TeV}$ )

pPb ( $174 \text{ nb}^{-1}$ ,  $\sqrt{s_{\text{NN}}} = 8.16 \text{ TeV}$ )





- **Decay channel:**  $\geq 4$  jets + lepton ( $\mu$  or  $e$ ) + missing momentum
- **Jet Selection:** Anti- $k_T$  ( $\Delta R=0.4$ ) jet with  $p_T > 25 \text{ GeV}/c$  &  $|\eta| < 2.5$
- **B-quark tagger:** Based on combined secondary vertex
- **Lepton Selection:** Isolated lepton with  $p_T > 30 \text{ GeV}/c$  &  $|\eta| < 2.1$
- **Extraction:** Fits of the  $W \rightarrow jj'$  mass using functional forms in different b-jet and lepton flavor categories, **without relying on simulation**





# Top quarks: cross section

Phys. Rev. Lett. 119 (2017), 242001



- **First observation ( $>5\sigma$ ) of top quark production in pPb collisions!**
- Inclusive cross section ( $45 \pm 8 \text{ nb}$ ) in agreement with NNLO+NNLL pQCD interfaced to NLO proton/nuclear PDF calculations


# W bosons in pPb at 8.16 TeV

# W bosons in pPb


CMS-PAS-HIN-17-007



- **Decay channel:** muon + missing momentum




- Objects reconstructed with the **Particle Flow (PF) algorithm**
- **Muon selection:** Leading isolated muon with  $p_T > 25$  GeV/c and  $|\eta| < 2.4$
- **Muon isolation:** Sum of  $p_T$  of PF particles ( $\gamma$ ,  $h^\pm$  &  $h^0$ ) around the muon  $< 15\%$  of muon  $p_T$  (**suppress multi-jet bkg**)
- **Event selection:**
  - **Veto  $Z \rightarrow \mu^- \mu^+$ :** Reject events with  $\mu^- \mu^+$  pairs,  $p_T^{\mu^-} > 15$  &  $p_T^{\mu^+} > 15$  GeV/c
- **Dominant backgrounds:** QCD jet  $\rightarrow$  muon passing isolation, and  $Z \rightarrow$  muon + (missing muon)



# W bosons in pPb

CMS-PAS-HIN-17-007





- **Simulation:** pPb NLO POWHEG v2 including CT14+EPPS16 nPDF
- **Signal and electroweak background:** Template from **simulation**
- **QCD multi-jet background:** **Data-driven** functional form
- **Extraction:** Fits of the **missing  $p_T$**  distribution in 24 bins of muon  $\eta_{CM}$

# W boson: cross section

CMS-PAS-HIN-17-007



$P(\chi^2) = <0.01\%$  CT14 , 79% nCTEQ15 , 96% EPPS16





- $\eta_{CM} < 0$  (large  $x_{pb}$ ): Results agree with PDF and nPDF calculations
- $\eta_{CM} > 0$  (small  $x_{pb}$ ): Results favor the nuclear PDF calculations

# W boson: forward-backward ratio

CMS-PAS-HIN-17-007



$P(\chi^2) = <0.01\%$  CT14 , 83% nCTEQ15 , 95% EPPS16



- Uncertainties fully correlated in  $\eta_{CM}$  cancels (correlations included)
- **Exclude ( $>7\sigma$ ) free-nucleon PDF calculations**
- Experimental uncertainties smaller than nPDF uncertainties

# W boson: charge asymmetry


CMS-PAS-HIN-17-007



$P(\chi^2) = 54\% \text{ CT14}, 23\% \text{ nCTEQ15}, 80\% \text{ EPPS16}$

pPb 173.4 nb<sup>-1</sup>


$\sqrt{s_{NN}} = 8.16 \text{ TeV}$



- nPDF effects cancel, except for different up / down quark modification
- **All (n)PDF calculations reproduce the measurements**

# W boson: PDF comparison

CMS-PAS-HIN-17-007




- Compute  $\chi^2$  test between the measurements and each individual PDF set
- Good agreement between data and EPPS16 nPDF





# Summary

## Probing gluon nPDF:

- First observation of top quark production in pPb . Opens a new window of opportunities



## Probing quark nPDF:



- Observation of nuclear modifications of the (anti) quark PDFs
- Experimental uncertainties smaller than nPDF uncertainties
- Good agreement with EPPS16 nPDF calculations


**Thank you for your attention!**



# BACKUP



# CMS detector



# W boson: forward-backward ratio




CMS-PAS-HIN-17-007

$P(\chi^2) = <0.01\%$  CT14 , 90% nCTEQ15 , 99% EPPS16

pPb 173.4 nb<sup>-1</sup>

$\sqrt{s_{NN}} = 8.16$  TeV




- **Strongly deviate from CT14 calculations favoring EPPS16**
- Experimental uncertainties significantly smaller than nPDF uncertainties

# W boson: charge asymmetry



CMS-PAS-HIN-17-007




- Good agreement between measurement at 8.16 TeV and 5.02 TeV after shifting the  $\eta_{\text{CM}}^{\mu}$  taking into account the difference in energy

# W boson: correlation matrix



CMS-PAS-HIN-17-007



- W boson cross section measurement almost uncorrelated in muon pseudorapidity, while a bit correlated in muon charge