

Exclusive Photoproduction of $2\pi^+2\pi^-$ Final State at HERA

Friday, July 6, 2018 8:15 PM (15 minutes)

Exclusive production of four charged pions at the ep collider HERA is studied at small photon virtualities $Q^2 < 2 \text{ GeV}^2$. The data were taken with the H1 detector in the years 2006 and 2007 at a centre-of-mass energy of $\sqrt{s} = 319 \text{ GeV}$ and correspond to an integrated luminosity of 7.6 pb^{-1} . The cross section of the reaction $\gamma p \rightarrow 2(\pi^+\pi^-)Y$ is determined in the phase space of $45 < W_{\gamma p} < 100 \text{ GeV}$, $|t| < 1 \text{ GeV}^2$ and $M_Y < 1.6 \text{ GeV}$. The 4π mass spectra indicate that the reaction proceeds predominately via production and decay of $\rho'(1450)$ and $\rho''(1700)$ resonances. Parameters of these resonances as well as production cross sections times branching ratio into four charged pions are estimated from the mass fit, which includes contributions from non-resonant 4π channel and interference terms.

Primary authors: SCHMITT, Stefan (Deutsches Elektronen-Synchrotron (DE)); COLLABORATION, H1 (DESY); BOLZ, Arthur (Ruprecht Karls Universitaet Heidelberg (DE))

Presenter: BOLZ, Arthur (Ruprecht Karls Universitaet Heidelberg (DE))

Session Classification: POSTER

Track Classification: Strong Interactions and Hadron Physics