## A Minimal Model For Two-Component FIMP Dark Matter: A Basic Search

Friday 6 July 2018 20:15 (15 minutes)

In the multi-component configurations of dark matter phenomenology, we propose a minimal two-component one which is an extension of the Standard Model with only three new fields; one scalar and one fermion interact with the thermal soup through Higgs portal mediated by the other scalar in such a way that the stabilities of dark matter candidates are made simultaneously by an explicit Z2 symmetry. Against the most common freeze-out framework, we look for dark matter particle signatures in the freeze-in scenario by evaluating the relic density and detection signals. A simple distinguishing feature of the model is the lack of dark matter conversion, so the dark matter components act individually and the model can be adapted entirely to both singlet scalar and singlet fermionic models, separately. We find dark matter self-interaction as the most promising approach to probe such feeble models. Although, the scalar component adopts this constraint, the

fermionic one refuses it even in the resonant region.

**Authors:** Mr ZAKERI, Seyed Peyman (F aculty of P hysics, Y azd University, P.O. Box 89195 – 741, Y azd, Iran); Dr MOOSAVI NEJAD, Seyed Mohammad (School of P articles and Accelerators, Institute for Research in Fundamental Sciences (IPM), P.O. Box 19395 – 5531, T ehran, Iran); Dr ZAKERI, Mohammadreza (Physics and Astronomy Department, University of California, Riverside, California 92521, USA); Dr AYAZI, Seyed Yaser (Physics Department, Semnan University, P.O. Box 35131 – 19111, Semnan, Iran)

**Presenter:** Mr ZAKERI, Seyed Peyman (F aculty of P hysics, Y azd University, P.O. Box 89195 – 741, Y azd, Iran)

Session Classification: POSTER

Track Classification: Beyond the Standard Model