

P. Giacomelli INFN Bologna





• FCC complex and FCC-ee first stage



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement
- Higgs couplings



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement
- Higgs couplings
- s-channel Higgs production



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement
- Higgs couplings
- s-channel Higgs production
- BSM Higgs searches



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement
- Higgs couplings
- s-channel Higgs production
- BSM Higgs searches
- Higgs self-coupling



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement
- Higgs couplings
- s-channel Higgs production
- BSM Higgs searches
- Higgs self-coupling
- Synergy with FCC-hh



- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement
- Higgs couplings
- s-channel Higgs production
- BSM Higgs searches
- Higgs self-coupling
- Synergy with FCC-hh
- Conclusions




- FCC complex and FCC-ee first stage
- Higgs production at FCC-ee
- Higgs width measurement
- Higgs couplings
- s-channel Higgs production
- BSM Higgs searches
- Higgs self-coupling
- Synergy with FCC-hh
- Conclusions

Acknowledgments

M. Klute and P. Janot for material and useful discussions



FCC complex



International FCC collaboration


- pp-collider (FCC-hh)
- main emphasis, defining infrastructure requirements
- 16 T \Rightarrow 100 TeV pp, L~2x10³⁵, L_{int}=2 ab⁻¹/yr
- ~100 km tunnel infrastructure
- ^y e⁺e⁻ collider (FCC-ee),√s=90-365 GeV, L~10³⁵ 4x10³⁶, L_{int}=1-48 ab⁻¹/yr for H, Z as potential first step
- FCC-hh technology
- **p-e** (*FCC-he*) option, √s=3.5 TeV, L~10³⁴

FCC-ee operation mod	el

working point	nominal luminosity/IP [10 ³⁴ cm ⁻² s ⁻¹]	total luminosity (2 IPs)/ vr half luminosity in first two years (Z) and first year (ttbar) to account for initial operation	physics goal	run time [years]
Z first 2 years	100	26 ab ⁻¹ /year	150 - L-1	A
Z later	200	48 ab ⁻¹ /year	150 ab ⁻¹	4
W	25	6 ab ⁻¹ /year	10 ab ⁻¹	1 - 2
Н	7.0	1.7 ab ⁻¹ /year	5 ab ⁻¹	3
machine modification for	RF installation &	rearrangement: 1 year		
top 1st year (350 GeV)	0.8	0.2 ab ⁻¹ /year	0.2 ab ⁻¹	1
top later (365 GeV)	1.4	0.34 ab ⁻¹ /year	1.5 ab ⁻¹	4



Future lepton colliders luminosities




Significant advantage in luminosity for circular colliders vs. linear colliders, especially for FCC-ee. Linear colliders (esp. CLIC) have higher energy reach, but less than FCC-hh.




Higgs production at FCC-ee

FCC-ee
5 ab-1@240 GeV
~1.5 ab-1@365 GeV



Higgs Factory!





	FCC-ee 240 GeV	FCC-ee 365 GeV
Total Integrated Luminosity (ab ⁻¹)	5	1.5
# Higgs bosons from e+e-→HZ	1,000,000	180,000
# Higgs bosons from fusion process	25,000	45,000



Higgs couplings to Z

- → Recoil method provides a unique opportunity for a decay-mode independent measurement of the HZ coupling
 - Figgs events are tagged with the Z boson decays, independently of Higgs decay mode, m_{recoil} = m_H
 - Expected precision 0.5% at 240 GeV, 0.9% at 365 GeV on the ZH cross section
 - Using only leptonic Z decays







Higgs boson width

- Total Higgs boson width can be extracted from a combination of measurements in a model independent way
 - 1) tagging Higgs final states

$$\sigma(\text{ee} \to \text{ZH}) \cdot \text{BR}(\text{H} \to \text{ZZ}) \propto \frac{g_{\text{HZ}}^4}{\Gamma}$$

2) measurements of vector boson fusion production at 350-365 GeV

$$\frac{\sigma(\text{ee} \rightarrow \text{ZH}) \cdot \text{BR}(\text{H} \rightarrow \text{WW}) \cdot \sigma(\text{ee} \rightarrow \text{ZH}) \cdot \text{BR}(\text{H} \rightarrow \text{bb})}{\sigma(\text{ee} \rightarrow \nu \nu \text{H}) \cdot \text{BR}(\text{H} \rightarrow \text{bb})}$$

$$\propto \frac{g_{\text{HZ}}^2 \cdot g_{\text{HW}}^2}{\Gamma} \cdot \frac{g_{\text{HZ}}^2 \cdot g_{\text{Hb}}^2}{\Gamma} \cdot \frac{\Gamma}{g_{\text{HW}}^2 \cdot g_{\text{Hb}}^2} = \frac{g_{\text{HZ}}^4}{\Gamma}$$

3) combination of all measurements

Precision obtainable on $\delta\Gamma_H/\Gamma_H \sim 1.6\%$



Higgs boson couplings

→ Precision Higgs coupling measurements

- Absolute coupling measurements enabled by HZ cross section and total width measurement
- Data at 365 GeV constrain total width
- Fragging individual Higgs final states to extract various Higgs couplings
- Couplings extracted from model-independent fit
- Statistical uncertainties are shown for 5 ab-1@240 GeV and 1.5 ab-1@365 GeV (from FCC-ee CDR)
 - all measurements are under review / are being redone

Several couplings ($g_{H\mu\mu}$, $g_{H\gamma\gamma}$, g_{Htt}) improve further by doing a combined fit with HL-LHC.


The combined fit makes the HL-LHC measurements model-independent.

in %	FCC-ee 240 GeV	+FCC-ee 365 GeV	+HL- LHC
δ g Hzz	0.25	0.22	0.21
δ g нww	1.3	0.47	0.44
δ G Hbb	1.4	0.68	0.58
δ g Hcc	1.8	1.23	1.20
δ g Hgg	1.7	1.03	0.83
$\delta g_{ extsf{H}_{ au au}}$	1.4	0.8	0.71
δ g нμμ	9.6	8.6	3.4
$\delta \mathbf{g}_{H\gamma\gamma}$	4.7	3.8	1.3
δ g Htt			3.3
δΓΗ	2.8	1.56	1.3

relative meas., e.g. $\delta g_H/g_H$



Higgs boson couplings, ZH→{\lambda}X




CLD: CLIC-inspired Lepton Detector, optimised for FCC-ee

C. Bernet FCC week 2018




Higgs boson couplings, ZH→llbb




C. Bernet FCC week 2018



Higgs boson couplings, ZH→qqbb



Using CMS detector



Using CLD detector


C. Bernet FCC week 2018




electron Yukawa coupling

⇒ s-channel Higgs production

- unique opportunity for measurement close to SM sensitivity
- highly challenging; $\sigma(ee \rightarrow H) = 1.6$ fb;
- various Higgs decay channels studied
- studied monochromatization scenarios
 - baseline: 6 MeV energy spread, L = 2 ab⁻¹
 - optimized: 10 MeV energy spread, L = 7 ab⁻¹
 - limit ~2.5 times SM in both cases








Higgs invisible decays

→ Higgs boson to invisible decays are predicted for instance in the Higgs-portal model of Dark Matter.




- Follows FCC-ee ZH cross section measurement, events with a Z and nothing else are selected
- for visualization BR(H->inv) = 100%
- 95%CL upper limit using 5ab-1 is 0.3% at 240
 GeV, 0.6% with 1.5ab-1 at 365 GeV
- Study using leptonic Z decays in Eur. Phys. J. C (2017) 77: 116
- Hadronic Z decays also included



BSM Higgs


→ Excellent opportunities for BSM Higgs searches







Higgs self-coupling

- → Very large HZ datasets allow $σ_{ZH}$ and $σ_{vvH}$ measurements of extreme precision at √s = 240, 350 and 365 GeV
- → Model-independent bounds obtained from global-fit on Higgs self coupling $κ_λ$ and the coupling to SM bosons c_Z








A precision on $\delta \kappa_{\lambda}$ of $\pm 40\%$ can be achieved, and of $\pm 35\%$ in combination with HL-LHC.

If c_Z if fixed to its SM value, then the precision on $\delta \kappa_\lambda$ improves to $\pm 20\%$



Higgs self-coupling at FCC-hh



→ Enormous di-Higgs samples produced at FCC-hh


- \circ σ (100 TeV) / σ (14 TeV) \simeq 40
- \bigcirc L (FCC-hh) / L (HL-LHC) \cong 10
- Naively, factor 20 smaller statistical uncertainty

→ Studied a number of final states

bbyy most sensitive channel

→ Per-cent level precision of HHH coupling

- Only possible at a 100 TeV hadron machine



$$\delta\mu \cong 2-4\%$$

$$\delta \kappa \simeq 5\%$$

Details in arXiv:1606.09408 and arXiv:1802.01607





INFN Conclusions

→ Fantastic prospects to probe the Higgs sector with FCC



- **→ Fantastic** prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities



- **→ Fantastic** prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee



- **→** Fantastic prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee
 - Precision measurements of Higgs boson couplings, including self couplings, mass, CP



- **→ Fantastic** prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee
 - Precision measurements of Higgs boson couplings, including self couplings, mass, CP
 - sub-percent precision on several Higgs couplings only possible with FCC-ee



- **→** Fantastic prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee
 - Precision measurements of Higgs boson couplings, including self couplings, mass, CP
 - sub-percent precision on several Higgs couplings only possible with FCC-ee
 - BSM Higgs physics through direct and indirect measurements



- **→** Fantastic prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee
 - Precision measurements of Higgs boson couplings, including self couplings, mass, CP
 - sub-percent precision on several Higgs couplings only possible with FCC-ee
 - BSM Higgs physics through direct and indirect measurements
 - Synergy between FCC-ee and FCC-hh Higgs physics



- **→** Fantastic prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee
 - Precision measurements of Higgs boson couplings, including self couplings, mass, CP
 - sub-percent precision on several Higgs couplings only possible with FCC-ee
 - BSM Higgs physics through direct and indirect measurements
 - Synergy between FCC-ee and FCC-hh Higgs physics
 - More precise Higgs measurements are foreseen for the FCC CDR.



- **→** Fantastic prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee
 - Precision measurements of Higgs boson couplings, including self couplings, mass, CP
 - sub-percent precision on several Higgs couplings only possible with FCC-ee
 - BSM Higgs physics through direct and indirect measurements
 - Synergy between FCC-ee and FCC-hh Higgs physics
 - More precise Higgs measurements are foreseen for the FCC CDR.
 - FCC-ee and FCC-hh will provide <u>by far</u> the best possible Higgs measurements of any accelerator.



- **→** Fantastic prospects to probe the Higgs sector with FCC
 - Large Higgs samples open new possibilities
 - Model-independent measurements of gzH and total width with FCC-ee
 - Precision measurements of Higgs boson couplings, including self couplings, mass, CP
 - sub-percent precision on several Higgs couplings only possible with FCC-ee
 - BSM Higgs physics through direct and indirect measurements
 - Synergy between FCC-ee and FCC-hh Higgs physics
 - More precise Higgs measurements are foreseen for the FCC CDR.
 - FCC-ee and FCC-hh will provide <u>by far</u> the best possible Higgs measurements of any accelerator.

Stay tuned!

Backup



FCC-ee parameters

parameter	Z	ww	H (ZH)	ttbar
beam energy [GeV]	45	80	120	182.5
beam current [mA]	1390	147	29	5.4
no. bunches/beam	16640	2000	393	48
bunch intensity [10 ¹¹]	1.7	1.5	1.5	2.3
SR energy loss / turn [GeV]	0.036	0.34	1.72	9.21
total RF voltage [GV]	0.1	0.44	2.0	10.9
long. damping time [turns]	1281	235	70	20
horizontal beta* [m]	0.15	0.2	0.3	1
vertical beta* [mm]	0.8	1	1	1.6
horiz. geometric emittance [nm]	0.27	0.28	0.63	1.46
vert. geom. emittance [pm]	1.0	1.7	1.3	2.9
bunch length with SR / BS [mm]	3.5 / 12.1	3.0 / 6.0	3.3 / 5.3	2.0 / 2.5
luminosity per IP [10 ³⁴ cm ⁻² s ⁻¹]	>200	>25	>7	>1.4
beam lifetime rad Bhabha / BS [min]	68 / >200	49 / >1000	38 / 18	40 / 18




Higgs CP studies

- → H→ττ decay is a promising channel to study
 CP violation
 - Free level couplings to quarks and leptons
 - CP-even and CP-odd couplings induced at the same order
- → CP violation can be probed through τ polarization
 - τ decays clean enough that the spin information is not washed out by hadronization effects

 - $\mathcal{L}_{hff} \propto h \bar{f} (\cos \Delta + \mathrm{i} \gamma_5 \sin \Delta) f$
 - model using effective lagrangian

Andres Rios (MIT), Aram Apyan (FNAL) following arXiv:1308.1094




- 920 signal event in 5 ab⁻¹
- expected 68% CL
 - 0.17 radian (0.05 in GEN level study)
 - 9.7 degree (2.9 in GEN level study)



Higgs self-coupling at hadron colliders

→ Probing triple-Higgs coupling with double Higgs production

- Consistency of check of EWSB
- Reconstructing the Higgs potential
- Sensitivity through yields and kinematics
- Large enhancement through BSM possible
- Exhaustive program at the (HL-)LHC



EFT Lagrangian