

Hadronic Decays of $D^{0(+)}$ and D_s^+ at BESIII

Liaoyuan Dong

Institute of High Energy Physics, Beijing

(On behalf of the BESIII collaboration)



ICHEP2018 SEOUL

XXXIX INTERNATIONAL CONFERENCE ON

JULY 4 - 11, 2018 COEX, SEOUL

high energy PHYSICS

Outline

- Introduction

- The BESIII experiment
- Production near threshold and tag technique

- Hadronic decays of $D_{(s)}$

- Observation of pure W -annihilation decays:

$$D_s^+ \rightarrow p\bar{n}, D_s^+ \rightarrow \omega\pi^+, D_s^+ \rightarrow a_0(980)\pi$$

- Amplitude analysis of $D \rightarrow K\pi\pi\pi$:

$$D^0 \rightarrow K^-\pi^+\pi^+\pi^-, D^0 \rightarrow K^-\pi^+\pi^0\pi^0, D^+ \rightarrow K_s^0\pi^+\pi^+\pi^-$$

- Measurement of the branching fractions(BFs) of

$$D \rightarrow PP \quad (P=\text{pseudo-scalar})$$


- Summary

The BESIII Experiment

Beijing-Electron-Positron Collider II (BEPCII)


- e^+e^- collisions with
 $\sqrt{s} = 2.0 - 4.6 \text{ GeV}$
- Direct production of charmonia
- Designed Luminosity

$\mathcal{L} = 1 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
was achieved in April 2016.



BESIII detector

- 93% coverage of the full solid angle
- Main drift chamber $\sigma_p/p = 0.5\% @ 1 \text{ GeV}$
- Time-of-flight system $\sigma_T = 100 \text{ ps}$ in Barrel
- Eimg. Calorimeter $\Delta E/E = 2.5\% @ 1 \text{ GeV}$
- Superconducting 1T magnet
- Muon system (RPC)



Production near threshold and tag technique

Dataset used in this talk:


- 2.93 fb^{-1} at $E_{\text{cm}} = 3.773 \text{ GeV}$ ($\sim 3.6\times$ larger than CLEO's):
 D are produced via $e^+e^- \rightarrow \psi(3770) \rightarrow D\bar{D}$
- 3.19 fb^{-1} at $E_{\text{cm}} = 4.178 \text{ GeV}$ ($\sim 5.3\times$ larger than CLEO's):
 D_s are produced mostly via $e^+e^- \rightarrow D_s^\pm D_s^{*\mp}, D_s^{*\mp} \rightarrow \pi^0/\gamma D_s^\mp$

Two ways to study $D_{(s)}$ decays:

- **Single Tag (ST):** reconstruct only one of the $D\bar{D}$ ($D_s^+ D_s^-$)
- **Double Tag (DT):** reconstruct both of $D\bar{D}$ ($D_s^+ D_s^-$)

DT provides access to absolute BF's.

DT provides clean samples for amplitude analysis.



(Charge-conjugate states are implied throughout this talk)

Observation of pure W -annihilation decay

$$D_s^+ \rightarrow p\bar{n}$$

- $D_s^+ \rightarrow p\bar{n}$ is the only baryonic decay of charmed meson and can proceed only through W -annihilation process,
 - Short-distance expected: $\text{BF} \sim 10^{-6}$
 - Long-distance enhance to: $\text{BF} \sim 10^{-3}$

PLB663(2008)326

The large BF ($\sim 10^{-3}$) indicates large final state interaction(FSI) effect and is important to understand the dynamical enhancement of W -annihilation.

- First evidence was reported by CLEO with a signal of 13.0 ± 3.6 events with $\text{BF} = (1.30 \pm 0.36_{-0.16}^{+0.12}) \times 10^{-3}$ (PRL100, 181802(2008)).


Observation of $D_s^+ \rightarrow p\bar{n}$

Tag modes:

$K_S^0 K^-$, $K^- K^+ \pi^-$, $\bar{K}_S^0 K^- \pi^0$, $K^- K^+ \pi^- \pi^0$,
 $K_S^0 K^+ \pi^- \pi^-$, $\pi^- \pi^+ \pi^-$, $\pi^- \eta$, $\rho^- \eta$, $\pi^- \eta'$ (with $\eta' \rightarrow$
 $\pi^+ \pi^- \eta$), $\pi^- \eta'$ (with $\eta' \rightarrow \gamma \pi^+ \pi^-$) and $K^- \pi^+ \pi^-$.

Total ST yield $\sim 0.35\text{M}$

Fit to the mass of missing particle
 M_{miss} to get the DT yield.



Preliminary result:

$$\mathcal{B}_{D_s^+ \rightarrow p\bar{n}} = (1.22 \pm 0.10) \times 10^{-3}$$

Statistical uncertainty only,

Statistically limited. Sys. dominated by baryon PID.

- Confirm CLEO's measurement with greatly improved precision.
- Consistent with “long-distance” expectation (PLB663, 326).

Observation of pure W -annihilation decays

$$D_s^+ \rightarrow \omega \pi^+, D_s^+ \rightarrow a_0(980) \pi$$

- $D_s^+ \rightarrow \omega \pi^+, D_s^+ \rightarrow a_0(980) \pi$ can proceed only via W -annihilation process:
 - factorizable short-distance contribution is helicity suppressed,
 - non-factorizable long-distance contribution induced by FSI dominate,which makes the input from experimental measurement to be the unique method to determine the W -annihilation amplitude.

- With the measured BF of $D_s^+ \rightarrow \omega \pi^+$ as one of the inputs ,
Q. Qin et al. (PRD89, 054006) predicts:

$$\mathcal{B}(D_s^+ \rightarrow \omega K^+) = 0.6 \times 10^{-3}, A_{CP}(D_s^+ \rightarrow \omega K^+) = -0.6 \times 10^{-3} \text{ (without } \rho - \omega \text{ mixing)}$$

$$\mathcal{B}(D_s^+ \rightarrow \omega K^+) = 0.07 \times 10^{-3}, A_{CP}(D_s^+ \rightarrow \omega K^+) = -2.3 \times 10^{-3} \text{ (with } \rho - \omega \text{ mixing)}$$


Among the largest expected A_{CP} observed in charmed decays

- $D_s^+ \rightarrow \omega \pi^+$: Evidence by CLEO, $\text{BF} = (2.1 \pm 0.9 \pm 0.1) \times 10^{-3}$ with a signal of 6.0 ± 2.4 events.
- $D_s^+ \rightarrow \omega K^+$: CLEO set an UL = 2.4×10^{-3} @90% C.L. (PRD80, 051102(R) (2009))

Observation of $D_s^+ \rightarrow \omega \pi^+$ and $D_s^+ \rightarrow \omega K^+$

- Tag modes: $D_s^- \rightarrow K_S^0 K^-$, $D_s^- \rightarrow K^+ K^- \pi^-$. Total ST yield $\sim 0.167\text{M}$.
- Double tag: average mass of two D_s mesons closest to the PDG value.

Fit to the invariant mass $M_{\pi^+ \pi^- \pi^0}$ to get the DT yield:



Preliminary results:

Consistent with CLEO's measurement, but more precise.

$$\mathcal{B}(D_s^+ \rightarrow \omega \pi^+) = (1.85 \pm 0.30_{stat.} \pm 0.19_{sys.}) \times 10^{-3}$$

$$\mathcal{B}(D_s^+ \rightarrow \omega K^+) = (1.13 \pm 0.24_{stat.} \pm 0.14_{sys.}) \times 10^{-3}$$

First observation !

- The measurement of $D_s^+ \rightarrow \omega K^+$ implies the $\rho - \omega$ mixing is negligible.

Amplitude analysis of $D_s^+ \rightarrow \pi^+ \pi^0 \eta$

Events are selected with double tag:

- Tag modes:

$$D_s^- \rightarrow K_S^0 K^-, D_s^- \rightarrow K^+ K^- \pi^-, D_s^- \rightarrow K_S^0 K^- \pi^0, D_s^- \rightarrow K^+ K^- \pi^- \pi^0, \\ D_s^- \rightarrow K_S^0 K^+ \pi^- \pi^-, D_s^- \rightarrow \pi^- \eta_{\gamma\gamma}, D_s^- \rightarrow \pi^- \eta'_{\pi^+ \pi^- \eta}$$

Data sample for amplitude analysis:

- A Multi-variate analysis is performed to suppress the background from fake η .
- The retained data sample has 1239 events with a purity of $(97.7 \pm 0.5)\%$.


Observation of $D_s^+ \rightarrow a_0(980)\pi$

Preliminary: significances, phases, and fit fractions (FFs) for intermediate processes:

Amplitude	Significance (σ)	Phase	FF
$D_s^+ \rightarrow \rho^+\eta$	> 20	0.0 (fixed)	$0.783 \pm 0.050 \pm 0.021$
$D_s^+ \rightarrow (\pi^+\pi^0)_V\eta$	5.7	$0.612 \pm 0.172 \pm 0.342$	$0.054 \pm 0.021 \pm 0.026$
$D_s^+ \rightarrow a_0(980)\pi$	16.2	$2.794 \pm 0.087 \pm 0.041$	$0.232 \pm 0.023 \pm 0.034$

The amplitudes agree with: $A(D_s^+ \rightarrow a_0(980)^+\pi^0) = -A(D_s^+ \rightarrow a_0(980)^0\pi^+)$ within stat. uncertainty, thus we set the magnitudes to be the same with the phase difference fixed to π .

Dalitz plot & Projections:



Dots: data; solid: total fit; dashed: $D_s^+ \rightarrow \rho^+\eta$; long dashed: $D_s^+ \rightarrow a_0(980)\pi$ 10

Branching Fraction Results of $D_s^+ \rightarrow \pi^+ \pi^0 \eta$

Fit to the invariant mass $M_{\pi^+ \pi^0 \eta}$ to get the DT yield.

Preliminary results:

$$\mathcal{B}(D_s^+ \rightarrow \pi^+ \pi^0 \eta) = (9.50 \pm 0.28_{stat.} \pm 0.41_{sys.})\%$$

PDG value = $(9.2 \pm 1.2)\%$

$$BF(\text{sub-mode } n) = \mathcal{B}(D_s^+ \rightarrow \pi^+ \pi^0 \eta) FF(n)$$


Branching fraction (%)	BESIII Preliminary
$\mathcal{B}(D_s^+ \rightarrow \rho^+ \eta) = 7.44 \pm 0.48_{stat.} \pm 0.44_{sys.}$	
$\mathcal{B}(D_s^+ \rightarrow a_0(980)\pi)^* = 2.20 \pm 0.22_{stat.} \pm 0.34_{sys.}$	
$\mathcal{B}(D_s^+ \rightarrow a_0(980)^+ \pi^0)^* = 1.46 \pm 0.15_{stat.} \pm 0.22_{sys.}$	
$\mathcal{B}(D_s^+ \rightarrow a_0(980)^0 \pi^+)^* = 1.46 \pm 0.15_{stat.} \pm 0.22_{sys.}$	

*here, $a_0(980) \rightarrow \pi\eta$

Sys. dominated by π^0 and η reconstruction (4%).

- The measured $\mathcal{B}(D_s^+ \rightarrow a_0(980)^+ \pi^0)$ is larger than other measured pure W -annihilation decays ($D_s^+ \rightarrow p\bar{n}, D_s^+ \rightarrow \omega\pi^+$) by one order.

This provides theoretical challenge in understanding such a large W -annihilation contribution in $D \rightarrow SP$ (S =scalar; P =pseudo-scalar).



PDG value = $(8.9 \pm 0.9)\%$

First observation !

Amplitude analysis of $D \rightarrow K\pi\pi\pi$

- The measurement of the sub-modes in $D \rightarrow K\pi\pi\pi$ provides a window to study the decays $D \rightarrow AP$ and $D \rightarrow VV$ (A =axial-vector, V =vector), both of them are important in learning the CPV in charm decays but less effective experimental measurements.
- The knowledge of sub-modes can be widely used in many measurements:
 - Branching fraction measurement
 - Strong phase measurement
 - CKM unitary triangle measurement
- There are seven $D \rightarrow K\pi\pi\pi$ modes:
 $D^0 \rightarrow K^-\pi^+\pi^+\pi^-, K^-\pi^+\pi^0\pi^0, K_S^0\pi^+\pi^-\pi^0, K_S^0\pi^0\pi^0\pi^0$ and $D^+ \rightarrow K^-\pi^+\pi^+\pi^0, K_S^0\pi^+\pi^+\pi^-, K_S^0\pi^+\pi^0\pi^0$.

Previous measurements of sub-modes in $D^0 \rightarrow K^-\pi^+\pi^+\pi^-, K_S^0\pi^+\pi^-\pi^0$ and $D^+ \rightarrow K^-\pi^+\pi^+\pi^0, K_S^0\pi^+\pi^+\pi^-$ have been performed by Mark III and E691.
Both measurements are affected by low statistics.
- In this talk, we report the amplitude analysis results of
 $D^0 \rightarrow K^-\pi^+\pi^+\pi^-, D^0 \rightarrow K^-\pi^+\pi^0\pi^0, D^+ \rightarrow K_S^0\pi^+\pi^+\pi^-$

Amplitude Analysis Results of $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$

Double tag: The D^0 is reconstructed by $K^- \pi^+ \pi^+ \pi^-$ with \bar{D}^0 reconstructed by $K^+ \pi^-$. PRD95,072010

A sample of 15912 events with purity ~99.4% is used to perform the amplitude analysis.


The data can be described with 23 amplitudes:

Amplitude	ϕ_i	Fit fraction (%)
$D^0[S] \rightarrow \bar{K}^* \rho^0$	$2.35 \pm 0.06 \pm 0.18$	$6.5 \pm 0.5 \pm 0.8$
$D^0[P] \rightarrow \bar{K}^* \rho^0$	$-2.25 \pm 0.08 \pm 0.15$	$2.3 \pm 0.2 \pm 0.1$
$D^0[D] \rightarrow \bar{K}^* \rho^0$	$2.49 \pm 0.06 \pm 0.11$	$7.9 \pm 0.4 \pm 0.7$
$D^0 \rightarrow K^- a_1^+(1260), a_1^+(1260)[S] \rightarrow \rho^0 \pi^+$	0(fixed)	$53.2 \pm 2.8 \pm 4.0$
$D^0 \rightarrow K^- a_1^+(1260), a_1^+(1260)[D] \rightarrow \rho^0 \pi^+$	$-2.11 \pm 0.15 \pm 0.21$	$0.3 \pm 0.1 \pm 0.1$
$D^0 \rightarrow K_1^-(1270) \pi^+, K_1^-(1270)[S] \rightarrow \bar{K}^{*0} \pi^-$	$1.48 \pm 0.21 \pm 0.24$	$0.1 \pm 0.1 \pm 0.1$
$D^0 \rightarrow K_1^-(1270) \pi^+, K_1^-(1270)[D] \rightarrow \bar{K}^{*0} \pi^-$	$3.00 \pm 0.09 \pm 0.15$	$0.7 \pm 0.2 \pm 0.2$
$D^0 \rightarrow K_1^-(1270) \pi^+, K_1^-(1270) \rightarrow K^- \rho^0$	$-2.46 \pm 0.06 \pm 0.21$	$3.4 \pm 0.3 \pm 0.5$
$D^0 \rightarrow (\rho^0 K^-)_A \pi^+, (\rho^0 K^-)_A [D] \rightarrow K^- \rho^0$	$-0.43 \pm 0.09 \pm 0.12$	$1.1 \pm 0.2 \pm 0.3$
$D^0 \rightarrow (K^- \rho^0)_P \pi^+$	$-0.14 \pm 0.11 \pm 0.10$	$7.4 \pm 1.6 \pm 5.7$
$D^0 \rightarrow (K^- \pi^+)_{S\text{-wave}} \rho^0$	$-2.45 \pm 0.19 \pm 0.47$	$2.0 \pm 0.7 \pm 1.9$
$D^0 \rightarrow (K^- \rho^0)_V \pi^+$	$-1.34 \pm 0.12 \pm 0.09$	$0.4 \pm 0.1 \pm 0.1$
$D^0 \rightarrow (\bar{K}^{*0} \pi^-)_P \pi^+$	$-2.09 \pm 0.12 \pm 0.22$	$2.4 \pm 0.5 \pm 0.5$
$D^0 \rightarrow \bar{K}^{*0} (\pi^+ \pi^-)_S$	$-0.17 \pm 0.11 \pm 0.12$	$2.6 \pm 0.6 \pm 0.6$
$D^0 \rightarrow (\bar{K}^{*0} \pi^-)_V \pi^+$	$-2.13 \pm 0.10 \pm 0.11$	$0.8 \pm 0.1 \pm 0.1$
$D^0 \rightarrow ((K^- \pi^+)_{S\text{-wave}} \pi^-)_A \pi^+$	$-1.36 \pm 0.08 \pm 0.37$	$5.6 \pm 0.9 \pm 2.7$
$D^0 \rightarrow K^- ((\pi^+ \pi^-)_S \pi^+)_A$	$-2.23 \pm 0.08 \pm 0.22$	$13.1 \pm 1.9 \pm 2.2$
$D^0 \rightarrow (K^- \pi^+)_{S\text{-wave}} (\pi^+ \pi^-)_S$	$-1.40 \pm 0.04 \pm 0.22$	$16.3 \pm 0.5 \pm 0.6$
$D^0[S] \rightarrow (K^- \pi^+)_V (\pi^+ \pi^-)_V$	$1.59 \pm 0.13 \pm 0.41$	$5.4 \pm 1.2 \pm 1.9$
$D^0 \rightarrow (K^- \pi^+)_{S\text{-wave}} (\pi^+ \pi^-)_V$	$-0.16 \pm 0.17 \pm 0.43$	$1.9 \pm 0.6 \pm 1.2$
$D^0 \rightarrow (K^- \pi^+)_V (\pi^+ \pi^-)_S$	$2.58 \pm 0.08 \pm 0.25$	$2.9 \pm 0.5 \pm 1.7$
$D^0 \rightarrow (K^- \pi^+)_T (\pi^+ \pi^-)_S$	$-2.92 \pm 0.14 \pm 0.12$	$0.3 \pm 0.1 \pm 0.1$
$D^0 \rightarrow (K^- \pi^+)_{S\text{-wave}} (\pi^+ \pi^-)_T$	$2.45 \pm 0.12 \pm 0.37$	$0.5 \pm 0.1 \pm 0.1$

Amplitude Analysis Results of $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$

PRD95,072010

Fit projections:



Points with error bars: data, curves: fit, red histograms: background.

For the two identical π^+ , we require $m(\pi_1^+ \pi^-) > m(\pi_2^+ \pi^-)$.

Branching Fraction Results of $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$

PRD95,072010

Results of branching fractions for different components:

Component	Branching fraction (%)	PDG value (%)
$D^0 \rightarrow \bar{K}^{*0} \rho^0$	$0.99 \pm 0.04 \pm 0.04 \pm 0.03$	1.05 ± 0.23
$D^0 \rightarrow K^- a_1^+(1260) (\rho^0 \pi^+)$	$4.41 \pm 0.22 \pm 0.30 \pm 0.13$	3.6 ± 0.6
$D^0 \rightarrow K_1^-(1270) (\bar{K}^{*0} \pi^-) \pi^+$	$0.07 \pm 0.01 \pm 0.02 \pm 0.00$	0.29 ± 0.03
$D^0 \rightarrow K_1^-(1270) (K^- \rho^0) \pi^+$	$0.27 \pm 0.02 \pm 0.04 \pm 0.01$	
$D^0 \rightarrow K^- \pi^+ \rho^0$	$0.68 \pm 0.09 \pm 0.20 \pm 0.02$	0.51 ± 0.23
$D^0 \rightarrow \bar{K}^{*0} \pi^+ \pi^-$	$0.57 \pm 0.03 \pm 0.04 \pm 0.02$	0.99 ± 0.23
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	$1.77 \pm 0.05 \pm 0.04 \pm 0.05$	1.88 ± 0.26

stat. uncertainty from FF

sys. uncertainty from FF

uncertainties related to $\text{BF}(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)$ in PDG

Amplitude Analysis Results of $D^0 \rightarrow K^- \pi^+ \pi^0 \pi^0$

Double tag: The D^0 is reconstructed by $K^- \pi^+ \pi^0 \pi^0$ with \bar{D}^0 reconstructed by $K^+ \pi^-$.

A sample of 5950 events with purity ~99% is used to perform the amplitude analysis.

The data can be described with 26 amplitudes:









Amplitude mode	FF(%)	Phase (ϕ)	BESIII Preliminary
$D \rightarrow SS$			
$D \rightarrow (K^- \pi^+)_{S\text{-wave}}(\pi^0 \pi^0)_S$	$6.92 \pm 1.44 \pm 2.86$	$-0.75 \pm 0.15 \pm 0.47$	
$D \rightarrow (K^- \pi^0)_{S\text{-wave}}(\pi^+ \pi^0)_S$	$4.18 \pm 1.02 \pm 1.77$	$-2.90 \pm 0.19 \pm 0.47$	
$D \rightarrow AP, A \rightarrow VP$			
$D \rightarrow K^- a_1(1260)^+, \rho^+ \pi^0[S]$	$28.36 \pm 2.50 \pm 3.53$	0 (fixed)	
$D \rightarrow K^- a_1(1260)^+, \rho^+ \pi^0[D]$	$0.68 \pm 0.29 \pm 0.30$	$-2.05 \pm 0.17 \pm 0.25$	
$D \rightarrow K_1(1270)^- \pi^+, K^{*-} \pi^0[S]$	$0.15 \pm 0.09 \pm 0.18$	$1.84 \pm 0.34 \pm 0.43$	
$D \rightarrow K_1(1270)^0 \pi^0, K^{*0} \pi^0[S]$	$0.39 \pm 0.18 \pm 0.30$	$-1.55 \pm 0.20 \pm 0.26$	
$D \rightarrow K_1(1270)^0 \pi^0, K^{*0} \pi^0[D]$	$0.11 \pm 0.11 \pm 0.13$	$-1.35 \pm 0.43 \pm 0.48$	
$D \rightarrow K_1(1270)^0 \pi^0, K^- \rho^+[S]$	$2.71 \pm 0.38 \pm 0.29$	$-2.07 \pm 0.09 \pm 0.20$	
$D \rightarrow (K^{*-} \pi^0)_A \pi^+, K^{*-} \pi^0[S]$	$1.85 \pm 0.62 \pm 1.11$	$1.93 \pm 0.10 \pm 0.15$	
$D \rightarrow (K^{*0} \pi^0)_A \pi^0, K^{*0} \pi^0[S]$	$3.13 \pm 0.45 \pm 0.58$	$0.44 \pm 0.12 \pm 0.21$	
$D \rightarrow (K^{*0} \pi^0)_A \pi^0, K^{*0} \pi^0[D]$	$0.46 \pm 0.17 \pm 0.29$	$-1.84 \pm 0.26 \pm 0.42$	
$D \rightarrow (\rho^+ K^-)_A \pi^0, K^- \rho^+[D]$	$0.75 \pm 0.40 \pm 0.60$	$0.64 \pm 0.36 \pm 0.53$	
$D \rightarrow AP, A \rightarrow SP$			
$D \rightarrow ((K^- \pi^+)_{S\text{-wave}} \pi^0)_A \pi^0$	$1.99 \pm 1.08 \pm 1.55$	$-0.02 \pm 0.25 \pm 0.53$	
$D \rightarrow VS$			
$D \rightarrow (K^- \pi^0)_{S\text{-wave}} \rho^+$	$14.63 \pm 1.70 \pm 2.41$	$-2.39 \pm 0.11 \pm 0.35$	
$D \rightarrow K^{*-}(\pi^+ \pi^0)_S$	$0.80 \pm 0.38 \pm 0.26$	$1.59 \pm 0.19 \pm 0.24$	
$D \rightarrow K^{*0}(\pi^0 \pi^0)_S$	$0.12 \pm 0.27 \pm 0.27$	$1.45 \pm 0.48 \pm 0.51$	
$D \rightarrow VP, V \rightarrow VP$			
$D \rightarrow (K^{*-} \pi^+)_V \pi^0$	$2.25 \pm 0.43 \pm 0.45$	$0.52 \pm 0.12 \pm 0.17$	BESIII Preliminary
$D \rightarrow VV$			
$D[S] \rightarrow K^{*-} \rho^+$	$5.15 \pm 0.75 \pm 1.28$	$1.24 \pm 0.11 \pm 0.23$	
$D[P] \rightarrow K^{*-} \rho^+$	$3.25 \pm 0.55 \pm 0.41$	$-2.89 \pm 0.10 \pm 0.18$	
$D[D] \rightarrow K^{*-} \rho^+$	$10.90 \pm 1.53 \pm 2.36$	$2.41 \pm 0.08 \pm 0.16$	
$D[P] \rightarrow (K^- \pi^0)_V \rho^+$	$0.36 \pm 0.19 \pm 0.27$	$-0.94 \pm 0.19 \pm 0.28$	
$D[D] \rightarrow (K^- \pi^0)_V \rho^+$	$2.13 \pm 0.56 \pm 0.92$	$-1.93 \pm 0.22 \pm 0.25$	
$D[D] \rightarrow K^{*-}(\pi^+ \pi^0)_V$	$1.66 \pm 0.52 \pm 0.61$	$-1.17 \pm 0.20 \pm 0.39$	
$D[S] \rightarrow (K^- \pi^0)_V(\pi^+ \pi^0)_V$	$5.17 \pm 1.91 \pm 1.82$	$-1.74 \pm 0.20 \pm 0.31$	
$D \rightarrow TS$			
$D \rightarrow (K^- \pi^+)_{S\text{-wave}}(\pi^0 \pi^0)_T$	$0.30 \pm 0.21 \pm 0.32$	$-2.93 \pm 0.31 \pm 0.82$	
$D \rightarrow (K^- \pi^0)_{S\text{-wave}}(\pi^+ \pi^0)_T$	$0.14 \pm 0.12 \pm 0.10$	$2.23 \pm 0.38 \pm 0.65$	

Amplitude Analysis Results of $D^0 \rightarrow K^- \pi^+ \pi^0 \pi^0$

Fit projections:

BESIII Preliminary

BESIII Preliminary




Points with error bars: data, red histograms: fit.

Branching Fraction Results of $D^0 \rightarrow K^- \pi^+ \pi^0 \pi^0$


Fits to M_{BC} distributions of DT and ST data:

$$\text{Beam-Constrained Mass: } M_{BC} = \sqrt{E_{\text{beam}}^2 - |\vec{p}_D|^2}$$


Signal: M_{BC} peaks at D mass



DT ($K^- \pi^+ \pi^0 \pi^0$)



DT ($K^+ \pi^-$)



ST

DT yield = 6101 ± 83 ; ST yield = 534581 ± 769 .

The amplitude analysis result is used to determine the detection efficiency.

Preliminary result:

$$\mathcal{B}(D^0 \rightarrow K^- \pi^+ \pi^0 \pi^0) = (8.98 \pm 0.13_{\text{stat.}} \pm 0.40_{\text{sys.}})\%$$

First measurement

Amplitude Analysis Results of $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$

Double tag: The D^+ is reconstructed by $K_S^0 \pi^+ \pi^+ \pi^-$ with D^- reconstructed by $K^+ \pi^- \pi^-$.

A sample of 4559 events with purity ~99% is used to perform the amplitude analysis.

The data can be described with 12 amplitudes:


BESIII Preliminary

Amplitude	ϕ	Fit fraction
$D^+ \rightarrow K_S^0 a_1(1260)^+, a_1(1260)^+ \rightarrow \rho^0 \pi^+[S]$	0.000(fixed)	$0.567 \pm 0.020 \pm 0.044$
$D^+ \rightarrow K_S^0 a_1(1260)^+, a_1(1260)^+ \rightarrow f_0(500) \pi^+$	$-2.023 \pm 0.068 \pm 0.113$	$0.050 \pm 0.006 \pm 0.007$
$D^+ \rightarrow \bar{K}_1(1400)^0 \pi^+, \bar{K}_1(1400)^0 \rightarrow K^{*-} \pi^+[S]$	$-2.714 \pm 0.038 \pm 0.051$	$0.380 \pm 0.013 \pm 0.014$
$D^+ \rightarrow \bar{K}_1(1400)^0 \pi^+, \bar{K}_1(1400)^0 \rightarrow K^{*-} \pi^+[D]p$	$3.431 \pm 0.137 \pm 0.117$	$0.015 \pm 0.004 \pm 0.005$
$D^+ \rightarrow \bar{K}_1(1270)^0 \pi^+, \bar{K}_1(1270)^0 \rightarrow K_S^0 \rho^0[S]$	$-0.418 \pm 0.070 \pm 0.087$	$0.036 \pm 0.004 \pm 0.002$
$D^+ \rightarrow \bar{K}(1460)^0 \pi^+, \bar{K}(1460)^0 \rightarrow K_S^0 \rho^0$	$-1.850 \pm 0.120 \pm 0.223$	$0.014 \pm 0.004 \pm 0.003$
$D^+ \rightarrow (K_S^0 \rho^0)_A[D] \pi^+$	$2.328 \pm 0.097 \pm 0.068$	$0.011 \pm 0.003 \pm 0.002$
$D^+ \rightarrow K_S^0 (\rho^0 \pi^+)_P$	$1.656 \pm 0.083 \pm 0.056$	$0.031 \pm 0.004 \pm 0.010$
$D^+ \rightarrow (K^{*-} \pi^+)_A[S] \pi^+$	$1.962 \pm 0.047 \pm 0.073$	$0.132 \pm 0.011 \pm 0.011$
$D^+ \rightarrow (K^{*-} \pi^+)_A[D] \pi^+$	$0.989 \pm 0.158 \pm 0.229$	$0.013 \pm 0.004 \pm 0.004$
$D^+ \rightarrow (K_S^0 (\pi^+ \pi^-)_S)_A \pi^+$	$-2.935 \pm 0.060 \pm 0.125$	$0.051 \pm 0.004 \pm 0.003$
$D^+ \rightarrow ((K_S^0 \pi^-)_S \pi^+)_P \pi^+$	$1.864 \pm 0.069 \pm 0.288$	$0.022 \pm 0.003 \pm 0.003$

Amplitude Analysis Results of $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$

Fit projections:

BESIII Preliminary



Points with error bars: data, red histograms: fit, green histograms: background estimated from MC.

For the two identical π^+ , we require $m(\pi_1^+ \pi^-) < m(\pi_2^+ \pi^-)$.

Branching fraction results of $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$

The preliminary results of branching fractions for different components:

BESIII Preliminary

Component	Branching fraction (%)
$D^+ \rightarrow K_S^0 a_1(1260)^+ (\rho^0 \pi^+)$	$1.684 \pm 0.059 \pm 0.131 \pm 0.062$
$D^+ \rightarrow K_S^0 a_1(1260)^+ (f_0(500) \pi^+)$	$0.149 \pm 0.018 \pm 0.021 \pm 0.006$
$D^+ \rightarrow K_1(1400)^0 (K^{*-} \pi^+) \pi^+$	$1.105 \pm 0.045 \pm 0.048 \pm 0.041$
$D^+ \rightarrow \bar{K}_1(1270)^0 (K_S^0 \rho^0) \pi^+$	$0.107 \pm 0.012 \pm 0.006 \pm 0.004$
$D^+ \rightarrow \bar{K}(1460)^0 (K_S^0 \rho^0) \pi^+$	$0.042 \pm 0.012 \pm 0.009 \pm 0.002$
$D^+ \rightarrow K_S^0 \pi^+ \rho^0$	$0.131 \pm 0.015 \pm 0.015 \pm 0.005$
$D^+ \rightarrow K^{*-} \pi^+ \pi^+$	$0.413 \pm 0.036 \pm 0.059 \pm 0.015$
$D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$	$0.220 \pm 0.015 \pm 0.024 \pm 0.008$

stat. uncertainty from FF

sys. uncertainty from FF

uncertainties related to $\text{BF}(D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-)$ in PDG

- The measurements of the decays with $K_1(1270)$ and $K_1(1400)$ involved provide some experimental information in understanding the mixture of the two excited Kaons.

(PLB707,116).

Measurement of the Branching Fractions of $D \rightarrow PP$

- The analysis of $D \rightarrow PP$ modes provides materials for the study of SU(3) breaking effect.
- Most of the $D \rightarrow PP$ decays have been studied by CLEO in 2010 , other measurements come from Belle , BABAR and CDF , etc.

Branching fraction results of $D \rightarrow PP$

PRD97,072004

BFs of $D \rightarrow PP$ are obtained using ST method:

Mode	$\mathcal{B} (\times 10^{-3})$	$\mathcal{B}_{\text{PDG}} (\times 10^{-3})$
$D^+ \rightarrow \pi^+ \pi^0$	$1.259 \pm 0.033 \pm 0.023$	1.24 ± 0.06
$D^+ \rightarrow K^+ \pi^0$	$0.232 \pm 0.021 \pm 0.006$	0.189 ± 0.025
$D^+ \rightarrow \pi^+ \eta$	$3.790 \pm 0.070 \pm 0.068$	3.66 ± 0.22
$D^+ \rightarrow K^+ \eta$	$0.151 \pm 0.025 \pm 0.014$	0.112 ± 0.018
$D^+ \rightarrow \pi^+ \eta'$	$5.12 \pm 0.14 \pm 0.024$	4.84 ± 0.31
$D^+ \rightarrow K^+ \eta'$	$0.164 \pm 0.051 \pm 0.024$	0.183 ± 0.023
$D^+ \rightarrow K_S^0 \pi^+$	$15.91 \pm 0.06 \pm 0.30$	15.3 ± 0.6
$D^+ \rightarrow K_S^0 K^+$	$3.183 \pm 0.029 \pm 0.060$	2.95 ± 0.15
$D^0 \rightarrow \pi^+ \pi^-$	$1.508 \pm 0.018 \pm 0.022$	1.421 ± 0.025
$D^0 \rightarrow K^+ K^-$	$4.233 \pm 0.021 \pm 0.064$	4.01 ± 0.07
$D^0 \rightarrow K^\mp \pi^\pm$	$38.98 \pm 0.06 \pm 0.51$	39.4 ± 0.4
$D^0 \rightarrow K_S^0 \pi^0$	$12.39 \pm 0.06 \pm 0.27$	12.0 ± 0.4
$D^0 \rightarrow K_S^0 \eta$	$5.13 \pm 0.07 \pm 0.12$	4.85 ± 0.30
$D^0 \rightarrow K_S^0 \eta'$	$9.49 \pm 0.20 \pm 0.36$	9.5 ± 0.5

- The results from BESIII are consistent with the world average values within uncertainties. The BFs of $D^+ \rightarrow \pi^+ \pi^0, K^+ \pi^0, \pi^+ \eta, \pi^+ \eta', K_S^0 \pi^+, K_S^0 K^+$ and $D^0 \rightarrow K_S^0 \pi^0, K_S^0 \eta, K_S^0 \eta'$ are determined with improved precision.

Summary

BESIII provides large data samples close to charm related threshold to study the $D_{(s)}$ hadronic decays:

- Observation of pure W -annihilation decays $D_s^+ \rightarrow p\bar{n}, D_s^+ \rightarrow \omega\pi^+, D_s^+ \rightarrow a_0(980)\pi$
 - Our preliminary results on $D_s^+ \rightarrow p\bar{n}, D_s^+ \rightarrow \omega\pi^+$ confirm CLEO's measurements with greatly improved precision.
 - Our preliminary results on $D_s^+ \rightarrow a_0(980)\pi$ are larger than other measured pure W -annihilation decays ($D_s^+ \rightarrow p\bar{n}, D_s^+ \rightarrow \omega\pi^+$) by one order.
- Amplitude analysis of $D \rightarrow K\pi\pi\pi$:
 - $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ is published in PRD95,072010.
 - Preliminary results of $D^0 \rightarrow K^-\pi^+\pi^0\pi^0, D^+ \rightarrow K_S^0\pi^+\pi^+\pi^-$ are obtained.
$$\mathcal{B}(D^0 \rightarrow K^-\pi^+\pi^0\pi^0) = (8.98 \pm 0.13_{\text{stat.}} \pm 0.40_{\text{sys.}})\%$$
- Branching fractions of $D \rightarrow PP$:

The BF's of 14 decay modes are published in PRD97,072004.

More measurements in $D_{(s)}$ hadronic decays are coming.

Thank you for your attention!