



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 2011

Type: **Invited Speaker / Conférencier(ère) invité(e)**

Application of the exchange-hole dipole moment dispersion model to surfaces and 2D materials (I)

Thursday, 14 June 2018 08:30 (30 minutes)

The exchange-hole dipole moment (XDM) method is a density-functional model of London dispersion based upon second-order perturbation theory. The XDM dispersion coefficients are non-empirical and depend directly on the electron density and related properties. XDM offers simultaneous high accuracy for a diverse range of systems due to the variation of the calculated atomic dispersion coefficients with the electronic environment. In this talk, we focus on applications of XDM to adsorption of small molecules and of graphene on noble metal surfaces. We also consider the interlayer interactions in 2D electrides, an unusual class of materials possessing interstitial electron layers sandwiched between cationic atomic layers of the solid.

Primary author: Prof. JOHNSON, Erin (Department of Chemistry, Dalhousie University)

Presenter: Prof. JOHNSON, Erin (Department of Chemistry, Dalhousie University)

Session Classification: R1-1 Theoretical Modeling of Materials 1 (DCMMP) | La modélisation théorique de matériaux 1 (DPMCM)

Track Classification: Condensed Matter and Materials Physics / Physique de la matière condensée et matériaux (DCMMP-DPMCM)