


The logo for the Deutsche Forschungsgemeinschaft (DFG), consisting of the letters 'DFG' in a bold, blue, sans-serif font.

Deutsche  
Forschungsgemeinschaft



**CONACYT**

Consejo Nacional de Ciencia y Tecnología

# Global fits of the SM parameters


**Jens Erler**

**JGU & Helmholtz Institute Mainz (on leave from IF-UNAM)**

**LHCP 2019**

**May 20–25, 2019**

**Puebla, Mexico**



**Weak mixing angle:**  
**global survey of**  
 **$\sin^2\theta_w$  determinations**

# Why pushing $\sin^2\theta_W$ ?

- compute and measure  $\sin^2\theta_W$  and relate to  $M_W$
- ➔ **doubly over-constrained** system at sub-‰ precision
- key test of EW symmetry breaking sector
- comparisons of different measurements, scales, and initial/final states provide window to physics beyond the SM
- ➔ global analysis

# $\sin^2\theta_w(0)$ : approaches

- tuning in on the Z resonance
  - FB and LR asymmetries in  $e^+e^-$  annihilation near  $s = M_Z^2$
  - FB asymmetries in  $pp$  ( $p\bar{p}$ ) Drell-Yan around  $m_{ll} = M_Z$

	$\nu$ scattering	PVES
leptonic	$\nu_\mu - e^-$	$e^- - e^-$
DIS	heavy nuclei (NuTeV)	deuteron (PVDIS, SoLID)
elastic	<b>CEvNS (COHERENT)</b>	<b>proton, <math>^{12}\text{C}</math> (Qweak, P2)</b>
APV	heavy alkali atoms and ions	<b>isotope ratios (Mainz)</b>

# $\sin^2\theta_w(0)$ : approaches


- tuning in on the Z resonance

- FB and LR asymmetries in  $e^+e^- \rightarrow e^+e^-$  at  $\sqrt{s} = M_Z^2$
- FB asymmetries in  $e^-p$  at  $\sqrt{s} \approx M_Z$

**very recent first measurements**

	$\nu$ scattering	PVES
leptonic	$\nu_\mu - e^-$	$e^- - e^-$
DIS	heavy nuclei (NuTeV)	deuteron (PVDIS, SoLID)
elastic	<b>CEvNS (COHERENT)</b>	<b>proton, <math>^{12}\text{C}</math> (Qweak, P2)</b>
APV	heavy alkali atoms and ions	<b>isotope ratios (Mainz)</b>

# $\sin^2\theta_W$ measurements



**LEP & SLC:**

**$0.23153 \pm 0.00016$**

**Tevatron:**

**$0.23148 \pm 0.00033$**

**LHC:**

**$0.23131 \pm 0.00033$**


**average direct**

**$0.23149 \pm 0.00013$**

**global fit**

**$0.23153 \pm 0.00004$**

# $\sin^2\theta_W$ measurements



**LEP & SLC:**  
 **$0.23153 \pm 0.00016$**

**Tevatron:**

**$0.23148 \pm 0.00033$**

**LHC:**

**$0.23131 \pm 0.00033$**


**average direct**

**$0.23149 \pm 0.00013$**

**global fit**

**$0.23153 \pm 0.00004$**

# $\sin^2\theta_W$ measurements



**LEP & SLC:**  
 **$0.23153 \pm 0.00016$**

**Tevatron:**

**$0.23148 \pm 0.00033$**

**LHC:**

**$0.23131 \pm 0.00033$**

**average direct**


**$0.23149 \pm 0.00013$**

**global fit**

**$0.23153 \pm 0.00004$**



# $M_W$ measurements



*plenary talks by  
Daniele Zanzi &  
Alessandro Vicini*

**average direct**

**$80.379 \pm 0.012$  GeV**

**indirect**

**$80.357 \pm 0.006$  GeV**


**(incl. correlated theory errors)**

**Theoretical uncertainties:**  
**correlations in**  
**precision observables**

# Theory errors

- hadronic vacuum polarization and light-by-light ( $g_\mu - 2$ )
- non-factorizable QCD corrections ( $\Gamma_Z^{\text{had}}$ )
- non-resonant corrections to Breit-Wigner shape ( $\sigma_{\text{had}}$ )  
*Grassi, Kniehl & Sirlin, PRL 86 (2001)*
- W & Z self-energies
  - loop factors including enhancement factors such as  $N_C = N_F = 3$  or  $\sin^{-2}\theta_W \approx m_t^2/M_W^2 \approx 4$   
amount to **0.020** (QED), **0.116** (QCD), **0.032** (CC), **0.029** (NC)
  - parametrized by  $\Delta S_Z = \pm 0.0034$  (may be combined with  $\Delta\alpha_{\text{had}}$ ),  
 $\Delta T = \pm 0.0073$  (t-b doublet) and  $\Delta U = S_W - S_Z = \pm 0.0051$
  - assuming  $\Delta S_Z$ ,  $\Delta T$  and  $\Delta U$  to be sufficiently different (uncorrelated) induces **theory correlations** between different observables *Schott & JE, PPNP 106 (2019)*

# $M_H - m_t$



indirect  $m_t$ :  
 $176.4 \pm 1.8$  GeV  
 (2.0  $\sigma$  high)

indirect  $M_H$ :  
 $90^{+17}_{-15}$  GeV  
 (1.9  $\sigma$  low)

incl. theory error:

indirect  $M_H$ :  
 $91^{+18}_{-16}$  GeV  
 (1.8  $\sigma$  low)

# **Vacuum polarizations in global fits:**

**$\alpha(M_Z)$   $\sin^2\theta_w(0)$   $g_{\mu-2}$   $m_{b,c}$**

# $\alpha(M_Z)$

- Dispersive approach: integral over  $\sigma(e^+e^- \rightarrow \text{hadrons})$  and  $\tau$ -decay data
- $\alpha^{-1}(M_Z) = 128.947 \pm 0.012$  *Davier et al., EPJC 77 (2017)*
- $\alpha^{-1}(M_Z) = 128.958 \pm 0.016$  *Jegerlehner, arXiv:1711.06089*
- $\alpha^{-1}(M_Z) = 128.946 \pm 0.015$  *Keshavarzi et al., PRD 97 (2018)*
- $\alpha^{-1}(M_Z) = 128.949 \pm 0.010$  *Ferro-Hernández & JE, JHEP 03 (2018)*
  - **This value** is converted from the  $\overline{MS}$  scheme and uses both  $e^+e^-$  annihilation and  $\tau$  decay spectral functions  
*Davier et al., EPJC 77 (2017)*
  - PQCD for  $\sqrt{s} > 2$  GeV (using  $\bar{m}_c$  &  $\bar{m}_b$ )
- (anti)correlation with  $g_\mu - 2$  at two (three) loop order and with  $\sin^2\theta_W(0)$

# $g_\mu - 2$


PQCD:

*Luo & JE, PRL 87 (2001)*


$$(a_\mu^{\text{hvp}})^c = (14.6 \pm 0.5_{\text{theory}} \pm 0.2_{\text{mc}} \pm 0.1_{\alpha_s}) \times 10^{-10} \quad (a_\mu^{\text{hvp}})^b = 0.3 \times 10^{-10}$$

Lattice gauge theory:

*A. Gérardin et al., arXiv:1904.03120*




# $\sin^2\theta_w(\mu)$



Ferro-Hernández & JE, JHEP 03 (2018)




# $\sin^2\theta_w(\mu)$



*Ferro-Hernández & JE, JHEP 03 (2018)*

# $\sin^2\theta_w(0)$ and $\Delta\alpha(M_Z)$



$$\mu^2 \frac{d\hat{v}_f}{d\mu^2} = \frac{\hat{\alpha} Q_f}{24\pi} \left[ \sum_i K_i \gamma_i \hat{v}_i Q_i + 12\sigma \left( \sum_q Q_q \right) \left( \sum_q \hat{v}_q \right) \right]$$

$$\mu^2 \frac{d\hat{\alpha}}{d\mu^2} = \frac{\hat{\alpha}^2}{\pi} \left[ \frac{1}{24} \sum_i K_i \gamma_i Q_i^2 + \sigma \left( \sum_q Q_q \right)^2 \right]$$


- coupled system of differential equations *Ramsey-Musolf & JE, PRD 72 (2005)*

- $\Delta\alpha(M_Z)_{\text{had}}$  errors in  $\sin^2\theta_w(0) = \kappa(0) \sin^2\theta_w(M_Z)$  **add** since

$$M_Z^2 \propto g_Z^2(M_Z) v^2 \propto [\alpha / s_w^2 c_w^2](M_Z) G_F^{-1}$$

# $\bar{m}_c(\bar{m}_c)$

- derived from another set of dispersion integrals
- input: **electronic widths** of  $J/\psi$  and  $\psi(2S)$
- continuum contribution from **self-consistency between sum rules**



- $\bar{m}_c(\bar{m}_c) = 1272 \pm 8 + 2616 [\bar{\alpha}_s(M_Z) - 0.1182] \text{ MeV}$   
*Masjuan, Spiesberger & JE, EPJC 77 (2017)*

# **Fit Results**

Performed with package **GAPP**  
(**G**lobal **A**nalysis of **P**article **P**roperties)

# Standard global fit

$M_H$	$125.14 \pm 0.15 \text{ GeV}$
$M_Z$	$91.1884 \pm 0.0020 \text{ GeV}$
$\bar{m}_b(\bar{m}_b)$	$4.180 \pm 0.021 \text{ GeV}$
$\Delta\alpha_{\text{had}}^{(3)}(2 \text{ GeV})$	$(59.0 \pm 0.5) \times 10^{-4}$

$\bar{m}_t(\bar{m}_t)$	$163.28 \pm 0.44 \text{ GeV}$	1.00	-0.13	-0.28
$\bar{m}_c(\bar{m}_c)$	$1.275 \pm 0.009 \text{ GeV}$	-0.13	1.00	0.45
$\alpha_s(M_Z)$	$0.1187 \pm 0.0016$	-0.28	0.45	1.00


other correlations small

*Freitas & JE, PDG 2018*

# $\rho_0$ fit

- $\Delta\rho_0 = G_F \sum_i C_i / (8\sqrt{2}\pi^2) \Delta m_i^2$ 
  - where  $\Delta m_i^2 \geq (m_1 - m_2)^2$
  - despite appearance there is decoupling (see-saw type suppression of  $\Delta m_i^2$ )
- $\rho_0 = 1.00039 \pm 0.00019$  (2.0  $\sigma$ )
  - $(16 \text{ GeV})^2 \leq \sum_i C_i / 3 \Delta m_i^2 \leq (48 \text{ GeV})^2$  @ 90% CL
  - $Y = 0$  Higgs triplet VEVs  $v_3$  strongly disfavored ( $\rho_0 < 1$ )
  - consistent with  $|Y| = 1$  Higgs triplets if  $v_3 \sim 0.01 v_2$

# S and T



S	$0.02 \pm 0.07$
T	$0.06 \pm 0.06$
$\Delta\chi^2$	-4.2


- $M_{\text{KK}} \approx 3.2 \text{ TeV}$  in warped extra dimension models
- $M_V \approx 4 \text{ TeV}$  in minimal composite Higgs models *Freitas & JE, PDG (2018)*

# Conclusions and outlook


- LHC & low-energy experiments approaching LEP precision in  $\sin^2\theta_w$
- new players:
  - coherent  $\nu$ -scattering
  - ultra-high precision PVES
  - APV isotope ratios
- at ultra-high precision not only theoretical uncertainties are relevant, but also their **correlations** (hard to estimate)
  - example: vacuum polarization uncertainties enter correlated in an increasing number of quantities



**Backups**




# $m_c$



- $\alpha(M_Z)$  and  $\sin^2\theta_W(0)$ : can use PQCD for heavy quark contribution if masses are known.
- $g-2$ : c quark contribution to muon  $g-2$  similar to  $\gamma^*\gamma$ ;  $\pm 70$  MeV uncertainty in  $m_c$  induces an error of  $\pm 1.6 \times 10^{-10}$  comparable to the projected errors for the FNAL and J-PARC experiments.
- Yukawa coupling – mass relation (in single Higgs doublet SM):  $\Delta m_b = \pm 9$  MeV and  $\Delta m_c = \pm 8$  MeV to match precision from HiggsBRs @ FCC-ee
- QCD sum rule:  $m_c = 1272 \pm 8$  MeV  
*Masjuan, Spiesberger & JE, EPJC 77 (2017)*  
(expect about twice the error for  $m_b$ )

# Effective couplings



# $m_t$ measurements


	central	statistical	systematic	total
<b>Tevatron</b>	<b>174.30</b>	<b>0.35</b>	<b>0.54</b>	<b>0.64</b>
<b>ATLAS</b>	<b>172.51</b>	<b>0.27</b>	<b>0.42</b>	<b>0.50</b>
<b>CMS</b>	<b>172.43</b>	<b>0.13</b>	<b>0.46</b>	<b>0.48</b>
<b>CMS Run 2</b>	<b>172.25</b>	<b>0.08</b>	<b>0.62</b>	<b>0.63</b>
<b>grand average</b>	<b>172.74</b>	<b>0.11</b>	<b>0.31</b>	<b>0.33</b>

*JE, EPJC 75 (2015)*

- $m_t = 172.74 \pm 0.25_{\text{uncorr.}} \pm 0.21_{\text{corr.}} \pm 0.32_{\text{QCD}} \text{ GeV} = 172.74 \pm 0.46 \text{ GeV}$
- somewhat larger shifts and smaller errors conceivable in the future  
*Butenschoen et al., PRL 117 (2016); Andreassen & Schwartz, JHEP 10 (2017)*
- 2.8  $\sigma$  discrepancy between lepton + jet channels from DØ and CMS Run 2
- **indirectly** from EW fit:  $m_t = 176.4 \pm 1.8 \text{ GeV} (2 \sigma)$  *Freitas & JE (PDG 2018)*

# Features of our approach

- only experimental input: **electronic widths** of  $J/\psi$  and  $\psi(2S)$
- continuum contribution from **self-consistency between sum rules**
- include  $M_0 \rightarrow$  stronger (milder) sensitivity to continuum ( $m_c$ )
- quark-hadron duality needed only in finite region (**not locally**)





- $\bar{m}_c(\bar{m}_c) = 1272 \pm 8 + 2616 [\bar{\alpha}_s(M_Z) - 0.1182] \text{ MeV}$   
*Masjuan, Spiesberger & JE, EPJC 77 (2017)*

# $\sin^2\theta_W(0)$ : flavor separation

strange quark external current	ambiguous external current
$\Phi$	$K\bar{K}$ (non - $\Phi$ )
$K\bar{K}\pi$ [almost saturated by $\Phi(1680)$ ]	$K\bar{K}2\pi$ , $K\bar{K}3\pi$
$\eta\Phi$	$K\bar{K}\eta$ , $K\bar{K}\omega$

- use of result for  $\alpha(2 \text{ GeV})$  also needs isolation of strange contribution  $\Delta_s\alpha$
- left column assignment assumes OZI rule
- expect right column to originate mostly from strange current ( $m_s > m_{u,d}$ )
- quantify expectation using averaged  $\Delta_s(g_\mu-2)$  from lattices as Bayesian prior  
*RBC/UKQCD, JHEP 04 (2016); HPQCD, PRD 89 (2014)*
- $\Delta_s\alpha(1.8 \text{ GeV}) = (7.09 \pm 0.32) \times 10^{-4}$  (threshold mass  $\bar{m}_s = 342 \text{ MeV} \approx \bar{m}_s^{\text{disc}}$ )

# $\sin^2\theta_W(0)$ : singlet separation



*Ferro-Hernández & JE, JHEP 03 (2018)*  
 adapted from lattice  $g_{\mu-2}$  calculation  
*RBC/UKQCD, PRL 116 (2016)*

- use of result for  $\alpha(2 \text{ GeV})$  needs singlet piece isolation  $\Delta_{\text{disc}} \alpha(2 \text{ GeV})$
- then  $\Delta_{\text{disc}} \bar{s}^2 = (\bar{s}^2 \pm 1/20) \Delta_{\text{disc}} \alpha(2 \text{ GeV}) = (-6 \pm 3) \times 10^{-6}$
- **step function**  $\Rightarrow$  singlet threshold mass  $\bar{m}_s^{\text{disc}} \approx 350 \text{ MeV}$



# S fit

- S parameter rules out QCD-like technicolor models
- S also constrains extra degenerate fermion families:
  - ➔  $N_F = 2.75 \pm 0.14$  (assuming  $T = U = 0$ )
- compare with  $N_\nu = 2.991 \pm 0.007$  from  $\Gamma_Z$

# STU fit

$\sin^2\theta_W(M_Z)$	$0.23113 \pm 0.00014$
$\alpha_s(M_Z)$	$0.1189 \pm 0.0016$

S	$0.02 \pm 0.10$	1.00	0.92	-0.66
T	$0.07 \pm 0.12$	0.92	1.00	-0.86
U	$0.00 \pm 0.09$	-0.66	-0.86	1.00

- $M_{KK} \approx 3.2 \text{ TeV}$  in warped extra dimension models
- $M_V \approx 4 \text{ TeV}$  in minimal composite Higgs models *Freitas & JE (PDG 2018)*