


Search for axions with the International AXion Observatory (IAXO)

Igor G. Irastorza (U. Zaragoza)
on behalf of the IAXO collaboration


Dark Matter Review Meeting, LSC, February 7th 2018





Axion motivation in a nutshell



Axion/ALP searches motivation



Solar Axions



Standard Primakoff spectrum


Robust prediction from axion models. Conversion of solar plasma photons into axion (only axion-photon coupling involved)

van Bibber PRD 39 (89)
CAST JCAP 04(2007)010

Non-hadronic “ABC” Solar axion flux at Earth (only if axion couples to electron)

Redondo, JCAP 1312 008

Axion helioscopes



Axion helioscope concept
 P. Sikivie, 1983
 + K. van Bibber, G. Raffelt,
 et al. (1989)
 (use of buffer gas)



X-ray Focussing &
 low-background
 Later innovations by
 CAST

$$g_{a\gamma}^4 \propto \underbrace{b^{1/2} \epsilon^{-1}}_{\text{detectors}} \times \underbrace{a^{1/2} \epsilon_o^{-1}}_{\text{optics}} \times \underbrace{(BL)^{-2} A^{-1}}_{\text{magnet}} \times \underbrace{t^{-1/2}}_{\text{exposure}}$$

Pioneer implementations of helioscope concept:
 Brookhaven (just few hours of data) [Lazarus et al. PRL 69 (92)]
 TOKYO Helioscope (SUMICO): 2.3 m long 4 T magnet

IAXO experiment reminder

- Next generation “axion helioscope” after CAST
- Purpose-built large-scale magnet
 - >300 times larger B^2L^2A than CAST magnet
 - Toroid geometry
 - 8 conversion bores of 60 cm \varnothing , ~20 m long
- Detection systems (XRT+detectors)
 - Scaled-up versions based on experience in CAST
 - Low-background techniques for detectors
 - Optics based on slumped-glass technique used in NuStar
- ~50% Sun-tracking time
- Large magnetic volume available for additional “axion” physics (e.g. DM setups)




IAXO pathfinder system at CAST

In operation in 2014-15

Last CAST results published in Nature Physics last May
Nature Phys. 13 (2017) 584-590

Latest CAST limit

Enabled by the IAXO pathfinder system



X-ray optics specifically built for axions

Low background Micromegas

nature
physics

ARTICLES


PUBLISHED ONLINE: 1 MAY 2017 | DOI: 10.1038/NPHYS4109

OPEN

New CAST limit on the axion-photon interaction

CAST Collaboration[†]

Hypothetical low-mass particles, such as axions, provide a compelling explanation for the dark matter in the universe. Such particles are expected to emerge abundantly from the hot interior of stars. To test this prediction, the CERN Axion Solar Telescope (CAST) uses a 9 T refurbished Large Hadron Collider test magnet directed towards the Sun. In the strong magnetic 2013-2015 run, thanks about a factor of three, confidence level) set by





$$g_{ay} < 0.66 \times 10^{-10} \text{ GeV}^{-1} \text{ at 95\% CL}$$

IAXO pathfinder system in CAST


Detector: JCAP12 (2015)

Physics: Nature Physics (10.1038/nphys4109)



Test MM detector + slumped-glass x-ray optics together
Best SNR than any other CAST detector ever!!



X-ray optics specifically built for axions



Low background Micromegas at the focal point



- Best SNR of any previous detector
- 290 tracking hour acquired (6.5 months operation)
- 3 counts observed in RoI (1 expected)

IAXO-D0 at UNIZAR

- Based on IAXO pathfinder (improved):
 - Electroformed Cu chamber and pipes
 - Enlarged Pb shielding
 - 4π active shielding
 - AGET electronics (auto-trigger)
 - Operation with Xe
- New lab at UNIZAR
 - Comissioning of the system underway
- Software tools available
 - Background model underway

Goals: Bkg level: 10^{-7} - 10^{-8} keV $^{-1}$ cm $^{-2}$ s $^{-1}$
Energy threshold: ~ 0.1 keV

Test facility for other IAXO detector candidates




IAXO most recent milestones

- Collaboration formally established in the last collaboration meeting at DESY (last July):
 - 17 institutions to sign IAXO bylaws
 - Management structure defined. Positions being elected.
- Near term goal: **BabyIAXO** → Full experiment “scaled-down IAXO”
 - Test bench for magnet, optics and detectors. Risk reduction, but also opportunity to go beyond baseline FOM for IAXO (→ IAXO+)
 - Intermediate scale, but representative of full IAXO dimensions
 - **Relevant physics outcome**
- Experimental site for IAXO: Strong interest from DESY
 - BabyIAXO: DESY, INR
- Magnet under design with CERN support. (Applied Fellow started last month)
- IAXO progress is being followed by “Physics Beyond Colliders” process at CERN (to provide feedback for the European Strategy Part Phy)
- Construction of BabyIAXO could happen in 2-3 years. MoU under preparation, collaboration addressing funding bodies.

IAXO collaboration

17 institutions from Germany, Spain, US, France, Russia, Croatia, S. Africa, CERN.




- Likely to grow in the future, more groups showing interest...
- New partners welcome...

BabyIAXO


- Single bore magnet
- Bore dimensions similar to full IAXO bores → detection line representative of final ones.
- New magnet configuration (saddle dipole). Potential to go to higher B.
- Test & improve all systems. Risk mitigation for full IAXO
- Produce relevant physics
- More staged access to funds
- Move earlier to “experiment mode”

- BabyIAXO CDR finished. Moving to Technical Design



Free bore [m]	0.6
Magnetic length [m]	10
Field in bore [T]	2.5
Stored energy [MJ]	27
Peak field [T]	4.1

10x CAST MFOM



IAXO physics potential ("white paper")

Physics potential of the International Axion Observatory (IAXO)


IAXO Collaboration

- **Comprehensive review of IAXO physics case:**
 - Theory
 - Cosmology
 - Astro hints
 - Updated IAXO sensitivity plots
- **Very advanced draft...**
- **Many contributors from the theory/pheno community**

Contents

1	Introduction	2
2	Axions from high-energy physics	6
3	A meV mass QCD axion as dark matter candidate	8
3.1	The pre-inflationary Peccei-Quinn symmetry breaking scenario	11
3.2	The post-inflationary Peccei-Quinn symmetry breaking scenario	13
3.3	Production of dark matter axions from topological defects	17
3.3.1	Models with $N_{\text{DW}} = 1$	19
3.3.2	Models with $N_{\text{DW}} > 1$	19
4	Axions as dark radiation	22
4.1	Small amounts of thermally produced axion dark radiation	22
4.2	Large amounts of non-thermal axion dark radiation from modulus decay	23
4.3	Additional observational hints of axion dark radiation	25
5	Axions and inflation	27
5.1	Axion hilltop inflation	28
5.2	Reheating of the ALP inflaton into photons	29
5.3	The ALP miracle: axionic unification of inflaton and DM	30
6	Hints for axions in the anomalous cooling of stellar objects	31
6.1	Summary of cooling anomalies	32
6.1.1	Pulsating White Dwarfs	32
6.1.2	The White Dwarf Luminosity Function	32
6.1.3	Globular Clusters	33
6.1.4	He-burning Supergiants	33
6.1.5	Neutron Star in CAS A	34
6.2	Interpretation of the cooling anomalies in terms of axions/ALPs and the IAXO potential	34
7	Axion-like particles in the propagation of photons over astronomical distances	38
7.1	Conversion between gamma-ray photons and ALPs	38
7.2	Hints for anomalies in the gamma-ray opacity	40
7.2.1	flat-spectrum radio quasars	40
7.2.2	Statistical distributions of X-ray source luminosities	40
7.2.3	Extragalactic background light	41
7.3	Constraints from high-energy astronomy and remaining parameter space	42
7.4	Conversion between X-ray photons and ALPs	43
8	Updated sensitivity prospects for IAXO	44
9	Discussion and conclusions	50

Updated IAXO sensitivity regions




babyIAXO prospects:
 10x FOM magnet + optics and detector from conservative scenario of Lol


IAXO phase II:
 exposure rearranged to optimize sensitivity to DSFZ models

IAXO+: enhanced scenario with x4 higher MFOM with respect Lol

IAXO & meV axion cosmology



IAXO & astrophysics hints





IAXO will fully explore ALP models invoked to solve the “transparency hint”

... as well as a large fraction of the axion & ALP models invoked in the “stellar cooling anomaly”
 But for this the g_{ae} is particularly interesting

IAXO & stellar cooling

M. Giannotti et al.
 JCAP 1710 (2017) 010
[arXiv:1708.02111](https://arxiv.org/abs/1708.02111)

- Multiple stellar anomalies (HB, RG, WD, NS,..). Overall 3σ effect.



- IAXO will explore most of the relevant models (especially with IAXO+)
- Only experiment with such capability


Region of QCD axion models that solve the stellar anomaly

IAXO physics reach summary

- IAXO will improve the experimental “helioscope frontier” by more than 1 order of magnitude in sensitivity to $g_{a\gamma\gamma}$
(= by more than 10000 in terms of signal to noise ratio)
- IAXO will probe a large fraction of **QCD axion models** in the meV to eV mass band.
- **No other proposed technique can probe QCD axions in this region.**
← Uniqueness of IAXO.
- IAXO will fully probe the ALP region invoked to solve the transparency anomaly
- IAXO will largely probe the axion region invoked to solve the stellar cooling anomaly
- IAXO will partially explore viable **QCD axion DM** models, and largely probe the **“ALP miracle”** models solving both DM and inflation.
- IAXO will probe a large unexplored ALP parameter space, also generically motivated dark radiation, string theory,...
- **All this, independent of the axion-as-DM hypothesis.**
- **Large complementarity with other detection strategies**

IAXO-DM

- IAXO large magnetic volume offers an appealing environment for competitive axion DM setups.
- Various possible arrangements in IAXO. Leverage the huge magnetic volume available:
 1. Single large cavity tuned to low masses
 2. Thin long cavities tuned to mid-high masses. Possibility for directionality. Add several coherently? → **RADES R&D**
 3. Pick-up coils for vely low mass detection (toroidal geometry preferred)
- **RADES** (Relic Axion Detector Exploratory Setup): testing multicell cavity geometry inside the CAST Magnet





RADES

Haloscope FOM:


$$\mathcal{F} \sim g_{a\gamma}^4 m_a^2 B^4 V^2 T^{-2} C^2 Q$$

V is usually linked with m_a ($V \sim 1/m_a^3$)
how to go to much higher V
and higher m_a ?

- RADES concept: array of small cavities interconnected with irises:
 - No need to phase match externally
 - Clear prescription to optimize coupling to axion:
 - Robust and flexible scaling up
 - Preference for long aspect-ratio: CAST (and IAXO) magnets
 - Problem: how to tune.
- First RADES cavity with 5 subcavities, non-tunable ($f = 8.5$ GHz; $m_a = 34.75$ μeV)




RADES in CAST



Installed successfully in CAST last year, and first engineering run took place Nov-Dec2017

RADES in CAST



Sensitivity to (optimistically coupled) QCD axions with this very simple setup.

Only a factor of a few away from KSVZ.

[But at a fixed m_a value]

Challenge for next runs:

- Larger cavities
- Tuning

IAXO near-term milestones

- 1) Finish definition of management structures of the collaboration (almost there)
- 2) Finish and release “physics potential” paper (next weeks).
- 3) Technical Design of BabyIAXO (2018 – CERN support)
- 4) Site definition (2018)
 - DESY: possible sites identified and “reserved”
 - INR: interest expressed, possible synergy with local resources
- 5) Build BabyIAXO (2-3 years). Collaboration addressing funding bodies
 - MoU under preparation to cover BabyIAXO phase, especially magnet construction.
 - Optics: aim is building full optics. Back-up plan with existing optics (first contacts with ESA Science Directorate already ongoing)
- 6) Full IAXO TDR with feedback from BabyIAXO phase

IAXO and CERN

[from PBC technology subgroup meetings]

- IAXO crucially relies on CERN's expertise on **large-scale superconducting detector magnets**. Both BabyIAXO and IAXO detector magnets are systems of a scale for which CERN's expertise and infrastructure is almost unique.
- The collaboration expects support and significant participation of CERN, in particular, but not exclusively, with knowledge and expert personnel to the **Technical Design Report** of magnets and related infrastructure, as well as the **technical follow-up during construction, testing and commissioning**, first of BabyIAXO and then for IAXO.
- CAST is still active at CERN hosting a number of small-scale prototypes for testing novel concepts or performing R&D. There is potential in further using CAST infrastructure for providing feedback during the preparatory phase of IAXO, for example, to improve the insight on detector backgrounds


IAXO in Spain

- Axion & IAXO in Spain meeting organized in 2016 to raise awareness in our community
- Coordinated project active (FPA2016-76978-C3):
 - **UNIZAR**: physics / coordination / low background detectors (IAXO-D0)
 - **CNM**: R&D new readouts
 - **UPCT**: R&D RF cavities (RADES)
- New groups showed interest...
 - **ICCUB**: formally joined the collaboration. physics + low background electronics +...
- More groups are very welcome!



Conclusions


- 1) Important milestones regarding establishment of the collaboration
- 2) Near term planning better defined.
- 3) BabyIAXO → very appealing new experimental stage that can
 - 1) Enhance final FOM of experiment
 - 2) Catalyze near-term activities in the collaboration towards an intermediate experiment with relevant physics outcome
- 4) Physics case of IAXO getting sharpened by recent studies. Physics potential updated – new paper to be published soon



**IAXO → singular project in the axion experimental landscape with high visibility of Spanish participation.
Great opportunity for our community. New partners are welcome!**


Backup slides...

BabyIAXO & IAXO physics reach



Detecting axions

Source	Experiments	Model & Cosmology dependency	Technology
Relic axions	ADMX, ADMX-HF, Casper, CAPP, ...	High	New ideas emerging, Active R&D going on,...
Lab axions	ALPS, OSQAR, fifth force exps,...	Very low	
Solar axions	SUMICO, CAST, IAXO	Low	Ready for large scale experiment




- Helioscopes → do not rely on the axion being the dominant DM component. Solar axion emission robust prediction
- Helioscopes → No R&D needed. Technology mature enough for a large scale experiment (IAXO)
- Large complementarity with other detection strategies

IAXO status of project

- **2011: First studies concluded (JCAP 1106:013,2011)**
- **2013: Conceptual Design finished (arXiv:1401.3233) and Lol to CERN SPSC [CERN-SPSC-2013-022]**
- **2014: Positive recommendations from SPSC.**
- **2015-16: Transition phase towards TDR (technical design)**
 - Engaging funding agencies
 - Community meetings in many countries
 - Preparatory activities (IAXO pathfinder system at CAST)
 - Preparation of collaboration bylaws and MoU to carry out TDR work.
 - First discussions on experimental site options
 - First dicussion on long-term plan to built full experiment
- **2017: Formal constitution of collaboration (expected for summer)**
 - “Founding” IAXO collaboration meeting to happen in July 2017, DESY


IAXO in the axion landscape



Source	Experiments	Model & Cosmology dependency	Technology
Relic axions	ADMX, ADMX-HF, Casper, CAPP, ...	High	New ideas emerging, Active R&D going on,...
Lab axions	ALPS, OSQAR, fifth force exps,...	Very low	
Solar axions	SUMICO, CAST, IAXO	Low	Ready for large scale experiment



- Helioscopes → do not rely on the axion being the dominant DM component. Solar axion emission robust prediction
- Helioscopes → No R&D needed. Technology mature enough for a large scale experiment (IAXO)
- Large complementarity with other detection strategies

Axion/ALP parameter space






Transparency of the Universe

Stellar cooling hints





Solar Axions

- Solar axions produced by photon-to-axion conversion of the solar plasma photons in the solar core



Solar Axions

- Solar axions produced by photon-to-axion conversion of the solar plasma photons in the solar core




* if the axion couples with the electron (g_{ae})
(non hadronic axion)

Helioscopes


Axion helioscope
concept
P. Sikivie, 1983

+ K. van Bibber, G.
Raffelt, et al. (1989)
(use of buffer gas)




CAST experiment @ CERN

- Decommissioned LHC test magnet (L=10m, B=9 T)
- Moving platform $\pm 8^\circ V \pm 40^\circ H$ (to allow up to 50 days / year of alignment)
- 4 magnet bores to look for X rays
- 2 X ray telescopes to increase signal/noise ratio.



IAXO – Concept

Enhanced axion helioscope:
JCAP 1106:013,2011




$$g_{a\gamma}^4 \propto \underbrace{b^{1/2} \epsilon^{-1}}_{\text{detectors}} \times \underbrace{a^{1/2} \epsilon_o^{-1}}_{\text{optics}} \times \underbrace{(BL)^{-2} A^{-1}}_{\text{magnet}} \times \underbrace{t^{-1/2}}_{\text{exposure}}$$

4+ orders of magnitude better SNR than CAST (JCAP 1106:013)

IAXO – Conceptual Design



- Large toroidal 8-coil magnet $L \approx 20$ m
- 8 bores: 600 mm diameter each
- 8 x-ray telescopes + 8 detection systems
- Rotating platform with services



IAXO technologies – Baseline


IAXO telescopes

- Slumped glass technology with multilayers
- Cost-effective to cover large areas
- Based on NuSTAR developments
- Focal length ~5 m
- 60-70% efficiency
- LLNL+UC+DTU
+ MIT + INAF



IAXO magnet

- Superconducting “detector” magnet.
- Toroidal geometry (8 coils)
- Based on ATLAS toroid technical solutions.
- CERN+CEA expertise
- 8 bores / 20 m long / 60 cm Ø per bore



Baseline developed at:
IAXO Letter of Intent: CERN-SPSC-2013-022
IAXO Conceptual Design: JINST 9 (2014)
T05002 (arXiv:1401.3233)




Rotation System

IAXO technologies – Baseline

IAXO detectors

- **Micromegas** gaseous detectors
- Radiopure components + shielding
- Discrimination from event topology in gas
- Long trajectory in CAST
- Zaragoza + CEA + Bonn + others expertise

Optics+detector IAXO pathfinder system
(in operation in CAST during 2014-5)





Calibration photons (source 14 m away) focused onto the Micromegas

Services

IAXO detectors


Ingrid detectors

- Better threshold
- U. Bonn




MMC (Magnetic Metallic Calorimeters) TES (Transition Edge Sensors)

- Very good E resolution & threshold
- Heidelberg + CEA + CNSMS



Low noise CCDs




Solar axion spectroscopy:
Axion-electron ABC spectrum
Axion mass determination


Rotation System

Axion-electron coupling

ABC-produced solar axions



Sensitive to g_{ae} values down to $\sim 10^{-13}$




And more...

- Dark Matter Axions (\rightarrow use of IAXO magnet with cavities / RF antennas)
- ALPs from Dark Radiation
- ALPs from nearby Supernova (connection with SNEWS)
 \rightarrow MeV photon detector at the other end
- ...

Conclusions

- **IAXO**: best use of technologies (**magnet**, optics, detectors) and past trajectory of CAST at **CERN**
- **IAXO will probe deep into unexplored axion+ALP parameter space:**
 - QCD axions at the few meV scale → not at reach of any other technique
 - ALPs at the $g_{a\gamma\gamma} \sim 10^{-12} \text{ GeV}^{-1}$ scale
 - ALPs at the $g_{ae} \sim 10^{-13}$ scale
- **IAXO as a generic “axion/ALP facility”**
- **First steps towards TDR after the positive recommendation from CERN SPSC.**
- **Large community now endorsing the project.**
- **Longer-term strategy towards construction under discussion. Site: CERN but also DESY or LNF**
- **Looking forward to Study Group process**




Why to search for axions?

- Most compelling solution to the **Strong CP problem** of the SM
- Axion-like particles (ALPs) **predicted by many extensions** of the SM (e.g. string theory)
- Axions, like WIMPs, may **solve the DM problem for free**. (i.e. not *ad hoc* solution to DM)
- **Astrophysical hints** for axion/ALPs?
 - Transparency of the Universe to UHE gammas
 - Anomalous cooling of different types of star
- Relevant axion/ALP parameter space at **reach of current and near-future experiments**
- Still too little experimental effort devoted to axions when compared to WIMPs


An ALP with $g_{ay} \sim 10^{-11-12} \text{ GeV}^{-1}$

- Well beyond current upper bounds on the g_{ay} coupling (CAST & HB stars $\sim 10^{-10} \text{ GeV}^{-1}$)
- String theory ALPs
- Invoked to explain the anomalous transparency of the Universe to UHE gammas
- Could explain some anomalous stellar cooling observations



The multi-meV axion

- Is compatible with all current axion bounds.
- Is invoked in several anomalous stellar cooling scenarios
 - Gianotti et al. JCAP1605 (2016) 057 [arXiv:1512.08108]
- Can be the cold DM in some models
 - with $NDW > 1$ and bias term to break the discrete symmetry (Kawasaki et al. PRD91 (2015), Ringwald et al. arXiv:1512.06436)
 - or it can be a subdominant DM component
- Very hard to detect (\rightarrow IAXO!)
- SN axion background
 - Raffelt et al. PRD84 (11)




Astrophysical hints for axions

- Gama ray telescopes like MAGIC or HESS observe HE photons from very distant sources...





ALP: $g_{a\gamma} \sim 10^{-12} - 10^{-10} \text{ GeV}^{-1}$
 $m_a \lesssim 10^{-(10-7)} \text{ eV}$



Astrophysical hints for axions (II)

- Most stellar systems seem to cool down faster than expected.
- Presence of axions/ALPs offer a good joint explanation (Giannotti et al. JCAP05(2016)057 [arXiv:1512.08108])
- Parameters at reach of IAXO





Buffer gas for higher masses

Coherence condition ($qL \ll 1$) is recovered for a narrow mass range around m_γ

$$|q| = \frac{m_a^2 - m_\gamma^2}{2E}$$


$$m_\gamma \approx \sqrt{\frac{4\pi\alpha N_e}{m_e}} = 28.9 \sqrt{\frac{Z}{A} \rho} \text{ eV}$$

N_e : number of electrons/cm³
 ρ : gas density (g/cm³)




CAST results (solar axions)

2003 – 2004	CAST phase I <ul style="list-style-type: none"> • vacuum in the magnet bores
2006	CAST phase II - ^4He Run <ul style="list-style-type: none"> • axion masses explored up to 0.39 eV (160 P-steps)
2007	^3He Gas system implementation
2008 - 2011	CAST phase II - ^3He Run <ul style="list-style-type: none"> • axion masses explored up to 1.17 eV • bridging the dark matter limit
2012	<ul style="list-style-type: none"> • Revisit ^4He Run with improved detectors
2013-2015	<ul style="list-style-type: none"> • Revisit vacuum phase with improved detectors • Analisis ongoing. • New result soon available



IAXO sensitivity prospects



Astrophysical hints for ALPs

Much larger QCD axion region explored

CERN SPSC recommendations

SPSC Draft minutes [Jan 2014]

The Committee **recognises** the physics motivation of an International Axion Observatory as described in the Letter of Intent SPSC-I-242, and considers that the proposed setup makes appropriate use of state-of-the-art technologies i.e. magnets, x-ray optics and low-background detectors.

The Committee **encourages** the collaboration to take the next steps towards a **Technical Design Report**.

The Committee recommends that, in the process of preparing the TDR, the possibility to **extend the physics reach** with additional detectors compared to the baseline goal should be investigated. The collaboration should be further strengthened.

Considering the required funding, the **SPSC recommends** that the R&D for the TDR should be pursued within an MOU involving all interested parties.


This was endorsed by the Research Board in March 2014

Minutes of the 206th CERN Research Board held on March 2014:

<https://cds.cern.ch/record/1695812/files/M-207.pdf>

Mid term plans (+4 years) towards IAXO construction

- **Site:**
 - CERN is Plan A, but alternatives under consideration: it may affect funding strategies. Some preliminary interest from a number of institutions.
- **Magnet:**
 - Magnet construction → main resources challenge
 - Plan A: Form international consortium, coordinated by CERN (+CEA?) to collectively support the magnet construction, by means of inkind contributions to the effort.
 - Plan B: Large investment by host institution (IA XO being hosted outside CERN)
- **X-ray optics:**
 - Construction efforts distributed among groups pushing x-ray technologies for IAXO: US (LLNL, UC, MIT) + Europe (DTU, INAF-Milano)
- **Detectors:**
 - Efforts distributed among groups pushing each detector technology (actual decisions taken after TDR):
 - Micromegas: Zaragoza, CEA
 - Ingrid: Bonn
 - CCD: FNAL?, LPNHE?
 - MMCs/TES: Heidelberg, Orsay, CEA



IAXO costs


Item	Cost (MCHF)	Subtotals (MCHF)
Magnet		31.3
Eight coils based assembled toroid	28	
Magnet services	3.3	
Optics		16.0
Prototype Optic: Design, Fabrication, Calibration, Analysis	1.0	
IAXO telescopes (8 + 1 spare)	8.0	
Calibration	2.0	
Integration and alignment	5.0	
Detectors		5.8
Shielding & mechanics	2.1	
Readouts, DAQ electronics & computing	0.8	
Calibration systems	1.5	
Gas & vacuum	1.4	
Dome, base, services building and integration		3.7
Sum		56.8

Table 5: Estimated costs of the IAXO setup: magnet, optics and detectors. It does not include laboratory engineering, as well as maintenance & operation and physics exploitation of the experiment.

Comments/caveats:

- **Costs are for construction, and do not include operations and science support**
- **Costs based on initial estimates that need to be confirmed at TDR**
- **Labor for engineering, maintenance & operations not included**
- **Estimates do not include contingencies**

IAXO timeline



Years		1				2				3				4				5				6			
Months		3	6	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51	54	57	60	63	66	69	72
Magnet																									
Design	T0																								
	T1-T8																								
Demo coil																									
Production																									
Integration																									
Services																									
Optics																									
Optic design study																									
Prototype construction																									
Calibration																									
Finalize design																									
Build assembly machines																									
Procure mandrels & ovens																									
Build coating facilities																									
Slump glass																									
Deposit coatings																									
Assemble optics																									
Calibrate optics																									
Installation																									
Detectors																									
Prototype																									
Construction (incl. spares)																									
Installation & commissioning																									