



Contribution ID: 241

Type: **Poster submission**

## The ILC as a natural SUSY discovery machine and precision microscope: From light higgsinos to tests of unification

*Thursday, August 8, 2019 10:40 AM (20 minutes)*

### Summary

The requirement of electroweak naturalness in simple supersymmetric models motivates the existence of a cluster of four light higgsinos with mass 100-300 GeV, the lighter the better. While such light compressed spectra may be challenging to observe at LHC, future e+e- colliders with  $\sqrt{s} > 2m(\text{higgsino})$  would serve as both a SUSY discovery machine and a precision microscope.

We study higgsino pair production signatures at the ILC based on full, Geant4-based simulation of the ILD detector concept. We examine several benchmark scenarios that may or may not be accessible to HL-LHC searches, with mass differences between the higgsino states between 20 and 4 GeV. Assuming  $\sqrt{s} = 500$  GeV and  $1000 \text{ fb}^{-1}$  of integrated luminosity, the individual higgsino masses can be measured to 1-2% precision in case of the larger mass differences, and still at the level of 5% for the smallest mass difference case. The higgsino mass splittings are sensitive to the electroweak gaugino masses and can allow extraction of gaugino masses to  $\pm 3 - 20\%$  (depending on the model).

Extrapolation of gaugino masses via renormalization group running can test the hypothesis of gaugino mass unification. We also examine a case with natural generalized mirage mediation where the unification of gaugino masses at an intermediate scale apparently gives rise to a natural SUSY spectrum somewhat beyond the reach of HL-LHC.

**Primary author:** LIST, Jenny (Deutsches Elektronen-Synchrotron (DE))

**Co-author:** KAWAGOE, Kiyotomo (Kyushu University (JP))

**Presenter:** BERGGREN, Mikael (Deutsches Elektronen-Synchrotron (DE))

**Session Classification:** Poster Session (Thu/Fri)

**Track Classification:** Collider – Beyond Standard Model Searches