

26th International Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY2018)



Contribution ID: 208

Type: **Talk (closed)**

Can we discover a light singlet-like NMSSM Higgs boson at the LHC?

Monday 23 July 2018 15:00 (20 minutes)

In the next-to minimal supersymmetric standard model (NMSSM) an additional singlet-like Higgs boson with small couplings to the standard model (SM) particles is introduced. Although the mass can be well below the discovered 125 GeV Higgs boson its small couplings may make a discovery at the LHC difficult.

We use a novel scanning technique to efficiently scan the whole parameter space and determine the range of cross sections and branching ratios for a light singlet-like Higgs boson below 125 GeV.

This allows to determine the perspectives for the future discovery potential at the LHC. Specific LHC benchmark points are proposed. A discovery of such a light Higgs singlet would strongly point to a singlino as a dark matter candidate.

Parallel Session

Electroweak, Top and Higgs Physics

Authors: BESKIDT, Conny Renate (KIT - Karlsruhe Institute of Technology (DE)); DE BOER, Wim (KIT - Karlsruhe Institute of Technology (DE)); KAZAKOV, Dmitri (JINR/ITEP)

Presenter: BESKIDT, Conny Renate (KIT - Karlsruhe Institute of Technology (DE))

Session Classification: Electroweak, Top and Higgs Physics