

Small- q_T factorization and the soft function for top pair production at NNLO

Sebastian Sapeta

IFJ PAN Kraków


In collaboration with René Ángeles-Martinez and Michał Czakon

based on JHEP 1810 (2018) 201



Top pair production: the status of QCD calculations


- ▶ A single *complete* NNLO result for total and differential cross section obtained with STRIPPER methodology [Czakon, Fiedler, Mitov '13; Czakon, Heymes, Mitov '16]



- ▶ Flavour off-diagonal channels at NNLO from q_T subtraction [Bonciani, Catani, Grazzini, Sargsyan, Torre '15]
- ▶ Approximate NNLO [Broggio, Papanastasiou, Signer '14] and N³LO [Kidonakis '14]
- ▶ Soft and small-mass resummation at NNLL [Czakon, Ferroglia, Heymes, Mitov, Pecjak, Scott, Wang, Yang '18]
- ▶ Small- q_T resummation at NNLL [Li, Li, Shao, Yang, Zhu '13; Catani, Grazzini, Torre '14]

Top pair production: the status of QCD calculations

- ▶ A single *complete* NNLO result for total and differential cross section obtained with STRIPPER methodology [Czakon, Fiedler, Mitov '13; Czakon, Heymes, Mitov '16]



- ▶ Given the complexity of the calculation, a second result obtained with an *independent* method is highly desirable

- ▶ Flavour off-diagonal channels at NNLO from q_T subtraction [Bonciani, Catani, Grazzini, Sargsyan, Torre '15]
- ▶ Approximate NNLO [Broggio, Papanastasiou, Signer '14] and N³LO [Kidonakis '14]
- ▶ Soft and small-mass resummation at NNLL [Czakon, Ferroglia, Heymes, Mitov, Pecjak, Scott, Wang, Yang '18]
- ▶ Small- q_T resummation at NNLL [Li, Li, Shao, Yang, Zhu '13; Catani, Grazzini, Torre '14]

The q_T slicing method

[Catani, Grazzini '07, '15]

$$p + p \rightarrow F(q_T) + X$$

$$\sigma_{N^m\text{LO}}^F = \int_0^{q_{T,\text{cut}}} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T} + \int_{q_{T,\text{cut}}}^{\infty} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T}$$

The q_T slicing method

[Catani, Grazzini '07, '15]

$$p + p \rightarrow F(q_T) + X$$

$$\begin{aligned}\sigma_{N^m\text{LO}}^F &= \int_0^{q_{T,\text{cut}}} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T} + \int_{q_{T,\text{cut}}}^{\infty} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T} \\ &= \int_0^{q_{T,\text{cut}}} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T} + \int_{q_{T,\text{cut}}}^{\infty} dq_T \frac{d\sigma_{N^{m-1}\text{LO}}^{F+\text{jet}}}{dq_T}\end{aligned}$$

The q_T slicing method

[Catani, Grazzini '07, '15]

$$p + p \rightarrow F(q_T) + X$$

$$\begin{aligned}\sigma_{N^m\text{LO}}^F &= \int_0^{q_{T,\text{cut}}} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T} + \int_{q_{T,\text{cut}}}^{\infty} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T} \\ &= \int_0^{q_{T,\text{cut}}} dq_T \frac{d\sigma_{N^m\text{LO}}^F}{dq_T} + \int_{q_{T,\text{cut}}}^{\infty} dq_T \frac{d\sigma_{N^{m-1}\text{LO}}^{F+\text{jet}}}{dq_T}\end{aligned}$$

enough to know in
small- q_T approximation



known

Soft Collinear Effective Theory (SCET)

$$\text{SCET} \simeq \text{QCD} \Big|_{\text{IR limit}}$$

- ▶ Hard degrees of freedom are integrated out into Wilson coefficients, which are then used to adjust new couplings of the (effective) theory.

Soft Collinear Effective Theory (SCET)

$$\text{SCET} \simeq \text{QCD} \Big|_{\text{IR limit}}$$

- ▶ Hard degrees of freedom are integrated out into Wilson coefficients, which are then used to adjust new couplings of the (effective) theory.

QCD fields written as sums of collinear, anti-collinear and soft components:

$$\phi(x) = \phi_c(x) + \phi_{\bar{c}}(x) + \phi_s(x)$$

Soft Collinear Effective Theory (SCET)

$$\text{SCET} \simeq \text{QCD} \Big|_{\text{IR limit}}$$

- ▶ Hard degrees of freedom are integrated out into Wilson coefficients, which are then used to adjust new couplings of the (effective) theory.

QCD fields written as sums of collinear, anti-collinear and soft components:


$$\phi(x) = \phi_c(x) + \phi_{\bar{c}}(x) + \phi_s(x)$$

The new fields decouple in the Lagrangian

$$\mathcal{L}_{\text{SCET}} = \mathcal{L}_c + \mathcal{L}_{\bar{c}} + \mathcal{L}_s$$


- ▶ The separation of fields in the Lagrangian into collinear, anti-collinear and soft sectors, facilitates proofs of factorization theorems

Small- q_T factorization in SCET



where $F = H, Z, W, ZZ, WW, t\bar{t}, \dots$

Small- q_T factorization in SCET



where $F = H, Z, W, ZZ, WW, t\bar{t}, \dots$

$$\frac{d\sigma^F}{d\Phi} = \mathcal{B}_1 \otimes \mathcal{B}_2 \otimes \mathcal{H} \otimes \mathcal{S} + \mathcal{O}\left(\frac{q_T^2}{q^2}\right)$$

Small- q_T factorization in SCET

Gluons' momenta in light-cone coordinates

$$k_i^\mu = (k_i^+, k_i^-, \mathbf{k}_i^\perp) \quad \text{where} \quad k^\pm = k^0 \pm k^3$$

Expansion parameter

$$\lambda = \frac{q_T^2}{q^2} \ll 1$$

Small- q_T factorization in SCET

Gluons' momenta in light-cone coordinates

$$k_i^\mu = (k_i^+, k_i^-, \mathbf{k}_i^\perp) \quad \text{where} \quad k^\pm = k^0 \pm k^3$$

Expansion parameter

$$\lambda = \frac{q_T^2}{q^2} \ll 1$$


Regions

collinear $k_i^\mu \sim (1, \lambda^2, \lambda) Q^2 \quad \mathcal{B}_1$

anti-collinear $k_i^\mu \sim (\lambda^2, 1, \lambda) Q^2 \quad \mathcal{B}_2$

hard $k_i^\mu \sim (1, 1, 1) Q^2 \quad \mathcal{H}$

soft $k_i^\mu \sim (\lambda, \lambda, \lambda) Q^2 \quad \mathcal{S}$



Top pair production at small- q_T through NNLO

$$\frac{d\sigma^{\text{NNLO}}}{dq_T dy dM d\cos\theta} = \sum_{i,\bar{i}} \mathcal{B}_{i/h_1} \otimes \mathcal{B}_{\bar{i}/h_2} \otimes \text{Tr}[\mathcal{H}_{i\bar{i}} \otimes \mathcal{S}_{i\bar{i}}]$$

where

- q_T, y, M : transverse momentum, rapidity, mass of top quark pair
- θ : scattering angle of the top quark in $t\bar{t}$ rest frame

Top pair production at small- q_T through NNLO

$$\frac{d\sigma^{\text{NNLO}}}{dq_T dy dM d\cos\theta} = \sum_{i,\bar{i}} \mathcal{B}_{i/h_1} \otimes \mathcal{B}_{\bar{i}/h_2} \otimes \text{Tr}[\mathcal{H}_{i\bar{i}} \otimes \mathcal{S}_{i\bar{i}}]$$

where

q_T, y, M : transverse momentum, rapidity, mass of top quark pair
 θ : scattering angle of the top quark in $t\bar{t}$ rest frame

\mathcal{B} - known up to NNLO [Gehrmann, Lübbert, Yang '12, '14]

\mathcal{H} - known up to NNLO [Czakon '08; Baernreuther, Czakon, Fiedler '13]

\mathcal{S} - known up to NLO in small- q_T limit [Li, Li, Shao, Yan, Zhu '13; Catani, Grazzini, Torre '14] (and up to NNLO in the threshold limit [Ferroglia, Pecjak, Yang '12; Wang, Xu, Yang and Zhu '18])

Top pair production at small- q_T through NNLO

$$\frac{d\sigma^{\text{NNLO}}}{dq_T dy dM d\cos\theta} = \sum_{i,\bar{i}} \mathcal{B}_{i/h_1} \otimes \mathcal{B}_{\bar{i}/h_2} \otimes \text{Tr}[\mathcal{H}_{i\bar{i}} \otimes \mathcal{S}_{i\bar{i}}]$$



where

q_T, y, M : transverse momentum, rapidity, mass of top quark pair
 θ : scattering angle of the top quark in $t\bar{t}$ rest frame


- \mathcal{B} - known up to NNLO [Gehrmann, Lübbert, Yang '12, '14]
- \mathcal{H} - known up to NNLO [Czakon '08; Baernreuther, Czakon, Fiedler '13]
- \mathcal{S} - known up to NLO in small- q_T limit [Li, Li, Shao, Yan, Zhu '13; Catani, Grazzini, Torre '14] (and up to NNLO in the threshold limit [Ferrogli, Pecjak, Yang '12; Wang, Xu, Yang and Zhu '18])

Calculating the missing NNLO correction to the soft function in the small- q_T limit, \mathcal{S} , is the aim of this phase of our work.

Rapidity divergences and analytic regulator



Rapidity divergences and analytic regulator



Modification of the measure [Becher, Bell '12]

$$\int d^d k \delta^+(k^2) \rightarrow \int d^d k \left(\frac{\nu}{k_+} \right)^\alpha \delta^+(k^2)$$

- ▶ The regulator is necessary at intermediate steps of the calculation.
- ▶ Rapidity divergences do not appear in QCD, hence, the complete SCET result has to stay finite in the limit $\alpha \rightarrow 0$.

Soft function

- Represents corrections coming from exchanges of **real, soft gluons**, whose transverse momenta sum up to a fixed value q_T .

$$S_{\text{bare}}(q_T, \beta_t, \theta) \propto \sum \text{Diagram} \delta(q_T - |\sum_i k_{i\perp}|) \prod_i \delta^+(k_i^2)$$

Soft function

- Represents corrections coming from exchanges of **real, soft gluons**, whose transverse momenta sum up to a fixed value q_T .

$$\beta_t = \sqrt{1 - \frac{4m_t^2}{M^2}}$$

$$S_{\text{bare}}(q_T, \beta_t, \theta) \propto \sum \text{Diagram} \delta(q_T - |\sum_i k_{i\perp}|) \prod_i \delta^+(k_i^2)$$

Soft function

- Represents corrections coming from exchanges of **real, soft gluons**, whose transverse momenta sum up to a fixed value q_T .

$$S_{\text{bare}}(q_T, \beta_t, \theta) \propto \sum \text{Diagram} \delta(q_T - |\sum_i k_{i\perp}|) \prod_i \delta^+(k_i^2)$$

$$\beta_t = \sqrt{1 - \frac{4m_t^2}{M^2}}$$

- external momenta \rightarrow Wilson Lines along n, \bar{n}, v_3, v_4 (Born kinematics)

Soft function

- Represents corrections coming from exchanges of **real, soft gluons**, whose transverse momenta sum up to a fixed value q_T .

$$\beta_t = \sqrt{1 - \frac{4m_t^2}{M^2}}$$

$$S_{\text{bare}}(q_T, \beta_t, \theta) \propto \sum \text{Diagram} \delta(q_T - |\sum_i k_{i\perp}|) \prod_i \delta^+(k_i^2)$$

- external momenta \rightarrow Wilson Lines along n, \bar{n}, v_3, v_4 (Born kinematics)


$$\mathbf{S}_{i\bar{i}} = \sum_{n=0}^{\infty} \mathbf{S}_{i\bar{i}}^{(n)} \left(\frac{\alpha_s}{4\pi}\right)^n$$

$$\mathbf{S}_{i\bar{i}}^{(n)} = \sum_{\{j\}} \mathbf{w}_{\{j\}}^{i\bar{i}} I_{\{j\}}$$

colour matrices \uparrow \uparrow phase space integrals

Renormalization

separately divergent


$$\left[\frac{d\sigma}{d\Phi} \right] = \mathcal{B}_1^{(\text{bare})} \otimes \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} \left[\mathcal{H}^{(\text{bare})} \otimes \mathcal{S}^{(\text{bare})} \right]$$

finite

Renormalization

separately divergent



$$\begin{aligned} \text{finite} \quad \left[\frac{d\sigma}{d\Phi} \right] &= \mathcal{B}_1^{(\text{bare})} \otimes \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} \left[\mathcal{H}^{(\text{bare})} \otimes \mathcal{S}^{(\text{bare})} \right] \\ &= Z_B \mathcal{B}_1^{(\text{bare})} \otimes Z_B \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} \left[\mathbf{Z}_H^\dagger \mathcal{H}^{(\text{bare})} \mathbf{Z}_H \otimes \mathbf{Z}_S^\dagger \mathcal{S}^{(\text{bare})} \mathbf{Z}_S \right] \end{aligned}$$

Renormalization

separately divergent


$$\begin{aligned} \left. \begin{array}{l} \text{finite} \\ \rightarrow \end{array} \right\} \frac{d\sigma}{d\Phi} &= \mathcal{B}_1^{(\text{bare})} \otimes \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} \left[\mathcal{H}^{(\text{bare})} \otimes \mathcal{S}^{(\text{bare})} \right] \\ &= Z_B \mathcal{B}_1^{(\text{bare})} \otimes Z_B \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} \left[\mathbf{Z}_H^\dagger \mathcal{H}^{(\text{bare})} \mathbf{Z}_H \otimes \mathbf{Z}_S^\dagger \mathcal{S}^{(\text{bare})} \mathbf{Z}_S \right] \\ &= \mathcal{B}_1(\mu) \otimes \mathcal{B}_2(\mu) \otimes \text{Tr} \left[\mathcal{H}(\mu) \otimes \mathcal{S}(\mu) \right] \end{aligned}$$

Renormalization



separately divergent

$$\begin{aligned} \lrcorner \rightarrow \frac{d\sigma}{d\Phi} &= \mathcal{B}_1^{(\text{bare})} \otimes \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} [\mathcal{H}^{(\text{bare})} \otimes \mathcal{S}^{(\text{bare})}] \\ \text{finite} &= Z_B \mathcal{B}_1^{(\text{bare})} \otimes Z_B \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} [\mathbf{Z}_H^\dagger \mathcal{H}^{(\text{bare})} \mathbf{Z}_H \otimes \mathbf{Z}_S^\dagger \mathcal{S}^{(\text{bare})} \mathbf{Z}_S] \\ &= \mathcal{B}_1(\mu) \otimes \mathcal{B}_2(\mu) \otimes \text{Tr} [\mathcal{H}(\mu) \otimes \mathcal{S}(\mu)] \end{aligned}$$

separately finite

Renormalization

separately divergent

$$\begin{aligned} \Gamma \rightarrow \frac{d\sigma}{d\Phi} &= \mathcal{B}_1^{(\text{bare})} \otimes \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} [\mathcal{H}^{(\text{bare})} \otimes \mathcal{S}^{(\text{bare})}] \\ \text{finite} &= Z_B \mathcal{B}_1^{(\text{bare})} \otimes Z_B \mathcal{B}_2^{(\text{bare})} \otimes \text{Tr} [\mathbf{Z}_H^\dagger \mathcal{H}^{(\text{bare})} \mathbf{Z}_H \otimes \mathbf{Z}_S^\dagger \mathcal{S}^{(\text{bare})} \mathbf{Z}_S] \\ &= \mathcal{B}_1(\mu) \otimes \mathcal{B}_2(\mu) \otimes \text{Tr} [\mathcal{H}(\mu) \otimes \mathcal{S}(\mu)] \end{aligned}$$

separately finite

$$\frac{d}{d\mu} \frac{d\sigma}{d\Phi} = 0 \quad \rightarrow \quad \text{Renormalization Group Equations for } \mathcal{B}, \mathcal{H} \text{ and } \mathcal{S}$$

Renormalization

- ▶ RG equation

$$\frac{d}{d \ln \mu} \mathbf{S}_{i\bar{i}}(\mu) = -\gamma_{i\bar{i}}^{s\dagger} \mathbf{S}_{i\bar{i}}(\mu) - \mathbf{S}_{i\bar{i}}(\mu) \gamma_{i\bar{i}}^s$$

- ▶ Soft anomalous dimension

$$\gamma^s = -\mathbf{Z}_s^{-1} \frac{d\mathbf{Z}_s}{d \ln \mu}$$

Renormalization

- ▶ RG equation

$$\frac{d}{d \ln \mu} \mathbf{S}_{i\bar{i}}(\mu) = -\gamma_{i\bar{i}}^{s\dagger} \mathbf{S}_{i\bar{i}}(\mu) - \mathbf{S}_{i\bar{i}}(\mu) \gamma_{i\bar{i}}^s$$


- ▶ Soft anomalous dimension

$$\gamma^s = -\mathbf{Z}_s^{-1} \frac{d\mathbf{Z}_s}{d \ln \mu}$$


Specifically, at the order α_s^2 , we get

$$\underbrace{\mathbf{S}^{(2)}}_{\text{finite part only}} = \overbrace{\mathbf{Z}_s^{\dagger(2)} \mathbf{S}_{\text{bare}}^{(0)} + \mathbf{S}_{\text{bare}}^{(0)} \mathbf{Z}_s^{(2)} + \mathbf{Z}_s^{\dagger(1)} \mathbf{S}_{\text{bare}}^{(0)} \mathbf{Z}_s^{(1)}}^{\text{pole part only}} + \underbrace{\mathbf{Z}_s^{\dagger(1)} \mathbf{S}_{\text{bare}}^{(1)} + \mathbf{S}_{\text{bare}}^{(1)} \mathbf{Z}_s^{(1)} + \mathbf{S}_{\text{bare}}^{(2)} - \frac{\beta_0}{\epsilon} \mathbf{S}_{\text{bare}}^{(1)}}_{\text{finite + pole part}}$$

Soft function at NLO



Soft function at NLO



- Known in analytic form

[Li, Li, Shao, Yan, Zhu '13; Catani, Grazzini, Torre '13]


$$L_{\perp} = \ln \frac{x_T^2 \mu^2}{4e^{-2\gamma_E}}$$

$$\begin{aligned} \mathbf{S}_{ii}^{(1)} = & 4L_{\perp} \left(2\mathbf{w}_{ii}^{13} \ln \frac{-t_1}{m_t M} + 2\mathbf{w}_{ii}^{23} \ln \frac{-u_1}{m_t M} + \mathbf{w}_{ii}^{33} \right) \\ & - 4 \left(\mathbf{w}_{ii}^{13} + \mathbf{w}_{ii}^{23} \right) \text{Li}_2 \left(1 - \frac{t_1 u_1}{m_t^2 M^2} \right) + 4\mathbf{w}_{ii}^{33} \ln \frac{t_1 u_1}{m_t^2 M^2} \\ & - 2\mathbf{w}_{ii}^{34} \frac{1 + \beta_t^2}{\beta_t} \left[L_{\perp} \ln x_s - \text{Li}_2 \left(-x_s \text{tg}^2 \frac{\theta}{2} \right) + \text{Li}_2 \left(-\frac{1}{x_s} \text{tg}^2 \frac{\theta}{2} \right) \right. \\ & \left. + 4 \ln x_s \ln \cos \frac{\theta}{2} \right] + \mathcal{O}(\epsilon) \end{aligned}$$

Soft function at NNLO

Three distinct groups of diagrams:


► Bubble




Soft function at NNLO

Three distinct groups of diagrams:

► Bubble




► Single-cut




Soft function at NNLO

Three distinct groups of diagrams:



► Bubble



► Single-cut



► Double-cut




Soft function at NNLO

Three distinct groups of diagrams:


► Bubble

DIFFERENTIAL EQUATIONS

► Double-cut



► Single-cut



Soft function at NNLO

Three distinct groups of diagrams:



► Bubble

► Single-cut

**DIFFERENTIAL
EQUATIONS**

**DIRECT
INTEGRATION**

► Double-cut



Soft function at NNLO

Three distinct groups of diagrams:

▶ Bubble

▶ Single-cut

**DIFFERENTIAL
EQUATIONS**

**DIRECT
INTEGRATION**

▶ Double-cut

SECTOR DECOMPOSITION

Double-cut NNLO integrals

Example:

$$\tilde{I}_{3gV,ij} = \int \frac{d^d k_1 d^d k_2 \delta^+(k_1^2) \delta^+(k_2^2) \delta((k_1 + k_2)_T^2 - q_T^2)}{(n \cdot k_1)^\alpha (n \cdot k_2)^\alpha (n_i \cdot k_1) (n_j \cdot (k_1 + k_2)) (k_1 + k_2)^2}$$

Double-cut NNLO integrals

Example:

$$\tilde{I}_{3g\nu,ij} = \int \frac{d^d k_1 d^d k_2 \delta^+(k_1^2) \delta^+(k_2^2) \delta((k_1 + k_2)_T^2 - q_T^2)}{(n \cdot k_1)^\alpha (n \cdot k_2)^\alpha (n_i \cdot k_1) (n_j \cdot (k_1 + k_2)) (k_1 + k_2)^2}$$

- ▶ divergent in the limits $\epsilon \rightarrow 0$ and $\alpha \rightarrow 0$
- ▶ a range of overlapping singularities
- ▶ complication introduced by $\delta((k_1 + k_2)_T^2 - q_T^2)$ which additionally couples gluon's momenta

Double-cut NNLO integrals

Example:

$$\tilde{I}_{3g\nu,ij} = \int \frac{d^d k_1 d^d k_2 \delta^+(k_1^2) \delta^+(k_2^2) \delta((k_1 + k_2)_T^2 - q_T^2)}{(n \cdot k_1)^\alpha (n \cdot k_2)^\alpha (n_i \cdot k_1) (n_j \cdot (k_1 + k_2)) (k_1 + k_2)^2}$$

- ▶ divergent in the limits $\epsilon \rightarrow 0$ and $\alpha \rightarrow 0$
- ▶ a range of overlapping singularities
- ▶ complication introduced by $\delta((k_1 + k_2)_T^2 - q_T^2)$ which additionally couples gluon's momenta

To disentangle overlapping singularities and calculate regularized integrals we use the method of **sector decomposition** [Binoth, Heinrich, '00; Borowka, Heinrich, Jahn, Jones, Kerner, Schlenk, Zirke '17].

Sector decomposition

$$\int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x + y)^{2+\epsilon}}$$

Sector decomposition

$$\int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x + y)^{2+\epsilon}} = \int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x + y)^{2+\epsilon}} \left[\overbrace{\Theta(x - y)}^{(1)} + \overbrace{\Theta(y - x)}^{(2)} \right]$$

Sector decomposition

$$\int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x+y)^{2+\epsilon}} = \int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x+y)^{2+\epsilon}} \left[\overbrace{\Theta(x-y)}^{(1)} + \overbrace{\Theta(y-x)}^{(2)} \right]$$

$$(1) \quad y = x t \qquad (2) \quad x = y t$$

Sector decomposition

$$\int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x + y)^{2+\epsilon}} = \int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x + y)^{2+\epsilon}} \left[\overbrace{\Theta(x - y)}^{(1)} + \overbrace{\Theta(y - x)}^{(2)} \right]$$

.....

$$(1) \quad y = x t \qquad (2) \quad x = y t$$

.....

$$= \int_0^1 dx dt \frac{\mathcal{W}(x, tx)}{(1 + t)^{2+\epsilon} x^{1+\epsilon}} + \int_0^1 dt dy \frac{\mathcal{W}(ty, y)}{(1 + t)^{2+\epsilon} y^{1+\epsilon}}$$

Sector decomposition


$$\int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x + y)^{2+\epsilon}} = \int_0^1 dx dy \frac{\mathcal{W}(x, y)}{(x + y)^{2+\epsilon}} \left[\overbrace{\Theta(x - y)}^{(1)} + \overbrace{\Theta(y - x)}^{(2)} \right]$$

.....

$$(1) \quad y = x t \qquad (2) \quad x = y t$$

.....

$$= \int_0^1 dx dt \frac{\mathcal{W}(x, tx)}{(1+t)^{2+\epsilon} x^{1+\epsilon}} + \int_0^1 dt dy \frac{\mathcal{W}(ty, y)}{(1+t)^{2+\epsilon} y^{1+\epsilon}}$$



Sector decomposition

In general, each integral can be expressed as

$$\mathcal{I} = \sum_{i \in \text{sectors}} \int_0^1 \frac{dx_1}{x_1^{1+a_1\epsilon}} \frac{dx_2}{x_2^{1+a_2\epsilon}} \cdots \frac{dx_n}{x_n^{1+a_n\epsilon}} \mathcal{W}_i(x_1, x_2, \dots, x_n)$$

Sector decomposition

In general, each integral can be expressed as

$$\mathcal{I} = \sum_{i \in \text{sectors}} \int_0^1 \frac{dx_1}{x_1^{1+a_1\epsilon}} \frac{dx_2}{x_2^{1+a_2\epsilon}} \cdots \frac{dx_n}{x_n^{1+a_n\epsilon}} \mathcal{W}_i(x_1, x_2, \dots, x_n)$$

and then we use

$$\frac{1}{x_i^{1+a_i\epsilon}} = -\frac{1}{a_i\epsilon} \delta(x_i) + \sum_{n=0}^{\infty} \frac{a_i^n \epsilon^n}{n!} \left[\frac{\log^n(x_i)}{x_i} \right]_+$$

Sector decomposition

In general, each integral can be expressed as

$$\mathcal{I} = \sum_{i \in \text{sectors}} \int_0^1 \frac{dx_1}{x_1^{1+a_1\epsilon}} \frac{dx_2}{x_2^{1+a_2\epsilon}} \cdots \frac{dx_n}{x_n^{1+a_n\epsilon}} \mathcal{W}_i(x_1, x_2, \dots, x_n)$$

and then we use

$$\frac{1}{x_i^{1+a_i\epsilon}} = -\frac{1}{a_i\epsilon} \delta(x_i) + \sum_{n=0}^{\infty} \frac{a_i^n \epsilon^n}{n!} \left[\frac{\log^n(x_i)}{x_i} \right]_+$$

After the above procedure is performed, all divergences become explicit and are turned in to ϵ poles

Sector decomposition

Two types of singularities

- ▶ Endpoint, e.g. soft:

$$(k_1^+, k_1^-, k_1^\perp) \rightarrow 0$$

Sector decomposition


Two types of singularities

- ▶ Endpoint, e.g. soft:

$$(k_1^+, k_1^-, k_1^\perp) \rightarrow 0$$

- ▶ Manifold, e.g. collinear

$$k_1 \cdot k_2 \rightarrow 0$$



The strategy

Given the integral:

$$I_G = \int d^d k_1 d^d k_2 \mathcal{I}_G \times \mathcal{W}_G$$

The strategy

Given the integral:

boundary integral encodes all $\alpha, \epsilon \rightarrow 0$ singularities

$$I_G = \int d^d k_1 d^d k_2 \mathcal{I}_G \times \mathcal{W}_G$$

↑
finite weight

The strategy

Given the integral:

boundary integral encodes all $\alpha, \epsilon \rightarrow 0$ singularities

$$I_G = \int d^d k_1 d^d k_2 \mathcal{I}_G \times \mathcal{W}_G$$

↑
finite weight

- ▶ Analytically integrate 3 out of $2d$ dimensions
- ▶ Map the remaining variables to a unit hypercube (split the original integral into a sum if necessary)

$$= \sum_j \int_0^1 \prod_{i=1}^{2d-3} dx_i (\mathcal{I}_G \times \mathcal{W}_G)_j$$

The strategy

Given the integral:

boundary integral encodes all $\alpha, \epsilon \rightarrow 0$ singularities

$$I_G = \int d^d k_1 d^d k_2 \mathcal{I}_G \times \mathcal{W}_G$$

↑
finite weight

- ▶ Analytically integrate 3 out of $2d$ dimensions
- ▶ Map the remaining variables to a unit hypercube (split the original integral into a sum if necessary)
- ▶ Apply sector decomposition to disentangle overlapping singularities

$$= \sum_j \int_0^1 \prod_{i=1}^{2d-3} dx_i (\mathcal{I}_G \times \mathcal{W}_G)_j$$

$$= \sum_j \sum_{k \in \text{sectors}} \int_0^1 \prod_{i=1}^{2d-3} dx_i (\mathcal{I}_G \times \mathcal{W}_G)_{jk}$$

The strategy

Given the integral:

boundary integral encodes all $\alpha, \epsilon \rightarrow 0$ singularities

$$I_G = \int d^d k_1 d^d k_2 \mathcal{I}_G \times \mathcal{W}_G$$

↑
finite weight

- ▶ Analytically integrate 3 out of $2d$ dimensions
- ▶ Map the remaining variables to a unit hypercube (split the original integral into a sum if necessary)
- ▶ Apply sector decomposition to disentangle overlapping singularities
- ▶ Expand the result in Laurent series in ϵ and α

$$= \sum_j \int_0^1 \prod_{i=1}^{2d-3} dx_i (\mathcal{I}_G \times \mathcal{W}_G)_j$$

$$= \sum_j \sum_{k \in \text{sectors}} \int_0^1 \prod_{i=1}^{2d-3} dx_i (\mathcal{I}_G \times \mathcal{W}_G)_{jk}$$

$$= \sum_{j,m,n} \sum_{k \in \text{sectors}} \left(\int_0^1 a_{mn} \times \mathcal{W}_G \right) \alpha^m \epsilon^n$$

The strategy

Given the integral:

boundary integral encodes all $\alpha, \epsilon \rightarrow 0$ singularities

$$I_G = \int d^d k_1 d^d k_2 \mathcal{I}_G \times \mathcal{W}_G$$

↑
finite weight

- ▶ Analytically integrate 3 out of $2d$ dimensions
- ▶ Map the remaining variables to a unit hypercube (split the original integral into a sum if necessary)
- ▶ Apply sector decomposition to disentangle overlapping singularities
- ▶ Expand the result in Laurent series in ϵ and α
- ▶ Numerically integrate series coefficients


$$= \sum_j \int_0^1 \prod_{i=1}^{2d-3} dx_i (\mathcal{I}_G \times \mathcal{W}_G)_j$$

$$= \sum_j \sum_{k \in \text{sectors}} \int_0^1 \prod_{i=1}^{2d-3} dx_i (\mathcal{I}_G \times \mathcal{W}_G)_{jk}$$


$$= \sum_{j,m,n} \sum_{k \in \text{sectors}} \left(\int_0^1 a_{mn} \times \mathcal{W}_G \right) \alpha^m \epsilon^n$$

$$= \sum_{j,m,n} \sum_{k \in \text{sectors}} c_{mn}^{jk} \alpha^m \epsilon^n$$

Single-cut (real-virtual)




Single-cut (real-virtual)



$$S_{1\text{-cut}}^{(2)} = \sum_{ijk} \int d^d l \frac{\delta^+(l^2) \delta(l_T - q_T)}{l_+^\alpha n_k \cdot l} n_k^\mu T_k^a J_{ij,a}^\mu(l)$$


Single-cut (real-virtual)



$$S_{1\text{-cut}}^{(2)} = \sum_{ijk} \int d^d l \frac{\delta^+(l^2) \delta(l_T - q_T)}{l_+^\alpha n_k \cdot l} n_k^\mu T_k^a J_{ij,a}^\mu(l)$$

- ▶ The soft current $J_{ij,a}^\mu(l)$ is known up to NLO [Catani, Grazzini '00; Bierenbaum, Czakon, Mitov '12; Czakon, Mitov '18].


Single-cut (real-virtual)



$$S_{1\text{-cut}}^{(2)} = \sum_{ijk} \int d^d l \frac{\delta^+(l^2) \delta(l_T - q_T)}{l_+^\alpha n_k \cdot l} n_k^\mu T_k^a J_{ij,a}^\mu(l)$$

- ▶ The soft current $J_{ij,a}^\mu(l)$ is known up to NLO [Catani, Grazzini '00; Bierenbaum, Czakon, Mitov '12; Czakon, Mitov '18].
- ▶ $S_{1\text{-cut}}^{(2)}$ can be obtained by a relatively simple integration over l^μ .


Single-cut (real-virtual)




$$S_{1\text{-cut}}^{(2)} = \sum_{ijk} \int d^d l \frac{\delta^+(l^2) \delta(l_T - q_T)}{l_+^\alpha n_k \cdot l} n_k^\mu T_k^a J_{ij,a}^\mu(l)$$

- ▶ The soft current $J_{ij,a}^\mu(l)$ is known up to NLO [Catani, Grazzini '00; Bierenbaum, Czakon, Mitov '12; Czakon, Mitov '18].
- ▶ $S_{1\text{-cut}}^{(2)}$ can be obtained by a relatively simple integration over l^μ .
- ▶ Single-cut piece of the soft function exhibits both real and imaginary part. The latter when $i \neq j \neq k$, the former, otherwise.

Bubble



Bubble




- ▶ Solvable analytically: direct cross check of our sector decomposition-based implementation
- ▶ Non-trivial tensor structure \rightarrow challenging numerators
- ▶ Laboratory to stress-test sector decomposition-based methodology

Bubble part of the soft function from differential equations


$$\propto \int \frac{d^d q \delta(q_T - 1) \theta^+(q^2) n_i^\mu n_j^\nu}{q^4 (n_i \cdot q) (n_j \cdot q)} \left(\text{bubble diagram} \right)_{\mu\nu}$$

Bubble part of the soft function from differential equations



$$\text{Diagram} \propto \int \frac{d^d q \delta(q_T - 1) \theta^+(q^2) n_i^\mu n_j^\nu}{q^4 (n_i \cdot q) (n_j \cdot q)} \left(\text{Bubble Diagram} \right)_{\mu\nu}$$

where



$$\left(\text{Bubble Diagram} \right)_{\mu\nu} = \int \frac{d^d k N_{\mu\nu} \delta^+(k^2) \delta^+((q-k)^2)}{(n \cdot k)^\alpha (n \cdot (q-k))^\alpha k^2 (q-k)^2}$$

$$= T_{00} g^{\mu,\nu} + T_{qq} q^\mu q^\nu + T_{nn} n^\mu n^\nu + T_{qn} (n^\mu q^\nu + q^\mu n^\nu)$$

Bubble part of the soft function from differential equations

- ▶ Reverse unitarity [Anastasiou, Melnikov '02, Cutkosky '60]

$$2i\pi\delta(p^2 - m^2) \rightarrow \frac{1}{p^2 - m^2 + i\epsilon} - \frac{1}{p^2 - m^2 - i\epsilon}$$

Bubble part of the soft function from differential equations

- ▶ Reverse unitarity [Anastasiou, Melnikov '02, Cutkosky '60]

$$2i\pi\delta(p^2 - m^2) \rightarrow \frac{1}{p^2 - m^2 + i\epsilon} - \frac{1}{p^2 - m^2 - i\epsilon}$$

- ▶ real gluons are turned into virtual ones
- ▶ this allows us to utilize all the existing loop technology

Bubble part of the soft function from differential equations

- ▶ Reverse unitarity [Anastasiou, Melnikov '02, Cutkosky '60]

$$2i\pi\delta(p^2 - m^2) \rightarrow \frac{1}{p^2 - m^2 + i\epsilon} - \frac{1}{p^2 - m^2 - i\epsilon}$$

- ▶ real gluons are turned into virtual ones
- ▶ this allows us to utilize all the existing loop technology

- ▶ Topology:

$$\int \frac{d^d k}{(n \cdot k)^{a_1 + 2\alpha} (\bar{n} \cdot k)^{a_2} (v_3 \cdot k)^{a_3} (v_4 \cdot k)^{a_4} (k^2 - m^2)^{a_5} ((n \cdot k)(\bar{n} \cdot k) - m^2 - 1)^{a_6}}$$

Integration by parts and differential equations

► Identities

$$\int d^d k \frac{\partial}{\partial k^\mu} q^\mu I(a_1, a_2, \dots, a_6) = 0, \quad q^\mu = n^\mu, \bar{n}^\mu, v_3^\mu, v_4^\mu, k^\mu$$

Integration by parts and differential equations

- ▶ Identities

$$\int d^d k \frac{\partial}{\partial k^\mu} q^\mu I(a_1, a_2, \dots, a_6) = 0, \quad q^\mu = n^\mu, \bar{n}^\mu, v_3^\mu, v_4^\mu, k^\mu$$

- ▶ Reduction to a set of Master Integrals: $\{M_i\}$

Integration by parts and differential equations

- ▶ Identities

$$\int d^d k \frac{\partial}{\partial k^\mu} q^\mu I(a_1, a_2, \dots, a_6) = 0, \quad q^\mu = n^\mu, \bar{n}^\mu, v_3^\mu, v_4^\mu, k^\mu$$

- ▶ Reduction to a set of Master Integrals: $\{M_i\}$

- ▶ Differential equations

$$\frac{d}{d\beta_t} M_i = A_{ij} M_j$$

Integration by parts and differential equations

- ▶ Identities

$$\int d^d k \frac{\partial}{\partial k^\mu} q^\mu I(a_1, a_2, \dots, a_6) = 0, \quad q^\mu = n^\mu, \bar{n}^\mu, v_3^\mu, v_4^\mu, k^\mu$$

- ▶ Reduction to a set of Master Integrals: $\{M_i\}$
- ▶ Differential equations

$$\frac{d}{d\beta_t} M_i = A_{ij} M_j$$

- ▶ Straightforward integrations with $\beta_t = 0$ fix boundary conditions

Integration by parts and differential equations

- ▶ Identities

$$\int d^d k \frac{\partial}{\partial k^\mu} q^\mu I(a_1, a_2, \dots, a_6) = 0, \quad q^\mu = n^\mu, \bar{n}^\mu, v_3^\mu, v_4^\mu, k^\mu$$

- ▶ Reduction to a set of Master Integrals: $\{M_i\}$
- ▶ Differential equations

$$\frac{d}{d\beta_t} M_i = A_{ij} M_j$$

- ▶ Straightforward integrations with $\beta_t = 0$ fix boundary conditions
- ▶ Final set of bubble integrals: $\{I_{jk}(\beta_t, \theta)\}$

Complete Soft Function at NNLO: structure of the result

$$\mathcal{S}^{(2,\text{bare})}(L_{\perp}, \beta_t, \theta) = \left[\frac{1}{\epsilon} + L_{\perp} + L_{\perp}^2 + \dots \right] \\ \times \left[\mathcal{S}_{\text{bubble}}^{(2)}(\beta_t, \theta, \epsilon) + \mathcal{S}_{1\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) + \mathcal{S}_{2\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) \right]$$

Complete Soft Function at NNLO: structure of the result

$$\begin{aligned} S^{(2,\text{bare})}(L_{\perp}, \beta_t, \theta) &= \left[\frac{1}{\epsilon} + L_{\perp} + L_{\perp}^2 + \dots \right] \\ &\times \left[S_{\text{bubble}}^{(2)}(\beta_t, \theta, \epsilon) + S_{1\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) + S_{2\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) \right] \\ &= \frac{1}{\epsilon^2} S^{(2,-2)}(L_{\perp}) + \frac{1}{\epsilon} S^{(2,-1)}(L_{\perp}) + S^{(2,0)}(L_{\perp}) \end{aligned}$$

Complete Soft Function at NNLO: structure of the result

$$\begin{aligned} S^{(2,\text{bare})}(L_{\perp}, \beta_t, \theta) &= \left[\frac{1}{\epsilon} + L_{\perp} + L_{\perp}^2 + \dots \right] \\ &\times \left[S_{\text{bubble}}^{(2)}(\beta_t, \theta, \epsilon) + S_{1\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) + S_{2\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) \right] \\ &= \underbrace{\frac{1}{\epsilon^2} S^{(2,-2)}(L_{\perp}) + \frac{1}{\epsilon} S^{(2,-1)}(L_{\perp}) + S^{(2,0)}(L_{\perp})} \end{aligned}$$

can be cross-checked against RG; fixes all L_{\perp} -dependent terms in $S^{(2,0)}(L_{\perp})$

Complete Soft Function at NNLO: structure of the result

$$\begin{aligned} S^{(2,\text{bare})}(L_{\perp}, \beta_t, \theta) &= \left[\frac{1}{\epsilon} + L_{\perp} + L_{\perp}^2 + \dots \right] \\ &\times \left[S_{\text{bubble}}^{(2)}(\beta_t, \theta, \epsilon) + S_{1\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) + S_{2\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) \right] \\ &= \underbrace{\frac{1}{\epsilon^2} S^{(2,-2)}(L_{\perp}) + \frac{1}{\epsilon} S^{(2,-1)}(L_{\perp}) + S^{(2,0)}(L_{\perp})}_{\text{can be cross-checked against RG; fixes all } L_{\perp}\text{-dependent terms in } S^{(2,0)}(L_{\perp})} \end{aligned}$$

can be cross-checked against RG; fixes all L_{\perp} -dependent terms in $S^{(2,0)}(L_{\perp})$

- ▶ The only term that has to be obtained through direct calculation is the L_{\perp} -independent part of $S^{(2,0)}(L_{\perp})$.

Complete Soft Function at NNLO: structure of the result

$$\begin{aligned} S^{(2,\text{bare})}(L_{\perp}, \beta_t, \theta) &= \left[\frac{1}{\epsilon} + L_{\perp} + L_{\perp}^2 + \dots \right] \\ &\times \left[S_{\text{bubble}}^{(2)}(\beta_t, \theta, \epsilon) + S_{1\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) + S_{2\text{-cut}}^{(2)}(\beta_t, \theta, \epsilon) \right] \\ &= \underbrace{\frac{1}{\epsilon^2} S^{(2,-2)}(L_{\perp}) + \frac{1}{\epsilon} S^{(2,-1)}(L_{\perp}) + S^{(2,0)}(L_{\perp})} \end{aligned}$$

can be cross-checked against RG; fixes all L_{\perp} -dependent terms in $S^{(2,0)}(L_{\perp})$

- ▶ The only term that has to be obtained through direct calculation is the L_{\perp} -independent part of $S^{(2,0)}(L_{\perp})$.
- ▶ However, we calculate all terms and use the redundant ones for cross checks against Renormalization Group prediction.

Vanishing of higher order poles

Even though the NNLO Soft Function exhibits at most $\frac{1}{\epsilon^2}$ singularity, higher order poles appear in individual contributions.

Vanishing of higher order poles

Even though the NNLO Soft Function exhibits at most $\frac{1}{\epsilon^2}$ singularity, higher order poles appear in individual contributions.

- ▶ All α poles, including $\frac{\epsilon}{\alpha}$, as well as $\frac{1}{\epsilon^4}$ pole cancel within each colour structure, for example

$$\frac{1}{\epsilon^4} \begin{pmatrix} 0.00009 N_c^{-1} - 0.00009 N_c & -0.00002 N_c^2 - 0.00009 N_c^{-2} + 0.0001 \\ -0.00002 N_c^2 - 0.00009 N_c^{-2} + 0.0001 & 0.00008 N_c^3 - 0.00006 N_c + 0.00007 N_c^{-3} - 0.00009 N_c^{-1} \end{pmatrix}$$

Vanishing of higher order poles

Even though the NNLO Soft Function exhibits at most $\frac{1}{\epsilon^2}$ singularity, higher order poles appear in individual contributions.

- ▶ All α poles, including $\frac{\epsilon}{\alpha}$, as well as $\frac{1}{\epsilon^4}$ pole cancel within each colour structure, for example


$$\frac{1}{\epsilon^4} \begin{pmatrix} 0.00009 N_c^{-1} - 0.00009 N_c & -0.00002 N_c^2 - 0.00009 N_c^{-2} + 0.0001 \\ -0.00002 N_c^2 - 0.00009 N_c^{-2} + 0.0001 & 0.00008 N_c^3 - 0.00006 N_c + 0.00007 N_c^{-3} - 0.00009 N_c^{-1} \end{pmatrix}$$

- ▶ $\frac{1}{\epsilon^3}$ pole cancels between 1-cut and 2-cut contributions

$$\frac{1}{\epsilon^3} \begin{pmatrix} 0.0004 N_c^3 - 0.0007 N_c + 0.0004 N_c^{-1} & 0.0004 N_c^2 - 0.0004 N_c^{-2} - 7. \times 10^{-6} \\ 0.0004 N_c^2 - 0.0004 N_c^{-2} - 7. \times 10^{-6} & -0.0004 N_c^3 - 0.00001 N_c + 0.0003 N_c^{-3} + 0.0002 N_c^{-1} \end{pmatrix}$$

[†] We used $\beta = 0.4$, $\theta = 0.5$.

NNLO, small- q_T soft function for top pair production




Validation of the framework


- ▶ Perfect agreement of quark bubble results obtained from *differential equations* and *sector decomposition* for all terms in ϵ expansion
- ▶ Reproduction of the n_f part of Renormalization Group result




Imaginary part







($q\bar{q}$ channel)



(gg channel)







Conclusions

- ▶ Our goal: Use small- q_T factorization and q_T slicing to perform independent calculation of top pair production at NNLO

Conclusions

- ▶ Our goal: Use small- q_T factorization and q_T slicing to perform independent calculation of top pair production at NNLO
- ▶ The only missing component: the NNLO soft function

Conclusions

- ▶ Our goal: Use small- q_T factorization and q_T slicing to perform independent calculation of top pair production at NNLO
- ▶ The only missing component: the NNLO soft function
- ▶ We have constructed a framework based on sector decomposition and differential equations and used it to **complete the calculation of the the small- q_T soft function for top pair production at NNLO**

Conclusions

- ▶ Our goal: Use small- q_T factorization and q_T slicing to perform independent calculation of top pair production at NNLO
- ▶ The only missing component: the NNLO soft function
- ▶ We have constructed a framework based on sector decomposition and differential equations and used it to **complete the calculation of the the small- q_T soft function for top pair production at NNLO**
- ▶ The framework has been extensively validated and cross-checked:
 1. Cancellation of α poles, including ϵ/α , and ϵ poles beyond $1/\epsilon^2$
 2. Perfect agreement with analytic calculation for bubble graphs
 3. RG result for the complete NNLO soft function recovered: real and imaginary part

Conclusions

- ▶ Our goal: Use small- q_T factorization and q_T slicing to perform independent calculation of top pair production at NNLO
- ▶ The only missing component: the NNLO soft function
- ▶ We have constructed a framework based on sector decomposition and differential equations and used it to **complete the calculation of the the small- q_T soft function for top pair production at NNLO**
- ▶ The framework has been extensively validated and cross-checked:
 1. Cancellation of α poles, including ϵ/α , and ϵ poles beyond $1/\epsilon^2$
 2. Perfect agreement with analytic calculation for bubble graphs
 3. RG result for the complete NNLO soft function recovered: real and imaginary part → direct demonstration of validity of the small- q_T factorization for top pair production at NNLO

Conclusions

- ▶ Our goal: Use small- q_T factorization and q_T slicing to perform independent calculation of top pair production at NNLO
- ▶ The only missing component: the NNLO soft function
- ▶ We have constructed a framework based on sector decomposition and differential equations and used it to **complete the calculation of the the small- q_T soft function for top pair production at NNLO**
- ▶ The framework has been extensively validated and cross-checked:
 1. Cancellation of α poles, including ϵ/α , and ϵ poles beyond $1/\epsilon^2$
 2. Perfect agreement with analytic calculation for bubble graphs
 3. RG result for the complete NNLO soft function recovered: real and imaginary part \rightarrow direct demonstration of validity of the small- q_T factorization for top pair production at NNLO
- ▶ The soft function can now be used to obtain full $t\bar{t}$ cross section at NNLO as well for resummation up to NNLL'

Acknowledgements

This work has been supported by the National Science Centre, Poland grant POLONEZ No. 2015/19/P/ST2/03007. The project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement NO. 665778.

