

# Subleading power threshold resummation

Jian Wang

In coll. with M. Beneke, A. Broggio, M. Garny, S. Jaskiewicz, R. Szafron, L. Vernazza

arXiv:1809.10631



REF@Kraków

20/11/2018







# LP results

$$\hat{\sigma}(z) = H(Q^2) Q S_{\text{DY}}(Q(1-z))$$

P. A. Baikov, K. G. Chetyrkin, A. V. Smirnov, V. A. Smirnov and M. Steinhauser, R. N. Lee, '09

T. Gehrmann, E. W. N. Glover, T. Huber, N. Ikizlerli and C. Studerus, '10

Wilson lines

$$S_{\text{DY}}(\Omega) = \int \frac{dx^0}{4\pi} e^{ix^0\Omega/2} \frac{1}{N_c} \text{Tr} \langle 0 | \bar{\mathbf{T}}(Y_+^\dagger(x^0)Y_-(x^0)) \mathbf{T}(Y_-^\dagger(0)Y_+(0)) | 0 \rangle$$

C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog and B. Mistlberger, '13

Y. Li, A. von Manteuffel, R. M. Schabinger and H. X. Zhu, '13

# LP results

$$\hat{\sigma}(z) = H(Q^2) Q S_{\text{DY}}(Q(1-z))$$

P. A. Baikov, K. G. Chetyrkin, A. V. Smirnov, V. A. Smirnov and M. Steinhauser, R. N. Lee, '09

T. Gehrmann, E. W. N. Glover, T. Huber, N. Ikizlerli and C. Studerus, '10

Wilson lines

$$S_{\text{DY}}(\Omega) = \int \frac{dx^0}{4\pi} e^{ix^0\Omega/2} \frac{1}{N_c} \text{Tr} \langle 0 | \bar{\mathbf{T}}(Y_+^\dagger(x^0)Y_-(x^0)) \mathbf{T}(Y_-^\dagger(0)Y_+(0)) | 0 \rangle$$

C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog and B. Mistlberger, '13

Y. Li, A. von Manteuffel, R. M. Schabinger and H. X. Zhu, '13

In contrast, much less is understood at NLP.

# Structure of NLP logarithms

- the method of region approach, Bonocore et al 2014, Anastasiou et al 2014, Bahjat-Abbas et al 2018
- diagrammatic factorization techniques, Bonocore et al 2015, Bonocore et al 2016, Del Duca et al:2017

## Low-Burnett-Kroll theorem 1958,1968

$$-g_s \sum_{i=1}^N \mathbf{T}_i \left( \frac{p_i \cdot \epsilon(k)}{p_i \cdot k} + \frac{\epsilon_\mu(k) k_\nu J_i^{\mu\nu}}{p_i \cdot k} \right) A_0(\{p_i\})$$

$$J_i^{\mu\nu} = p_i^\mu \frac{\partial}{\partial p_{i\nu}} - p_i^\nu \frac{\partial}{\partial p_{i\mu}} + \Sigma_i^{\mu\nu}$$

# LP factorization

$$\hat{\sigma}(z) = H(Q^2) Q S_{\text{DY}}(Q(1-z))$$

$$(n_+ p, n_- p, p_\perp)$$

$$Q(1, 1, 1)$$

$$Q(1, \lambda^2, \lambda)$$


$$Q(\lambda^2, \lambda^2, \lambda^2)$$

$$\lambda = \sqrt{1-z}$$

$$\mu_h \sim Q$$

$$\mu_s \sim Q(1-z)$$

$$\mu_c \sim Q\sqrt{1-z}$$



# LP factorization

$$\hat{\sigma}(z) = H(Q^2) Q S_{\text{DY}}(Q(1-z))$$

$$(n_+ p, n_- p, p_\perp)$$

$$Q(1, 1, 1)$$

$$Q(1, \lambda^2, \lambda)$$


$$Q(\lambda^2, \lambda^2, \lambda^2)$$

$$\lambda = \sqrt{1-z}$$

$$\mu_h \sim Q$$


$$\mu_s \sim Q(1-z)$$

$$\mu_c \sim Q\sqrt{1-z}$$




Tricky point: no collinear function at LP

# NLP factorization




# NLP factorization



# NLP factorization

$$Q(1, \lambda^2, \lambda)$$

$$Q(\lambda^2, \lambda^2, \lambda^2)$$




# NLP factorization

$$\hat{\sigma}(z) = \sum_i \int d\omega_i d\bar{\omega}_i d\omega'_i d\bar{\omega}'_i D(-\hat{s}; \omega_i, \bar{\omega}_i) D^*(-\hat{s}; \omega'_i, \bar{\omega}'_i)$$

$$\times Q^2 \int \frac{d^3 \vec{q}}{(2\pi)^3 2\sqrt{Q^2 + \vec{q}^2}} \frac{1}{2\pi}$$


$$\int d^4 x e^{i(x_a p_A + x_b p_B - q) \cdot x} \tilde{S}(x; \omega_i, \bar{\omega}_i, \omega'_i, \bar{\omega}'_i)$$



$$D(-\hat{s}; \omega_i, \bar{\omega}_i) = \int d(n_+ p_i) d(n_- \bar{p}_i) C(n_+ p_i, n_- \bar{p}_i) \quad n_+ p_i, n_- \bar{p}_i \sim O(1)$$

$$\times J(n_+ p_i, x_a n_+ p_A; \omega_i) \bar{J}(n_- \bar{p}_i, -x_b n_- p_B; \bar{\omega}_i) \quad \omega_i, \bar{\omega}_i \sim O(\lambda^2)$$

# Hard function




$$\bar{\psi}\gamma_{\mu}\psi(0) = \int dt d\bar{t} \tilde{C}^{A0}(t, \bar{t}) J_{\mu}^{A0}(t, \bar{t})$$

$$C^{A0}(n+p, n-\bar{p}) = \int dt d\bar{t} e^{-itn+p - i\bar{t}n-\bar{p}} \tilde{C}^{A0}(t, \bar{t})$$

$$J_{\mu}^{A0}(t, \bar{t}) = \bar{\chi}_{\bar{c}}(\bar{t}n_{-})\gamma_{\perp\mu}\chi_c(tn_{+})$$

# Hard function



$$\bar{\psi} \gamma_{\mu} \psi(0) = \int dt d\bar{t} \tilde{C}^{A0}(t, \bar{t}) J_{\mu}^{A0}(t, \bar{t})$$

$$C^{A0}(n_+ p, n_- \bar{p}) = \int dt d\bar{t} e^{-itn_+ p - i\bar{t}n_- \bar{p}} \tilde{C}^{A0}(t, \bar{t})$$


$$J_{\mu}^{A0}(t, \bar{t}) = \bar{\chi}_{\bar{c}}(\bar{t}n_-) \gamma_{\perp \mu} \chi_c(tn_+)$$



$$\hat{\sigma}(z) = H(Q^2) Q S_{\text{DY}}(Q(1-z))$$

# NLP jet function

$\chi_c^{\text{PDF}}$



$$i \int d^4 z e^{i\omega(n_+ z)/2} \mathbf{T} \left[ \chi_{c,\alpha a}(tn_+) \bar{\chi}_{c,d}(z) \frac{\not{n}_+}{2} \chi_{c,e}(z) \right]$$


$$= 2\pi \int du \tilde{J}_{\alpha\beta,abde}(t, u; \omega) \chi_{c,\beta b}^{\text{PDF}}(un_+)$$

NLP quark-gluon interaction: Beneke et al 2002

$$\mathcal{L}_{2\xi}^{(2)} = \frac{1}{2} \bar{\chi}_c x_{\perp}^{\mu} x_{\perp}^{\nu} [i\partial_{\nu} i n_{-} \partial \mathcal{B}_{\mu}^{+}] \frac{\not{n}_+}{2} \chi_c \quad \mathcal{B}_{\pm}^{\mu} = Y_{\pm}^{\dagger} [iD_s^{\mu} Y_{\pm}]$$

# NLP jet function

$\chi_c^{\text{PDF}}$



$$i \int d^4 z e^{i\omega(n_+ z)/2} \mathbf{T} \left[ \chi_{c,\alpha a}(tn_+) \bar{\chi}_{c,d}(z) \frac{\not{n}_+}{2} \chi_{c,e}(z) \right]$$

$$\chi_c = 2\pi \int du \tilde{J}_{\alpha\beta,abde}(t, u; \omega) \chi_{c,\beta b}^{\text{PDF}}(un_+)$$

Field definition of radiative jet function


del Duca 1990,  
Bonocore et al '15,'16

NLP quark-gluon interaction: Beneke et al 2002

$$\mathcal{L}_{2\xi}^{(2)} = \frac{1}{2} \bar{\chi}_c x_{\perp}^{\mu} x_{\perp}^{\nu} [i\partial_{\nu} i n_{-} \partial \mathcal{B}_{\mu}^{+}] \frac{\not{n}_+}{2} \chi_c \quad \mathcal{B}_{\pm}^{\mu} = Y_{\pm}^{\dagger} [iD_s^{\mu} Y_{\pm}]$$

# NLP jet function

$\chi_c^{\text{PDF}}$



$$i \int d^4 z e^{i\omega(n_+ z)/2} \mathbf{T} \left[ \chi_{c,\alpha a}(tn_+) \bar{\chi}_{c,d}(z) \frac{\not{n}_+}{2} \chi_{c,e}(z) \right]$$

$$\chi_c = 2\pi \int du \tilde{J}_{\alpha\beta,abde}(t, u; \omega) \chi_{c,\beta b}^{\text{PDF}}(un_+)$$

Field definition of radiative jet function


del Duca 1990,  
Bonocore et al '15,'16

NLP quark-gluon interaction: Beneke et al 2002

$$\mathcal{L}_{2\xi}^{(2)} = \frac{1}{2} \bar{\chi}_c x_{\perp}^{\mu} x_{\perp}^{\nu} [i\partial_{\nu} i n_{-} \partial \mathcal{B}_{\mu}^{+}] \frac{\not{n}_+}{2} \chi_c \quad \mathcal{B}_{\pm}^{\mu} = Y_{\pm}^{\dagger} [iD_s^{\mu} Y_{\pm}]$$

LO:  $J_{2\xi;\alpha\beta,abde}^{\mu\rho}(n_+p, n_+p'; \omega) = -\frac{g_{\perp}^{\mu\rho}}{n_+p} \delta(n_+p - n_+p') \delta_{\alpha\beta} \delta_{ad} \delta_{eb}$


# NLP factorization



$$\bar{\psi}\gamma^\mu\psi(0) = \int dt d\bar{t} \tilde{C}^{A0}(t, \bar{t}) \left[ \boxed{O(1)} J_{A0}^\mu(t, \bar{t}) + \boxed{O(\lambda^2)} (J_{A0,2\xi}^{T2}(t, \bar{t}))^\mu + \bar{c}\text{-term} \right]$$

$$(J_{A0,2\xi}^{T2}(s, t))^\mu = i \int d^4x \mathbf{T} \left[ J_{A0}^\mu(s, t) \mathcal{L}_{2\xi}^{(2)}(x) \right]$$

# NLP factorization



$$\bar{\psi}\gamma^\mu\psi(0) = \int dt d\bar{t} \tilde{C}^{A0}(t, \bar{t}) \left[ \boxed{O(1)} J_{A0}^\mu(t, \bar{t}) + \boxed{O(\lambda^2)} (J_{A0,2\xi}^{T2}(t, \bar{t}))^\mu + \bar{c}\text{-term} \right]$$

$$(J_{A0,2\xi}^{T2}(s, t))^\mu = i \int d^4x \mathbf{T} \left[ J_{A0}^\mu(s, t) \mathcal{L}_{2\xi}^{(2)}(x) \right]$$


qg-channel:

$$\bar{\psi}\gamma^\mu\psi(0) = \int dt d\bar{t} \tilde{C}^{A0}(t, \bar{t}) \left[ (J_{A0,\xi q}^{T1}(t, \bar{t}))^\mu + \bar{c}\text{-term} \right]$$

$$(J_{A0,\xi q}^{T1}(s, t))^\mu = i \int d^4x \mathbf{T} \left[ J_{A0}^\mu(s, t) \mathcal{L}_{\xi q}^{(1)}(x) \right]$$

$$\mathcal{L}_{\xi q}^{(1)} = \bar{q}_+ \mathcal{A}_{c\perp} \chi_c + \text{h.c.}$$

# Soft function at NLP



LO:


$$\mathcal{B}_{\pm}^{\mu} = Y_{\pm}^{\dagger} [iD_s^{\mu} Y_{\pm}]$$

$$\tilde{\mathcal{S}}_{2\xi}(x, z_{-}) = \bar{\mathbf{T}} [Y_{+}^{\dagger}(x) Y_{-}(x)] \mathbf{T} [Y_{-}^{\dagger}(0) Y_{+}(0) \frac{i\partial_{\perp}^{\nu}}{in_{-}\partial} \mathcal{B}_{\perp\nu}^{+}(z_{-})]$$

$$S_{2\xi}(\Omega, \omega) = \int \frac{dx^0}{4\pi} \int \frac{d(n+z)}{4\pi} e^{ix^0\Omega/2 - i\omega(n+z)/2} \frac{1}{N_c} \text{Tr} \langle 0 | \tilde{\mathcal{S}}_{2\xi}(x^0, z_{-}) | 0 \rangle$$

$$S_{2\xi}(\Omega, \omega) = \frac{\alpha_s C_F}{2\pi} \left\{ \theta(\Omega) \delta(\omega) \left( -\frac{1}{\epsilon} + \ln \frac{\Omega^2}{\mu^2} \right) + \left[ \frac{1}{\omega} \right]_{+} \theta(\omega) \theta(\Omega - \omega) \right\}$$

# Soft function at NLP



LO:

$$\mathcal{B}_{\pm}^{\mu} = Y_{\pm}^{\dagger} [iD_s^{\mu} Y_{\pm}]$$

$$\tilde{\mathcal{S}}_{2\xi}(x, z_{-}) = \bar{\mathbf{T}} [Y_{+}^{\dagger}(x) Y_{-}(x)] \mathbf{T} [Y_{-}^{\dagger}(0) Y_{+}(0) \frac{i\partial_{\perp}^{\nu}}{in_{-}\partial} \mathcal{B}_{\perp\nu}^{+}(z_{-})]$$

$$S_{2\xi}(\Omega, \omega) = \int \frac{dx^0}{4\pi} \int \frac{d(n+z)}{4\pi} e^{ix^0\Omega/2 - i\omega(n+z)/2} \frac{1}{N_c} \text{Tr} \langle 0 | \tilde{\mathcal{S}}_{2\xi}(x^0, z_{-}) | 0 \rangle$$

$$S_{2\xi}(\Omega, \omega) = \frac{\alpha_s C_F}{2\pi} \left\{ \theta(\Omega) \delta(\omega) \left( -\frac{1}{\epsilon} + \ln \frac{\Omega^2}{\mu^2} \right) + \left[ \frac{1}{\omega} \right]_{+} \theta(\omega) \theta(\Omega - \omega) \right\}$$

A puzzle: divergence at LO

# RG condition

$$S_{2\xi}(\Omega, \omega) = \frac{\alpha_s C_F}{2\pi} \left\{ \theta(\Omega)\delta(\omega) \left( -\frac{1}{\epsilon} + \ln \frac{\Omega^2}{\mu^2} \right) + \left[ \frac{1}{\omega} \right]_+ \theta(\omega)\theta(\Omega - \omega) \right\}$$

$$S_{2\xi}(\Omega, \omega)|_{\text{ren}} = \int d\Omega' \int d\omega' Z_{2\xi, 2\xi}(\Omega, \omega; \Omega', \omega') S_{2\xi}(\Omega', \omega')|_{\text{bare}} \\ + \int d\Omega' Z_{2\xi, x_0}(\Omega, \omega; \Omega') S_{x_0}(\Omega')|_{\text{bare}}$$

$$Z_{2\xi, 2\xi}(\Omega, \omega; \Omega, \omega') = \delta(\Omega - \Omega')\delta(\omega - \omega') + \mathcal{O}(\alpha_s),$$

$$Z_{2\xi, x_0}(\Omega, \omega; \Omega') = \frac{\alpha_s C_F}{2\pi} \frac{1}{\epsilon} \delta(\Omega - \Omega')\delta(\omega) + \mathcal{O}(\alpha_s^2)$$

The mixing term subtracts the divergent part of the first term on the right-hand side, resulting in a finite, renormalized soft function

# auxiliary soft function

$$S_{x_0}(\Omega) = \theta(\Omega)$$

We propose

$$S_{x_0}(\Omega) = \int \frac{dx^0}{4\pi} e^{ix^0\Omega/2} \frac{-2i}{x^0 - i\varepsilon} \frac{1}{N_c} \text{Tr} \langle 0 | \bar{\mathbf{T}} \left[ Y_+^\dagger(x^0) Y_-(x^0) \right] \mathbf{T} \left[ Y_-^\dagger(0) Y_+(0) \right] | 0 \rangle$$

“Theta-soft function” in NLP thrust distribution,

Moult, Stewart, Vita, Zhu '18

# auxiliary soft function

$$S_{x_0}(\Omega) = \theta(\Omega)$$

We propose

$$S_{x_0}(\Omega) = \int \frac{dx^0}{4\pi} e^{ix^0\Omega/2} \frac{-2i}{x^0 - i\varepsilon} \frac{1}{N_c} \text{Tr} \langle 0 | \bar{\mathbf{T}} \left[ Y_+^\dagger(x^0) Y_-(x^0) \right] \mathbf{T} \left[ Y_-^\dagger(0) Y_+(0) \right] | 0 \rangle$$

“Theta-soft function” in NLP thrust distribution,

Moult, Stewart, Vita, Zhu '18

We check this form by requiring the poles cancel at two loop

# Check

$$S_{2\xi}^{(2)} + Z_{2\xi x_0}^{(1)} S_{x_0}^{(1)} + Z_{2\xi x_0}^{(2)} S_{x_0}^{(0)} + Z_{2\xi 2\xi}^{(1)} S_{2\xi}^{(1)} = \text{finite}$$

Under assumption that the off-diag has only  
subleading pole

$$S_{2\xi}^{(2)} - \frac{1}{4} Z_{2\xi x_0}^{(1)} \left( 3Z_{2\xi 2\xi}^{(1)} + Z_{x_0 x_0}^{(1)} \right) S_{x_0}^{(0)} = \mathcal{O} \left( \frac{1}{\epsilon^2} \right)$$

# Check

$$S_{2\xi}^{(2)} + Z_{2\xi x_0}^{(1)} S_{x_0}^{(1)} + Z_{2\xi x_0}^{(2)} S_{x_0}^{(0)} + Z_{2\xi 2\xi}^{(1)} S_{2\xi}^{(1)} = \text{finite}$$

Under assumption that the off-diag has only  
subleading pole

$$S_{2\xi}^{(2)} - \frac{1}{4} Z_{2\xi x_0}^{(1)} \left( 3Z_{2\xi 2\xi}^{(1)} + Z_{x_0 x_0}^{(1)} \right) S_{x_0}^{(0)} = \mathcal{O} \left( \frac{1}{\epsilon^2} \right)$$



Same as LP soft fun.



# Check


$$S_{2\xi}^{(2)} + Z_{2\xi x_0}^{(1)} S_{x_0}^{(1)} + Z_{2\xi x_0}^{(2)} S_{x_0}^{(0)} + Z_{2\xi 2\xi}^{(1)} S_{2\xi}^{(1)} = \text{finite}$$

Under assumption that the off-diag has only subleading pole


$$S_{2\xi}^{(2)} - \frac{1}{4} Z_{2\xi x_0}^{(1)} \left( 3Z_{2\xi 2\xi}^{(1)} + Z_{x_0 x_0}^{(1)} \right) S_{x_0}^{(0)} = \mathcal{O}\left(\frac{1}{\epsilon^2}\right)$$

Known from 1-loop Same as LP soft fun.


$$S_{2\xi}^{(2)}$$



$$Z_{2\xi}^{(1)} \quad 2\xi \quad 2\xi$$



$$\bar{\mathbf{T}} \left[ Y_+^\dagger(x) Y_-(x) \right] \mathbf{T} \left[ Y_-^\dagger(0) Y_+(0) \frac{i\partial_\perp^\nu}{in_- \partial} \mathcal{B}_{\perp\nu}^+(z_-) \right]$$



# RG eq. of soft fun.

$$\frac{d}{d \ln \mu} \begin{pmatrix} S_{2\xi}(\Omega, \omega) \\ S_{x_0}(\Omega) \end{pmatrix} = \frac{\alpha_s}{\pi} \begin{pmatrix} 4C_F \ln \frac{\mu}{\mu_s} & -C_F \delta(\omega) \\ 0 & 4C_F \ln \frac{\mu}{\mu_s} \end{pmatrix} \begin{pmatrix} S_{2\xi}(\Omega, \omega) \\ S_{x_0}(\Omega) \end{pmatrix}$$

# RG eq. of soft fun.

$$\frac{d}{d \ln \mu} \begin{pmatrix} S_{2\xi}(\Omega, \omega) \\ S_{x_0}(\Omega) \end{pmatrix} = \frac{\alpha_s}{\pi} \begin{pmatrix} 4C_F \ln \frac{\mu}{\mu_s} & -C_F \delta(\omega) \\ 0 & 4C_F \ln \frac{\mu}{\mu_s} \end{pmatrix} \begin{pmatrix} S_{2\xi}(\Omega, \omega) \\ S_{x_0}(\Omega) \end{pmatrix}$$



$$S_{2\xi}^{\text{LL}}(\Omega, \omega, \mu) = \frac{2C_F}{\beta_0} \ln \frac{\alpha_s(\mu)}{\alpha_s(\mu_s)} \exp \left[ -4S^{\text{LL}}(\mu_s, \mu) \right] \theta(\Omega) \delta(\omega)$$

---

 $\alpha_s \ln \frac{\mu}{\mu_s}$

---

 $\alpha_s \ln^2 \frac{\mu}{\mu_s}$

# RG eq. of soft fun.

$$\frac{d}{d \ln \mu} \begin{pmatrix} S_{2\xi}(\Omega, \omega) \\ S_{x_0}(\Omega) \end{pmatrix} = \frac{\alpha_s}{\pi} \begin{pmatrix} 4C_F \ln \frac{\mu}{\mu_s} & -C_F \delta(\omega) \\ 0 & 4C_F \ln \frac{\mu}{\mu_s} \end{pmatrix} \begin{pmatrix} S_{2\xi}(\Omega, \omega) \\ S_{x_0}(\Omega) \end{pmatrix}$$



$$S_{2\xi}^{\text{LL}}(\Omega, \omega, \mu) = \frac{2C_F}{\beta_0} \ln \frac{\alpha_s(\mu)}{\alpha_s(\mu_s)} \exp \left[ -4S^{\text{LL}}(\mu_s, \mu) \right] \theta(\Omega) \delta(\omega)$$

$$\alpha_s \ln \frac{\mu}{\mu_s} \qquad \alpha_s \ln^2 \frac{\mu}{\mu_s}$$

Similarly, for the hard function

$$H(Q^2, \mu) = \exp \left[ 4S(\mu_h, \mu) \right]$$

$$\alpha_s \ln^2 \frac{\mu}{\mu_h}$$

# Kinematic corrections

In the partonic c.o.m frame, the energy of the soft hadronic final state is expanded as

$$[x_1 p_1 + x_2 p_2 - q]^0 = p_{X_s}^0 = \sqrt{\hat{s}} - \sqrt{Q^2 + \vec{q}^2} = \frac{\Omega_*}{2} - \frac{\vec{q}^2}{2Q} + O(\lambda^6)$$

$$\Omega_* = 2Q \frac{1 - \sqrt{z}}{\sqrt{z}} = Q(1 - z) + \frac{3}{4}Q(1 - z)^2 + O(\lambda^6)$$

# Kinematic corrections

In the partonic c.o.m frame, the energy of the soft hadronic final state is expanded as

$$[x_1 p_1 + x_2 p_2 - q]^0 = p_{X_s}^0 = \sqrt{\hat{s}} - \sqrt{Q^2 + \vec{q}^2} = \frac{\Omega_*}{2} - \frac{\vec{q}^2}{2Q} + O(\lambda^6)$$

$$\Omega_* = 2Q \frac{1 - \sqrt{z}}{\sqrt{z}} = Q(1 - z) + \frac{3}{4}Q(1 - z)^2 + O(\lambda^6)$$

The soft function expands

$$S_{\text{DY}}(Q(1 - z)) + \frac{1}{Q} S_{K1}(Q(1 - z)) + \frac{1}{Q} S_{K2}(Q(1 - z)) + O(\lambda^4)$$

$$S_{K1}(\Omega) = \frac{\partial}{\partial \Omega} \partial_{\vec{x}}^2 S_0(\Omega, \vec{x})|_{\vec{x}=0} ,$$

$$S_{K2}(\Omega) = \frac{3}{4} \Omega^2 \frac{\partial}{\partial \Omega} S_0(\Omega, \vec{x})|_{\vec{x}=0}$$

# Kinematic corrections

In the partonic c.o.m frame, the energy of the soft hadronic final state is expanded as

$$[x_1 p_1 + x_2 p_2 - q]^0 = p_{X_s}^0 = \sqrt{\hat{s}} - \sqrt{Q^2 + \vec{q}^2} = \frac{\Omega_*}{2} - \frac{\vec{q}^2}{2Q} + O(\lambda^6)$$

$$\Omega_* = 2Q \frac{1 - \sqrt{z}}{\sqrt{z}} = Q(1 - z) + \frac{3}{4}Q(1 - z)^2 + O(\lambda^6)$$

The soft function expands

$$S_{\text{DY}}(Q(1 - z)) + \frac{1}{Q} S_{K1}(Q(1 - z)) + \frac{1}{Q} S_{K2}(Q(1 - z)) + O(\lambda^4)$$

$$\Delta_{ab}(z) = \frac{\hat{\sigma}_{ab}(z)}{z}$$

$$S_{K1}(\Omega) = \frac{\partial}{\partial \Omega} \partial_{\vec{x}}^2 S_0(\Omega, \vec{x})|_{\vec{x}=0},$$

$$S_{K2}(\Omega) = \frac{3}{4} \Omega^2 \frac{\partial}{\partial \Omega} S_0(\Omega, \vec{x})|_{\vec{x}=0}$$

$$\rightarrow S_{K3}(\Omega) = \Omega S_0(\Omega, \vec{x})|_{\vec{x}=0}$$

# Kinematic corrections

In the partonic c.o.m frame, the energy of the soft hadronic final state is expanded as

$$[x_1 p_1 + x_2 p_2 - q]^0 = p_{X_s}^0 = \sqrt{\hat{s}} - \sqrt{Q^2 + \vec{q}^2} = \frac{\Omega_*}{2} - \frac{\vec{q}^2}{2Q} + O(\lambda^6)$$

$$\Omega_* = 2Q \frac{1 - \sqrt{z}}{\sqrt{z}} = Q(1 - z) + \frac{3}{4}Q(1 - z)^2 + O(\lambda^6)$$

The soft function expands

$$S_{\text{DY}}(Q(1 - z)) + \frac{1}{Q} S_{K1}(Q(1 - z)) + \frac{1}{Q} S_{K2}(Q(1 - z)) + O(\lambda^4)$$

$$\Delta_{ab}(z) = \frac{\hat{\sigma}_{ab}(z)}{z}$$

$$S_{K1}(\Omega) = \frac{\partial}{\partial \Omega} \partial_{\vec{x}}^2 S_0(\Omega, \vec{x})|_{\vec{x}=0},$$

$$S_{K2}(\Omega) = \frac{3}{4} \Omega^2 \frac{\partial}{\partial \Omega} S_0(\Omega, \vec{x})|_{\vec{x}=0}$$

$$\rightarrow S_{K3}(\Omega) = \Omega S_0(\Omega, \vec{x})|_{\vec{x}=0}$$

$$\sum_{i=1}^3 S_{Ki}(\Omega) = 2 \frac{\alpha_s C_F}{\pi}$$

# Kinematic corrections

In the partonic c.o.m frame, the energy of the soft hadronic final state is expanded as

$$[x_1 p_1 + x_2 p_2 - q]^0 = p_{X_s}^0 = \sqrt{\hat{s}} - \sqrt{Q^2 + \vec{q}^2} = \frac{\Omega_*}{2} - \frac{\vec{q}^2}{2Q} + O(\lambda^6)$$

$$\Omega_* = 2Q \frac{1 - \sqrt{z}}{\sqrt{z}} = Q(1 - z) + \frac{3}{4}Q(1 - z)^2 + O(\lambda^6)$$

The soft function expands

$$S_{\text{DY}}(Q(1 - z)) + \frac{1}{Q} S_{K1}(Q(1 - z)) + \frac{1}{Q} S_{K2}(Q(1 - z)) + O(\lambda^4)$$

$$\Delta_{ab}(z) = \frac{\hat{\sigma}_{ab}(z)}{z}$$

$$S_{K1}(\Omega) = \frac{\partial}{\partial \Omega} \partial_{\vec{x}}^2 S_0(\Omega, \vec{x})|_{\vec{x}=0},$$

$$S_{K2}(\Omega) = \frac{3}{4} \Omega^2 \frac{\partial}{\partial \Omega} S_0(\Omega, \vec{x})|_{\vec{x}=0}$$

$$\rightarrow S_{K3}(\Omega) = \Omega S_0(\Omega, \vec{x})|_{\vec{x}=0}$$

$$\sum_{i=1}^3 S_{Ki}(\Omega) = 2 \frac{\alpha_s C_F}{\pi}$$

No kine.cor.

# Final results

$$\mu_h \sim Q$$

$$\mu_s \sim Q(1-z)$$

$$\mu_c \sim Q\sqrt{1-z}$$

$$\Delta_{\text{NLP}}^{\text{LL}}(z) = -\exp [4S^{\text{LL}}(\mu_h, \mu_c) - 4S^{\text{LL}}(\mu_s, \mu_c)] \times \frac{8C_F}{\beta_0} \ln \frac{\alpha_s(\mu_c)}{\alpha_s(\mu_s)} \theta(1-z)$$

Why we evolve the hard/soft function to the jet scale?

# Final results

$$\mu_h \sim Q$$

$$\mu_s \sim Q(1-z)$$

$$\mu_c \sim Q\sqrt{1-z}$$

$$\Delta_{\text{NLP}}^{\text{LL}}(z) = -\exp [4S^{\text{LL}}(\mu_h, \mu_c) - 4S^{\text{LL}}(\mu_s, \mu_c)] \times \frac{8C_F}{\beta_0} \ln \frac{\alpha_s(\mu_c)}{\alpha_s(\mu_s)} \theta(1-z)$$

Why we evolve the hard/soft function to the jet scale?

We use the LO jet function. Recover the general scale dependence by the AP splitting kernels

# Final results

$$\mu_h \sim Q$$

$$\mu_s \sim Q(1-z)$$

$$\mu_c \sim Q\sqrt{1-z}$$

$$\Delta_{\text{NLP}}^{\text{LL}}(z) = -\exp \left[ 4S^{\text{LL}}(\mu_h, \mu_c) - 4S^{\text{LL}}(\mu_s, \mu_c) \right] \times \frac{8C_F}{\beta_0} \ln \frac{\alpha_s(\mu_c)}{\alpha_s(\mu_s)} \theta(1-z)$$

Why we evolve the hard/soft function to the jet scale?

We use the LO jet function. Recover the general scale dependence by the AP splitting kernels

$$\frac{d}{d \ln \mu} \hat{\sigma}_{ab}(z, \mu) = - \sum_c \int_z^1 dx \left( P_{ca}(x) \hat{\sigma}_{cb} \left( \frac{z}{x}, \mu \right) + P_{cb}(x) \hat{\sigma}_{ac} \left( \frac{z}{x}, \mu \right) \right)$$

$$P_{ab}^{\text{LP}}(x) = \left( 2\Gamma_{\text{cusp}}(\alpha_s) \frac{1}{[1-x]_+} + 2\gamma^\phi(\alpha_s) \delta(1-x) \right) \delta_{ab}$$

$$P_{ab}^{\text{NLP}} = \gamma_{ab}^{\text{NLP}}(\alpha_s)$$

# Final results

$$\begin{aligned} & \frac{d}{d \ln \mu} \Delta_{\text{NLP}}(z, \mu) \\ &= -4 \left[ \Gamma_{\text{cusp}}(\alpha_s) \left( \ln(1-z) - \gamma_E - \psi \left( 1 + \frac{d}{d \ln(1-z)} \right) \right) + \gamma^\phi(\alpha_s) \right] \Delta_{\text{NLP}}(z, \mu) \\ &+ K(z, \mu) \end{aligned}$$

$$K(z, \mu) = -2\gamma_{qq}^{\text{NLP}}(\alpha_s) \int_z^1 dy \Delta_{\text{LP}}(y, \mu) - 4\Gamma_{\text{cusp}}(\alpha_s)(1-z)\Delta_{\text{LP}}(z, \mu)$$

# Final results

$$\begin{aligned} & \frac{d}{d \ln \mu} \Delta_{\text{NLP}}(z, \mu) \\ &= -4 \left[ \Gamma_{\text{cusp}}(\alpha_s) \left( \ln(1-z) - \gamma_E - \psi \left( 1 + \frac{d}{d \ln(1-z)} \right) \right) + \gamma^\phi(\alpha_s) \right] \Delta_{\text{NLP}}(z, \mu) \\ &+ K(z, \mu) \end{aligned}$$

$$K(z, \mu) = -2\gamma_{qq}^{\text{NLP}}(\alpha_s) \int_z^1 dy \Delta_{\text{LP}}(y, \mu) - 4\Gamma_{\text{cusp}}(\alpha_s)(1-z)\Delta_{\text{LP}}(z, \mu)$$

---


LL

# Final results

$$\begin{aligned} & \frac{d}{d \ln \mu} \Delta_{\text{NLP}}(z, \mu) \\ &= -4 \left[ \Gamma_{\text{cusp}}(\alpha_s) \left( \ln(1-z) - \gamma_E - \psi \left( 1 + \frac{d}{d \ln(1-z)} \right) \right) + \gamma^\phi(\alpha_s) \right] \Delta_{\text{NLP}}(z, \mu) \\ &+ K(z, \mu) \end{aligned}$$

$$K(z, \mu) = -2\gamma_{qq}^{\text{NLP}}(\alpha_s) \int_z^1 dy \Delta_{\text{LP}}(y, \mu) - 4\Gamma_{\text{cusp}}(\alpha_s)(1-z)\Delta_{\text{LP}}(z, \mu)$$

LL



$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = \exp [4S^{\text{LL}}(\mu_h, \mu) - 4S^{\text{LL}}(\mu_s, \mu)] \times \frac{-8C_F}{\beta_0} \ln \frac{\alpha_s(\mu)}{\alpha_s(\mu_s)} \theta(1-z)$$

# Expansion

$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = \exp \left[ -2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_h} \right] \exp \left[ +2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_s} \right] \\ \times (-4) \frac{\alpha_s C_F}{\pi} \ln \frac{\mu_s}{\mu} \theta(1-z)$$



$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = -\theta(1-z) \left\{ 4C_F \frac{\alpha_s}{\pi} \left[ \ln(1-z) - L_\mu \right] \right. \\ + 8C_F^2 \left( \frac{\alpha_s}{\pi} \right)^2 \left[ \ln^3(1-z) - 3L_\mu \ln^2(1-z) + 2L_\mu^2 \ln(1-z) \right] \\ + 8C_F^3 \left( \frac{\alpha_s}{\pi} \right)^3 \left[ \ln^5(1-z) - 5L_\mu \ln^4(1-z) + 8L_\mu^2 \ln^3(1-z) - 4L_\mu^3 \ln^2(1-z) \right] \\ + \frac{16}{3} C_F^4 \left( \frac{\alpha_s}{\pi} \right)^4 \left[ \ln^7(1-z) - 7L_\mu \ln^6(1-z) + 18L_\mu^2 \ln^5(1-z) - 20L_\mu^3 \ln^4(1-z) \right. \\ \left. + 8L_\mu^4 \ln^3(1-z) \right] \\ + \frac{8}{3} C_F^5 \left( \frac{\alpha_s}{\pi} \right)^5 \left[ \ln^9(1-z) - 9L_\mu \ln^8(1-z) + 32L_\mu^2 \ln^7(1-z) - 56L_\mu^3 \ln^6(1-z) \right. \\ \left. + 48L_\mu^4 \ln^5(1-z) - 16L_\mu^5 \ln^4(1-z) \right] \left. \right\} + \mathcal{O}(\alpha_s^6 \times (\log)^{11})$$

$$L_\mu = \ln \mu/Q$$

# Expansion

$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = \exp \left[ -2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_h} \right] \exp \left[ +2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_s} \right] \\ \times (-4) \frac{\alpha_s C_F}{\pi} \ln \frac{\mu_s}{\mu} \theta(1-z)$$



$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = -\theta(1-z) \left\{ 4C_F \frac{\alpha_s}{\pi} \left[ \ln(1-z) - L_\mu \right] \right. \\ \left. + 8C_F^2 \left( \frac{\alpha_s}{\pi} \right)^2 \left[ \ln^3(1-z) - 3L_\mu \ln^2(1-z) + 2L_\mu^2 \ln(1-z) \right] \right. \\ \left. + 8C_F^3 \left( \frac{\alpha_s}{\pi} \right)^3 \left[ \ln^5(1-z) - 5L_\mu \ln^4(1-z) + 8L_\mu^2 \ln^3(1-z) - 4L_\mu^3 \ln^2(1-z) \right] \right. \\ \left. + \frac{16}{3} C_F^4 \left( \frac{\alpha_s}{\pi} \right)^4 \left[ \ln^7(1-z) - 7L_\mu \ln^6(1-z) + 18L_\mu^2 \ln^5(1-z) - 20L_\mu^3 \ln^4(1-z) \right. \right. \\ \left. \left. + 8L_\mu^4 \ln^3(1-z) \right] \right. \\ \left. + \frac{8}{3} C_F^5 \left( \frac{\alpha_s}{\pi} \right)^5 \left[ \ln^9(1-z) - 9L_\mu \ln^8(1-z) + 32L_\mu^2 \ln^7(1-z) - 56L_\mu^3 \ln^6(1-z) \right. \right. \\ \left. \left. + 48L_\mu^4 \ln^5(1-z) - 16L_\mu^5 \ln^4(1-z) \right] \right\} + \mathcal{O}(\alpha_s^6 \times (\log)^{11})$$

Hamberg, van Neerven, Matsuura, 1991

$$L_\mu = \ln \mu/Q$$

# Expansión

$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = \exp \left[ -2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_h} \right] \exp \left[ +2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_s} \right] \\ \times (-4) \frac{\alpha_s C_F}{\pi} \ln \frac{\mu_s}{\mu} \theta(1-z)$$



$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = -\theta(1-z) \left\{ 4C_F \frac{\alpha_s}{\pi} \left[ \ln(1-z) - L_\mu \right] \right. \\ \left. + 8C_F^2 \left( \frac{\alpha_s}{\pi} \right)^2 \left[ \ln^3(1-z) - 3L_\mu \ln^2(1-z) + 2L_\mu^2 \ln(1-z) \right] \right. \\ \left. + 8C_F^3 \left( \frac{\alpha_s}{\pi} \right)^3 \left[ \ln^5(1-z) - 5L_\mu \ln^4(1-z) + 8L_\mu^2 \ln^3(1-z) - 4L_\mu^3 \ln^2(1-z) \right] \right. \\ \left. + \frac{16}{3} C_F^4 \left( \frac{\alpha_s}{\pi} \right)^4 \left[ \ln^7(1-z) - 7L_\mu \ln^6(1-z) + 18L_\mu^2 \ln^5(1-z) - 20L_\mu^3 \ln^4(1-z) \right. \right. \\ \left. \left. + 8L_\mu^4 \ln^3(1-z) \right] \right. \\ \left. + \frac{8}{3} C_F^5 \left( \frac{\alpha_s}{\pi} \right)^5 \left[ \ln^9(1-z) - 9L_\mu \ln^8(1-z) + 32L_\mu^2 \ln^7(1-z) - 56L_\mu^3 \ln^6(1-z) \right. \right. \\ \left. \left. + 48L_\mu^4 \ln^5(1-z) - 16L_\mu^5 \ln^4(1-z) \right] \right\} + \mathcal{O}(\alpha_s^6 \times (\log)^{11})$$

Hamberg, van Neerven, Matsuura, 1991

Florian, Mazzitelli, Moch, Vogt, '14

Kramer, Laenen, Spira, '96, Kidonakis '07

$$L_\mu = \ln \mu/Q$$

# Summary and outlook

- The LP threshold fact. & res. was developed in 1987/89, each part extended to higher accuracy later.
- We provide an NLP resummation of the leading logs in the soft-collinear effective theory.
- The LO divergences in the soft function are cancelled by an auxiliary soft function.
- There is no kinematic power correction.
- The resummed result has no leading log at the jet scale.
- At a general scale, we reproduce the first few orders.

# Summary and outlook

- Extension to NLL is interesting and will reveal the full difficulty and complexity of NLP resummation, which can be seen from the anomalous dimension of NLP operators. M.Beneke, M.Garny, R.Szafron, JW '18

# Summary and outlook

- Extension to NLL is interesting and will reveal the full difficulty and complexity of NLP resummation, which can be seen from the anomalous dimension of NLP operators. M.Beneke, M.Garny, R.Szafron, JW '18

Thank you for your attention!