



Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector

BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

# Baryon form factors at BESIII

Lei Xia (on behalf of BESIII Collaboration)

[jessemcc@mail.ustc.edu.cn](mailto:jessemcc@mail.ustc.edu.cn)

University of Science and Technology of China, Hefei 230026,  
Peoples Republic of China

State Key Laboratory of Particle Detection and Electronics, Beijing  
100049, Hefei 230026, Peoples Republic of China

16<sup>th</sup> Conference on Flavor Physics and CP Violation (FCPC 2018)

July 14th 2018, Indian Institute of Technology Hyderabad and  
University of Hyderabad, Hyderabad, India

BESIII



SFB 104  
THE LOW-ENERGY FRONTIER  
OF THE STANDARD MODEL

JGU  
JOHANNES  
GUTENBERG  
UNIVERSITÄT MAINZ

HIM  
Helmholtz-Institut Mainz



中国科学院大学  
U.S.T.C.  
1958-2018  
60th Anniversary





# Outline

Baryon form factors at BESIII

Lei Xia

Introduction

Preamble  
NEFF

Accelerator and Detector

BEPCII

BESIII

Data set

Measurement of baryon FF at BESIII

Proton FF

$\Lambda$  FFs

$\Lambda_c$  Xsec

Summary

Discussion

Future



## 1 Introduction

## 2 Accelerator and Detector

## 3 Measurement of baryon form factors at BESIII

## 4 Summary



# Outline

Baryon form factors at BESIII

Lei Xia

Introduction

Preamble  
NEFF

Accelerator and Detector

BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

## 1 Introduction

- Preamble
- Nucleon Electromagnetic Form Factor

## 2 Accelerator and Detector

- Beijing Electron Positron Collider
- Beijing Spectrometer III
- Data Set

## 3 Measurement of baryon form factors at BESIII

- Measurement of proton form factor on BESIII
- Measurement of  $\Lambda$  form factor on BESIII
- Measurement of  $\Lambda_c$  cross section on BESIII

## 4 Summary

- Discussion
- Future



# Preamble

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector

BEPCH

BESIII

Data set

Measurement of baryon FF at BESIII


Proton FF

$\Lambda$  FFs

$\Lambda_c$  Xsec

Summary  
Discussion  
Future

- Baryon mass is the **main component** of the mass of the universe. It comes from the **strong force**, not from the Higgs mechanism. (K.Huang, *Story of Gauge Fields*, 2007, F. Wilczek, *A beautiful question*, 2016).



- Baryons, what they really are, is far from being understood.
- Many meson features come from **QED to QCD**, once  $\alpha \rightarrow \alpha_s$ . Baryon: no analogue in QED and **unique QCD feature**.
- For instance:
  - ✓ A fermion with mass, magnetic moment and other parameters close to proton and neutron ones can be obtained as a soliton of a  $\pi$  point-like boson field, by means of a non linear Lagrangian with one free parameter only (**Skyrme model**, *Proc. Roy. Soc. A* **260**, (1961), 127)!
  - ✓ The baryon spin is not due to the spins of the valence quarks (**Proton Spin Crisis**, *PLB* **206**, 364, (1988))!
- Therefore it is meaningful to point out open questions, concerning baryon structure.



# Nucleon Electromagnetic Form Factor

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector

BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$ ,  $\Sigma$ sec

Summary  
Discussion  
Future

- Elastic scattering of electron and proton (Phys. Rev. **98**, 217 R. Hofstadter, Nobel Prize 1961).

✓ Theoretically, differential cross section is:

$$\left(\frac{d\sigma}{d\Omega}\right)_{ep} = \left(\frac{d\sigma}{d\Omega}\right)_{Mott}(1 + 2\tau \tan^2 \frac{\theta}{2})F(q^2)$$

✓ The deviation represents the effect of a form factor (FF) for the proton.

- The nucleon electromagnetic vertex  $\Gamma_\mu$  describing the hadron current:



$$\Gamma_\mu(p', p) = \gamma_\mu F_1(q^2) + \frac{i\sigma_{\mu\nu}q^\nu}{2m_p} F_2(q^2)$$

- Sachs FFs:

$$\text{ElectronFF} : G_E(q^2) = F_1(q^2) + \tau \kappa_p F_2(q^2)$$

$$\text{MagnetFF} : G_M(q^2) = F_1(q^2) + \kappa_p F_2(q^2)$$

$$\text{where } \tau = \frac{q^2}{4m^2}, \kappa = \frac{g-2}{2} \text{ and } g = \frac{\mu}{J}$$




- In the Breit frame:  
Nucleon spin flip:  $G_M$ ,  
non spin flip:  $G_E$





# Nucleon Electromagnetic Form Factor

- Measurement of baryon FF: **Space-like (SL)** and **Time-like (TL)**.



Form Factors	
Dirac:	$F_1(q^2)$
Pauli:	$F_2(q^2)$
$G_E = F_1 + \frac{\kappa q^2}{4M^2} F_2$	
$G_M = F_1 + \kappa F_2$	



- TL** process includes **energy scan** and **initial state radiation (ISR)**, both techniques can be used at BESIII.

	Energy scan	Initial state radiation
$E_{beam}$	discrete	fixed
$\mathcal{L}$	Low at each beam energy	High at one beam energy
$\sigma$	$\frac{d\sigma_{p\bar{p}}}{d\cos\theta} = \frac{\pi\alpha^2\beta C}{2q^2} [ G_M ^2(1 + \cos^2\theta) + \frac{4m_p^2}{q^2}  G_E ^2 \sin^2\theta]$	$\frac{d\sigma_{p\bar{p}\gamma}}{dq^2 d\theta_\gamma} = \frac{1}{s} W(s, x, \theta_\gamma) \sigma_{p\bar{p}}(q^2)$ $W(s, x, \theta_\gamma) = \frac{\alpha}{\pi x} \left( \frac{2-2x+x^2}{\sin^2\theta_\gamma} - \frac{x^2}{2} \right)$
$q^2$	Single at each beam energy	From threshold to $s$



# Accelerator and Detector

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

**Accelerator and Detector**

BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

## 1 Introduction

- Preamble
- Nucleon Electromagnetic Form Factor

## 2 Accelerator and Detector

- Beijing Electron Positron Collider
- Beijing Spectrometer III
- Data Set

## 3 Measurement of baryon form factors at BESIII

- Measurement of proton form factor on BESIII
- Measurement of  $\Lambda$  form factor on BESIII
- Measurement of  $\Lambda_c$  cross section on BESIII

## 4 Summary

- Discussion
- Future







# BEijing S spectrometer III

Baryon form factors at BESIII

Lei Xia

Introduction

Preamble  
NEFF

Accelerator and Detector

BEPCH


BESIII

Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future



## ■ Main Drift Chamber (MDC): (He/C<sub>3</sub>H<sub>8</sub>=60/40)

- $\sigma_{xy} \approx 130 \mu\text{m}$ ,  $dE/dx \sim 6\%$ ;
- $\sigma_p/p \approx 0.5\%$  at 1 GeV.

## ■ Time Of Flight (TOF): (plastic scintillator)

- $\sigma_{time}(\text{barrel}) \approx 80 \text{ ps}$ ,
- $\sigma_{time}(\text{endcap}) \approx 65 \text{ ps}$ .

## ■ ElectroMagnetic Calorimeter (EMC): (CsI(Tl))

- $\sigma_E/E(\text{barrel}) \approx 2.5\%$  at 1 GeV,
- $\sigma_E/E(\text{endcap}) \approx 5\%$  at 1 GeV.

## ■ Superconducting Magnet: B = 1T.

## ■ Muon Counter: Resistive Plate Chambers (RPC):

- barrel: 9 layers;
- endcap: 8 layers.
- $\sigma_{spatial} = 2 \text{ cm}$ .



# Data set

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector

BEPCH


BESIII

Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future



- World largest data for
  - Charmonium spectroscopy
  - Charm physics
  - $\tau$  and R-QCD physics
  - Light hadrons
  - New physics research



# Measurement of baryon form factors at BESIII

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector  
BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

## 1 Introduction

- Preamble
- Nucleon Electromagnetic Form Factor

## 2 Accelerator and Detector

- Beijing Electron Positron Collider
- Beijing Spectrometer III
- Data Set

## 3 Measurement of baryon form factors at BESIII

- Measurement of proton form factor on BESIII
- Measurement of  $\Lambda$  form factor on BESIII
- Measurement of  $\Lambda_c$  cross section on BESIII

## 4 Summary

- Discussion
- Future



# Measurement of proton form factor on BESIII

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF




Accelerator and Detector

BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future



## ■ Energy scan technique:

- 2012 data,  $156.9 \text{ pb}^{-1}$ : PRD 91, 112004 (2015).
- 2015 R-scan data,  $688.5 \text{ pb}^{-1}$ : under reviewing.

## ■ ISR techniques:

- Tagged ( $7.4 \text{ fb}^{-1}$  above  $3.773 \text{ GeV}$ ): under reviewing.
- Untagged ( $7.4 \text{ fb}^{-1}$  above  $3.773 \text{ GeV}$ ): preliminary results.

## ■ Precision: Improved!

In TL region, our results are unprecedented precision.

Especially  $|G_E/G_M|$  providing an uncertainty comparable to the SL region for the first time.



# Some unexpected features are proved

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector


BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII



Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

- A step in the cross section is very likely due to **Coulomb**, since in the **Coulomb factor** there is a factor  $1/\beta$  that cancels the factor beta in the cross section formula and produces a step.



- Andrea Bianconi and Egle Tomasi-Gustafsson [PRL 114, 232301 \(2015\)](#) discovered the **oscillations** in effective FF ( $|G|$ ) from BABAR [PRD 87, 092005 \(2013\)](#) and [PRD 88, 072009 \(2013\)](#), which was confirmed by BESIII.



- Plateau above threshold**, corresponding to  $|G|$  close to 1, like a point-like fermion, similar features is also shown  $e^+e^- \rightarrow \Lambda_c^+ \bar{\Lambda}_c^-$ .



# Measurement of $\Lambda$ cross section on BESIII

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector  
BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

- 2012 data,  $2.63 \text{ pb}^{-1}$ , PRD 97, 032013 (2018).

- Neutral baryon: no Coulomb, but again a **step at threshold!**

- The observed threshold enhancement implies a more complicated underlying physics scenario.


- The **Coulomb Enhancement Factor**  $C(s) = \frac{\pi\alpha}{\beta(s)} \frac{1}{1 - e^{-\frac{\pi\alpha}{\beta(s)}}}$ , cancel

the  $\beta$  for a charged  $B\bar{B}$  pair, equals to 1 for a neutral  $B\bar{B}$  pair.

- Recalling the baryon pair production cross section:

$$\sigma_{B\bar{B}}(s) = \frac{4\pi\alpha^2\beta(s)C(s)}{3s} [ |G_M(s)|^2 + \frac{2m_p^2}{s} |G_E(s)|^2 ]$$

- Help to **understand the mechanism** of baryon production and test the theory hypotheses based on the threshold enhancement effect.





# Measurement of $\Lambda$ $G_E/G_M$ phase on BESIII

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector  
BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

- 2015 data,  $66.9 \text{ pb}^{-1}$ , preliminary result.

- Complex form of FFs:


- $G_E = |G_E|e^{i\phi_E}$ ,  $G_M = |G_M|e^{i\phi_M}$
- Relative phase:  $\Delta\phi = \phi_E - \phi_M$

- A non-zero phase has polarization effect on the Baryons:  $P_y \propto \sin \Delta\phi$

- With hyperon weak decay to  $B + P$ , the polarization of hyperon can be measurement, so does the relative phase between  $G_E$  and  $G_M$ !



- The angular distribution of daughter baryon from Hyperon weak decay:

- $\frac{d\sigma}{d\Omega} \propto 1 + \alpha_\Lambda \mathbf{P}_y \cdot \hat{\mathbf{q}}$ ,      •  $\alpha_\Lambda$ : asymmetry parameter.
- $\hat{\mathbf{q}}$ : unit vector along the daughter baryon in hyperon rest frame.



$$|G_E/G_M| = 0.94 \pm 0.16 \pm 0.03 \pm 0.02(\alpha_\Lambda),$$

$$\Delta\phi = 42^\circ \pm 16^\circ \pm 8^\circ \pm 6^\circ(\alpha_\Lambda).$$





# Measurement of $\Lambda_c$ cross section on BESIII

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector  
BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda_c$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

- 2014 data,  $631.3 \text{ pb}^{-1}$ : PRL 120, 132001 (2018).

- Ten modes of  $\Lambda_c^+$  ( $\bar{\Lambda}_c^-$ ) are reconstructed.

- Measurement of the  $\sigma_{Born}$  at 4 energy points below 4.6 GeV with unprecedented statistical accuracy ( $\sim 1.3\%$  at 4.6 GeV).


- The  $\sigma_{Born}$  at near the threshold, indicates the complexity of production behavior of the  $\Lambda_c$ .

- At threshold, there is again a step in  $\sigma_{\Lambda_c^+\bar{\Lambda}_c^-}$ .

- Followed by a kind of a plateau.

- At threshold  $\sigma_{\Lambda_c^+\bar{\Lambda}_c^-}$  is close to the point-like value, once the Coulomb Enhancement Factor is taken into account:

$$\sigma_{\Lambda_c^+\bar{\Lambda}_c^-} (\text{point-like}) \approx \frac{\pi^2 \alpha^3}{2m_{\Lambda_c}} \approx 145 \text{ pb.}$$







# Summary

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector  
BEPCH  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary

Discussion  
Future

- 1 Introduction
  - Preamble
  - Nucleon Electromagnetic Form Factor
- 2 Accelerator and Detector
  - Beijing Electron Positron Collider
  - Beijing Spectrometer III
  - Data Set
- 3 Measurement of baryon form factors at BESIII
  - Measurement of proton form factor on BESIII
  - Measurement of  $\Lambda$  form factor on BESIII
  - Measurement of  $\Lambda_c$  cross section on BESIII
- 4 Summary
  - Discussion
  - Future



# Summary: Discussion

Baryon form factors at BESIII

Lei Xia

Introduction

Preamble  
NEFF

Accelerator and Detector

BEPCII  
BESIII  
Data set

Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary

Discussion  
Future

- Electromagnetic FFs provide a quantitative description of hadron structure and are basic observables of QCD.
- BESIII is unique in its capability to measure baryon FFs, from nucleons to  $\Lambda_c$  and use two complementary approaches: energy scan and ISR technique:
  - Proton FFs have been measured using a test energy scan of 2012 and 2015, for 2012 data have published (PRD 91, 112004 (2015)):
    - ✓ Precision improved:
    - ✓ 2015 R-scan data,  $688.5 \text{ pb}^{-1}$ : under reviewing.
    - ✓ In TL region, our results are unprecedented precision.
    - ✓ Especially  $|G_E/G_M|$  providing an uncertainty comparable to the SL region for the first time.
  - Very exciting results from tagged ISR on protons expected very soon, preliminary results on untagged ISR techniques.
  - Published results on  $\sigma$  and FFs from  $\Lambda$  (PRD 97, 032013 (2018)) and  $\Lambda_c$  (PRL 120, 132001 (2018)) close to threshold.
    - ✓ Preliminary results on  $\Lambda$  Electromagnetic FFs relative phase.



# Summary: Future

Baryon form factors at BESIII

Lei Xia

Introduction  
Preamble  
NEFF

Accelerator and Detector

BEPCLII  
BESIII  
Data set


Measurement of baryon FF at BESIII

Proton FF  
 $\Lambda$  FFs  
 $\Lambda_c$  Xsec

Summary  
Discussion  
Future

## ■ Near future:

- **Present theory is missing something.**
- **Proton**: more data from CMD3 and BESIII.
- **$\Lambda$  and  $\phi K^+ K^-$** : more data around  $\Lambda\bar{\Lambda}$  threshold.
- **$\Lambda_c$** : more data at threshold and above by BESIII.
- **Neutron**: more data from SND, CMD3. [Publication by BESIII.](#)
- **$Br(J/\psi \rightarrow \gamma n\bar{n})$** : [Publication by BESIII.](#)
- **$G_E/G_M$  phase**: more data from BESIII.



Thanks all for hard work!  
Thanks for your attention!