## Phenomenology 2018 Symposium



Contribution ID: 557

Type: parallel talk

## Hybrid seesaw leptogenesis and TeV singlets (I)

Tuesday 8 May 2018 14:30 (15 minutes)

The appealing feature of inverse seesaw models is that the Standard Model (SM) neutrino mass emerges from the exchange of TeV scale singlets with sizable Yukawa couplings, which can be tested at colliders. However, the tiny Majorana mass splitting between TeV singlets, introduced to accommodate small neutrino masses, is left unexplained. Moreover, we argue that these models suffer from a structural limitation that prevents a successful leptogenesis if one insists on having unsuppressed Yukawa couplings and TeV scale singlets. We propose a hybrid seesaw model, where we replace the mass splitting with a coupling to a high scale seesaw module including a TeV scalar. We show that this structure achieves the goal of filling both the above gaps with couplings of order unity. In this talk, I will briefly describe the model, while the companion talk will discuss leptogenesis from this model in detail.

## Summary

**Primary authors:** Prof. AGASHE, Kaustubh (University of Maryland); DU, Peizhi; EKHTERACHIAN, Majid (University of Maryland); Dr FONG, Chee Sheng (Universidade de São Paulo); HONG, Sungwoo (University of Maryland-College Park); VECCHI, Luca (EPFL - Ecole Polytechnique Federale Lausanne (CH))

Presenter: DU, Peizhi

Session Classification: Neutrinos II