IDM 2018



Contribution ID: 79

Type: Talk

Results on sub-GeV dark matter direct detection with LUX Run 3 data by using Bremsstrahlung and Migdal-effect signal

Tuesday 24 July 2018 15:20 (20 minutes)

Dual-phase xenon time projection chambers have been recently suggested to be sensitive to sub-GeV dark matter if the inelastic channels of Bremsstrahlung and the Migdal effect in the nuclear interaction are taken into account. Sub-GeV dark matter is difficult to probe due to the small energy transfer in dark matter-nucleus elastic scattering and the finite energy threshold of the detector. Considering photon emissions from Bremsstrahlung or electron emissions from Migdal effect in the dark matter nucleus inelastic scattering could circumvent this difficulty. This analysis utilizes both the scintillation (S1) and ionization (S2) signals in the LUX Run 3 data. New sub-GeV parameter space in the dark matter- nucleus interaction cross-section is explored using the Profile Likelihood Ratio method.

Authors: LIN, Junsong (Lawrence Berkeley National Laboratory); TVRZNIKOVA, Lucie (Yale/LBNL)

Presenter: LIN, Junsong (Lawrence Berkeley National Laboratory)

Session Classification: 2.3 Direct Detection

Track Classification: Direct Detection