



# Exotica and Dark Matter searches


Michele Gallinaro

*LIP Lisbon*

May 2, 2018

- ✓ Introduction
- ✓ Dark matter
- ✓ Exotica searches

# 2012: A new boson discovery




M. Gallinaro - "Exotica and Dark Matter searches" - May 2, 2018







# Standard Model theory of everything?

- Discovery of the Higgs boson marks the triumph of the SM
- However, even with the inclusion of the Higgs boson, SM is an incomplete theory



# Tests of the SM






# Beyond the Standard Model

The SM answers many of the questions about the structure of matter. But SM is not complete; still many unanswered questions:

- a) Why do we observe matter and almost no antimatter if we believe there is a symmetry between the two in the universe?
- b) What is this "dark matter" that we can't see that has visible gravitational effects in the cosmos?
- c) Are quarks and leptons actually fundamental, or made up of even more fundamental particles?
- d) Why are there three generations of quarks and leptons? What is the explanation for the observed pattern for particle masses?
- e) How does gravity fit into all of this?

# Dark matter and energy


- What is that accounts for 96% of the Universe?  
Nobody knows.
- It is one of the greatest mysteries of Science





# What can we look for?

A crowded field. At the LHC we can search for some of these



# How?

- Search for new phenomena
- Look for New Physics
- **Indirect searches**
  - precision measurements, event properties, etc.
- **Direct searches**
  - resonances, specific final states, model-(in)dependent searches, etc.
- Production and decay rates, event characteristics, advanced tools





# Dark Matter

## What is it?

- DM does not interact electromagnetically
- DM interacts gravitationally

Visual map



From P. Harris DM talk at Cern (July 2015)

# Dark Matter (cont.)

## Why is it interesting?

- We do not see it...but we feel it

Mass map




From P. Harris DM talk at Cern (July 2015)




# Dark Matter (cont.)

## How do we find DM?

- Need to understand how it interacts with Universe
- Traditionally through a mediator
- Yields at least two new particles




# Searching for DM



Stable(-ish) particles:

- Anti-nuclei
- Photons
- Anti-protons
- Positrons
- neutrinos



BSM:



- Supersymmetry, neutralinos, gravitinos
- Extra-dimensions
- Axions(-like) particles
- Sterile neutrinos

# Searching for DM




## Direct Detection



## Particle Colliders




## Indirect Detection






# How do we find it: @LHC

- Produced it through a mediator





# How do we find it: @underground

- Through a nuclear recoil



# How do we find it: @Space

- Through annihilation
  - Cosmic rays from DM






# How do we find it: @nearUniverse



- Back and forth, production and annihilation
- Measure density and set constraints

Early Universe



Relic density constraint

# Experimental Results




⇒ Direct searches less sensitive to low masses due to energy threshold on nuclear recoil


# Collider searches

## Weakly interacting massive particles

- Effective field theory, simplified models



## Model-dependent searches




Searches for particles stable within detector acceptance,  
sensitive to mediator mass



# DM at the LHC


- CMS/ATLAS experiments **not** designed for DM searches



# DM searches at LHC

How do we find DM at the LHC?


- DM production gives MET signature



# DM searches at LHC

How do we find DM at the LHC?

- Mono-photon: Can also tag events with a photon






# DM searches at LHC

How do we find DM at the LHC?


- Mono-V: Tag events with a boson



# DM searches at LHC

How do we find DM at the LHC?

- Mono-V with (pseudo-) scalars
  - Different production mechanisms




# DM searches: backgrounds

What are the backgrounds?

- $Z \rightarrow \nu\nu$


–very similar to signal



# DM searches: backgrounds (cont.)

How to discriminate signal against the background?

- Look for high MET:



Study hadronic recoil

$$MET = -\sum_{\text{All particles}} p_T$$

$$MET(Z \rightarrow \nu\nu) = -Z \text{ recoil} + p_T(\nu\nu)$$


$$MET(Z \rightarrow \nu\nu) = -Z p_T$$



# DM searches: backgrounds (cont.)

How to discriminate signal against the background?


- Cut and count events or...



# DM searches: backgrounds (cont.)


How to discriminate signal against the background?

- Can fit the shape and look for signal



# Build a V-tagger


- Two jets are more collimated at high  $p_T$




- At **low  $p_T$**  jets are “resolved”
  - Focus on reconstructing di-jets with mass near W mass
- At **high  $p_T$**  get one “fat” jet
  - Focus on identifying one jet with mass near W mass
- Use additional variables to improve discrimination

# DM+Z


signal



background



- Main background is from ZZ di-bosons
- Understanding ZZ di-boson  $p_T$  is critical






# DM+jets ( $j/V/\gamma$ )

CMS-EXO-16-030

- Search focused on light mass region (100-300 GeV)
- Experimental challenges
  - Large QCD background
  - Triggers





hole in collider dijet searches

# DM+jets (j/V/ $\gamma$ ): Motivation

CMS-EXO-16-030



- Search for  $Z'$  leptophobic vector
- Strategy:  $Z' \rightarrow qq$
- Multijet topology with high- $p_T$  jet
- Look at jet substructure
- Search for “bump” in jet mass distribution



# DM+jets (j/V/ $\gamma$ ): Analysis

CMS-EXO-16-030



- Signal region
  - $p_T > 500$  GeV
  - $\tau_{21}^{\text{DDT}} < 0.38$
  - lepton veto
- Soft drop mass  $m_{\text{SD}}$ : peaks at  $Z'$  mass
  - removes soft wide-angle radiation from jet
- QCD background estimated from sideband regions in data
- $\tau_{21}^{\text{DDT}}$  n-subjettiness: consistency with 2-prong structure
- $\tau_{21}^{\text{DDT}}$  defines “pass” or “fail” sidebands
  - Use “*transfer function*” from fail to pass region



# DM+jets (j/V/ $\gamma$ ): Results

CMS-EXO-16-030

- Jet has 2-prong sub-structure
- Identify jet substructure using  $\tau_{21}$
- Set limits on light  $Z' \rightarrow qq$  search (most sensitive at  $<140$  GeV)





- Search for low-mass boosted dijet resonances
- Explores uncovered regions
- Limits in  $Z'$  mass at low mass



# DM+photon

arXiv:1704.03848


- BSM theories predict events with photon+MET
- Small SM background





# DM+jet/V

CMS-EXO-16-048

## DM search in mono-jet/V




Need good control of systematics



# DM+Higgs



arXiv:1510.06218, arXiv:1506.01081

- Generic search:  $pp \rightarrow X + \text{MET}$
- Search for DM +  $h(\rightarrow bb)$
- Model-independent search
  - Signature:  $h(\rightarrow ZZ/bb/\gamma\gamma) + \text{MET}$
  - Simplified model with  $Z'$  or pseudo-scalar Higgs  $A(\rightarrow \chi\chi)$



DM particle ( $\chi$ ): can be scalar or fermion  
Pseudo-scalar Higgs A


## Signal events at large MET




# DM+Higgs (cont.)

arXiv:1703.05236

- DM search with  $H(\rightarrow bb, \gamma\gamma)$
- Model dependent search
- $Z'$  2HDM Model





- **No significant excess**
- Set limits for coupling  $g=0.8$







# BSM/SUSY searches

- Density of allowed supersymmetric models before and after Run 1




# Experimental results

- Limits for given couplings between SM and DM interaction
- **Competitive limits at low masses** wrt other experiments




# Experimental results (cont.)



# Search for heavy resonances

- Heavy BSM resonances ( $>1\text{TeV}$ ) may decay into SM bosons (W,Z, H)
- Several final states
- Experimental challenges
  - SM bosons decay mostly to quarks
  - Due to large Lorentz boost, decay products merge into single jet
  - Clustered within a large-cone jet ( $R=0.8$ )
- Look into jet substructure
  - **Jet “grooming”**: get rid of soft jet components from UE/pileup, keep constituents from hard scatter
  - Apply filters (mass drop, pruning, trimming)




# Jet grooming


arXiv:0802.2470

## Mass drop/filtering


- Identify approx. symmetric sub-jets (with smaller mass than sum)



**Mass drop:** define 2 sub-jets



**Filtering:** re-cluster  $j_1, j_2$  constituents






# Jet grooming (cont.)

arXiv:0912.1342, arXiv:0912.0033


## “Trimming”

- Uses kT algorithm to make subjets (subjets with  $p_T^i/p_T < \text{cut}$  removed)



## “Pruning”

- Recombine jet constituents, while veto wide-angle/softer constituents



# W, Z, H reconstruction



CMS-B2G-17-002

- Grooming and jet mass

- Pruning
- soft drop (stable w/pileup, and good jet mass resolution  $\sim 10\%$ )

- Vector boson tagging ( $V \rightarrow qq$ )


- n-subjettiness  $\tau_{21}$ : how consistent with 2 sub-jets
- Categorization according to purity: high ( $< 0.35$ ) and high ( $> 0.35$ )



# W, Z, H reconstruction (cont.)

CMS-B2G-17-002





- Higgs boson tagging ( $H \rightarrow b\bar{b}$ )
  - Double b-tagging
  - Exploit b-tagging to identify two b-quarks in same jet
  - Soft-lepton information
  - Combines tracking and vertexing in MVA



# Diboson resonances




CMS-HIG-16-034, B2G-16-022

- Search for resonance:  $X \rightarrow ZZ \rightarrow \ell\ell qq'$
- Use tools to identify jet substructure
  - N-subjettiness  $\tau_{21}$ :  $\tau_N \sim 1/d_0 \sum p_T$
  - Kinematic and flavor information to improve S-B separation
- Discriminant Z+JJ (using MELA)
- Upper limits on resonant spin-0/spin-2 hypotheses
- Cross section limits  $\sim 3\text{-}100\text{fb}$



# Heavy resonance: WH final state

- Search for massive resonance  $W' \rightarrow WH$
- Distinctive features of BSM models, i.e. composite/little Higgs, technicolor, etc.
- Lepton+jet final state
- Use jet substructure/btag for  $H \rightarrow bb$
- $2.2\sigma$  highest local significance at 1.8 TeV








# Search for multilepton final states

CMS-EXO-17-006


- Type-III extension to SM
- Search for 3 or more lepton final states
- Pair production of  $W/Z/H \rightarrow \Sigma\Sigma$
- Scalar sum of lepton  $p_T$  ( $L_T$ )
- Bin and count ( $L_T + \text{MET}$ )



# $X \rightarrow VV \rightarrow qqqq$


CMS-B2G-17-001

- All hadronic resonance search with single (qV) or double (VV) V-tag
  - At least 2 back-to-back jets  $p_T > 200 \text{ GeV}$
  - Categorization (jet mass,  $\tau_{21}$ )
- Background estimation: “bump hunt” fit data with power law




Candidate ZZ event  
Dijet mass: 3.2 TeV

Anti- $k_T$ R=0.8 jet	
$p_T$	1321 GeV
$\eta$	-0.40
$\phi$	-2.71
$M_{SD}$	103 GeV
$\tau_{21}$	0.23





Anti- $k_T$ R=0.8 jet	
$p_T$	1374 GeV
$\eta$	0.79
$\phi$	0.43
$M_{SD}$	94.8
$\tau_{21}$	0.29



# $X \rightarrow VH \rightarrow qqbb$

CMS-B2G-17-002

- All-hadronic search for  $V \rightarrow qq$  and  $H \rightarrow bb$  resonances
  - dedicated identification for  $H \rightarrow bb$  (b-tagging)
- Use categories
  - V-jet mass (W or Z), V-jet  $\tau_{21}$  (high-purity, low-purity), H-jet (tight and loose b-tag)






- Similar topology and background estimate to VV resonance search
- No significant excess found in data

# $X \rightarrow ZV \rightarrow \ell\ell qq$

CMS-B2G-16-022

- Search for resonances in  $Z \rightarrow ee/\mu\mu$ ,  $V \rightarrow qq$
- Clean final state (leptons)
  - Good mass resolution, good efficiency
- $\tau_{21}$  categorization (HP, LP)
- Parametrize main bkg ( $Z$ +jets), fit to data in sidebands, take shape from MC





• Data compatible with SM-only hypothesis

# $X \rightarrow WV \rightarrow \ell\nu qq$

CMS-B2G-16-020

- Search for a resonance decaying to  $WV$  in leptonic channel
- Categorization in  $\tau_{21}$  and  $W/Z$  mass
- Sideband+transfer function for bkg estimate



- Similar sensitivity to  $Z(\ell)V(qq)$  search
- Excluded up to 2 TeV






# $X \rightarrow VH \rightarrow \ell\nu qq$



PLB 768(2017)137

- Search for a resonance decaying to VH in leptonic channels
  - $Z \rightarrow \nu\nu$ : transverse mass  $m_T(VH)$
  - $W \rightarrow \ell\nu$ : top control region
  - $Z \rightarrow \ell\ell$ : high-efficiency dilepton ID
  - $H(bb)$  b-tagging
- Sideband bkg prediction

- Heavy vector triplet ( $Z'$ ,  $W'$ )
- $g_V, g_H$  ( $c_V, c_F$ ): couplings






# Combination of diboson searches



# Di-muon events



- Di-muon events: a re-discovery of the SM





# Dilepton resonance

arXiv:1803.06292






- Search for dilepton ( $ee, \mu\mu$ ) resonance



# Resonance searches: Summary



**CMS Preliminary**







# Looking forward: PPS

CERN-LHC-2014-021

- The Precision Proton Spectrometer is a joint CMS and TOTEM project that aims at measuring the surviving **scattered protons** on both sides of CMS in standard running conditions
- **Tracking** and **timing** detectors inside the beam pipe at ~210m from IP5
- Project approved in Dec. 2014 by LHCC
- Data taking started in 2016 (full scope from 2017)




CERN-LHC-2014-021



# PPS physics motivations

- **Central Exclusive Production**
  - photon-photon collisions
  - gluon-gluon fusion in color singlet,  $J^{PC}=0^+$
- **High- $p_T$  system in central detector, together with very forward protons in PPS**
  - momentum balance between central system and forward protons, provides strong kinematical constraints
  - Mass of central system measured by momentum loss of the two leading protons
- **Gauge boson production by photon-photon fusion and anomalous couplings ( $\gamma\gamma WW$ ,  $\gamma\gamma ZZ$ , and  $\gamma\gamma\gamma\gamma$ )**
- **Search for new BSM resonances**
- **Study of QCD in a new domain**





# LHC tunnel @ PPS location

215m

CT-PPS  
timing


214m

CT-PPS  
tracking

beam

220m

CT-PPS  
tracking





# PPS detectors



- Tracking detectors

- Goal: measure proton momentum
- Technology: silicon 3D pixels (6 planes per pot)

- Timing detectors

- Goal: identify primary vertex, reject “pileup”
- $\sigma_{\text{time}} \sim 10\text{ps} \Rightarrow \sigma_z \sim 2\text{mm}$
- Technology: silicon/diamond



“3D” pixel sensors with columnar electrodes



# Timing detectors

Use timing to reject pileup background

- Two scenarios studied:
  - 10ps and 30ps time resolution
- **Baseline: solid state detectors**
- **Detector options investigated:**
  - Diamond sensors
  - Fast silicon sensors (UFSD, HFS)
- **Status:**
  - Diamond and LGAD detectors installed






# WW production

JHEP 08(2016)119

- Study of process:  $pp \rightarrow pWWp$ 
  - Clean process: W in central detector and “nothing” else, intact protons can be detected far away from IP
  - Exclusive production of W pairs via photon exchange: QED process, cross section well known




- Backgrounds:
  - inclusive WW,  $\tau\tau$ , exclusive two-photon  $\gamma\gamma \rightarrow ll$ , etc.
- Events:
  - WW pair in central detector, leading protons in PPS

- SM observation of WW events

- Anomalous coupling study

- AQGCs predicted in BSM theories
- parameters:  $a_0^W/\Lambda^2$ ,  $a_c^W/\Lambda^2$





- Deviations from SM can be large

# AQGC expected limits

Expected limits @95%CL:

$$a_0^W / \Lambda^2 = 2 \times 10^{-6} \quad (3 \times 10^{-6}),$$


$$a_C^W / \Lambda^2 = 7 \times 10^{-6} \quad (10 \times 10^{-6}).$$




# Exclusive Dileptons

CMS-PPS-17-001


- Study exclusive processes at the EWK scale
- Search for two-photon production of opposite charge lepton pair with forward proton tagging



signal



SD



DD



Background: SD, DD, DY, dibosons, PU

- Signal selected with:
- at least one proton tagged, muons, kinematic selection

# Exclusive Dileptons (cont.)

CMS-PPS-17-001

- Correlation between the  $\xi$  values in central system vs RP
- $12\mu\mu$ ,  $8ee$  candidates observed ( $>5\sigma$  over expected bkg)
- First observation of two-photon production of a lepton pair at this mass range



# BSM searches: resonances, etc.


CMS-EXO-15-004, CERN-LHC-2014-021





CMS Experiment at the LHC, CERN  
Data recorded: 2015-Sep-11 22:46:54.589056 GMT  
Run / Event / LS: 256353 / 437637379 / 244

diphotons at PPS

$\sigma \sim 0.3 \text{ fb}$  a few 'clean' events with 20/fb




exclusive WW production





# Increased reach at 13 TeV



# Summary

- Excellent consistency of SM but **SM is incomplete**
- Direct and indirect searches for New Physics
  - Collected  $\sim 80/\text{fb}$  @13 TeV in 2015-2017
  - $\sim 300/\text{fb}$  to be collected in the next few years (up to LS3)
- Many studies performed with data collected so far
  - New dedicated algorithms being developed
  - Dark Matter, Exotica, signature-based searches
  - Other BSM searches
- Searches provide **no hints for BSM yet**



# backup



# Exotica and Dark Matter searches


Michele Gallinaro

*LIP Lisbon*

May 3, 2017

- ✓ Introduction
- ✓ Dark matter
- ✓ Exotica searches





# Exotica and Dark Matter searches

Michele Gallinaro

*LIP Lisbon*


May 3, 2017

- ✓ Introduction
- ✓ Dark matter
- ✓ Exotica processes
- ✓ Dilepton and diboson processes



# Resolved V-tagger


- For low  $p_T$  objects focus on **di-jet properties**



- Color flow, jet quark/gluon likelihood, mass drop

# Boosted V-tagger





- For high  $p_T$  objects focus on **jet sub-structure**



• ...





# Tests of the SM

- Various tests provide excellent self-consistency of the SM



# Tests of the SM

- Various tests provide excellent self-consistency of the SM



# Invisible Higgs as a portal to Dark Matter

ATLAS at  $300 \text{ fb}^{-1}$

➤ Indirect constraints on  $\text{BR}(H \rightarrow \text{inv})$ :

- from Higgs coupling fit
- $\text{BR}(H \rightarrow \text{inv}) < 28\% \text{ @ } 95\% \text{ CL}$

➤ Direct search



- $ZH \rightarrow ee/\mu\mu + E_{\text{miss}}$
- $\text{BR}(H \rightarrow \text{inv}) < 32\% \text{ @ } 95\% \text{ CL}$

➤ Possible to **convert** the limits on  $\text{BR}(H \rightarrow \text{inv})$  into the strength of the interaction between dark matter and Higgs boson,  $\lambda_{hcc}$

➤ Bound on  $\lambda_{hcc}$  can be mapped into scattering cross section of dark matter on a nuclei

➔ comparison with direct searches

- Limits from ATLAS at low mass **better** than those from direct detection limits
- degrade as  $m_c$  approaches  $m_h/2$




ATL-PHYS-PUB-2013-014




# Mono-higgs

- Higgs discovery provides new portal into DM coupling to SM
- DM searches at the LHC include analyses with mono- $X$  + MET signatures for  $X=W, Z, \text{jet}, \text{and } \gamma$ 
  - ATLAS 8 | TeV mono- $H \rightarrow \gamma\gamma$
- In general,  $X$  can be emitted as ISR or from the new vertex coupling DM to SM
- Higgs ISR is highly suppressed, so mono- $H$  can directly probe the effective DM-SM coupling



# Mono-higgs models


- Models consist of the union of models from Phenom papers [arXiv:1312.2592](https://arxiv.org/abs/1312.2592), [arXiv:1402.7074](https://arxiv.org/abs/1402.7074) and [ATLAS-CMS DM Forum](#), with phenomenology studies for new models coming.
- Six EFTs: dimension 4 to 8 contact operators valid below cutoff scale  $\Lambda$




$$\lambda |H|^2 \chi^2 \quad \frac{1}{\Lambda} |H|^2 \bar{\chi} \chi \quad \frac{1}{\Lambda} |H|^2 \bar{\chi} i \gamma_5 \chi \quad \frac{1}{\Lambda^2} \chi i \partial^\mu \chi H^\dagger i D_\mu H$$

$$\frac{1}{\Lambda^2} \chi^\dagger i \overleftrightarrow{\partial}^\mu \chi H^\dagger i D_\mu H \quad \frac{1}{\Lambda^4} \bar{\chi} \gamma^\mu \chi B_{\mu\nu} H^\dagger D^\nu H.$$



- Four simplified models: new massive mediator –  $Z'$ ,  $S$ ,  $A^0$  – for Higgs-DM coupling



larger cross section



# Experimental Results



- Competitive limits at low masses

# Summary for Higgs exotic decays

