

Luminosity Spectrum Reconstruction - Impact of Detector Resolution Mismodelling

Philipp Zehetner
Ludwig-Maximilians-Universität München

CERN Summer Student
Supervised by Esteban Fullana and Andre Sailer

August 28, 2018

What if we don't know the exact resolution of our detector?

What if we don't know the exact resolution of our detector?

Monte Carlo generator: 10M e^+e^- pairs


What if we don't know the exact resolution of our detector?

Monte Carlo generator: $10\text{M } e^+e^-$ pairs

BHWIDE


$10\text{M } e^+e^-$ pairs after Bhabha scattering

What if we don't know the exact resolution of our detector?




Analysis Steps


What if we don't know the exact resolution of our detector?




Plot Structure



Plot Structure




Plot Structure



χ^2 results for Θ and Energy resolution scan

Θ


Chi²/ndf of fit



31 × 31 = 961 data points

Energy


Chi²/ndf of Fit




41 × 41 = 1681 data points

$$\sigma_E = \frac{a}{\sqrt{E}} \oplus b$$


Impacts on top mass measurements: Theta




Impacts on top mass measurements: Theta



Luminosity Spectrum for mismodelling Θ




Problematic Missmodelling: Energy resolution



- 1 'Stable zone' around diagonal: small error and little deviation
- 2 off-diagonal corners: large deviation, errors drop to zero


Impacts on top mass measurements: Energy

Chi²/ndf of Fit




Impacts on top mass measurements: Energy


Chi²/ndf of Fit




Impacts on top mass measurements: Energy



Impacts on top mass measurements: Energy



How reliable are the 8 MeV?



- 1 If the resolution on Θ is reasonably well modelled, resulting errors on the top quark mass measurement are small


- 1 If the resolution on Θ is reasonably well modelled, resulting errors on the top quark mass measurement are small
- 2 The energy resolution is more sensitive.

- 1 If the resolution on Θ is reasonably well modelled, resulting errors on the top quark mass measurement are small
- 2 The energy resolution is more sensitive.
- 3 Errors stay relatively small, when mismodelling the resolution by $\pm 15\%$

- 1 If the resolution on Θ is reasonably well modelled, resulting errors on the top quark mass measurement are small
- 2 The energy resolution is more sensitive.
- 3 Errors stay relatively small, when mismodelling the resolution by $\pm 15\%$
- 4 Larger deviations either drastically increase the error, or yield unreliable results on error propagation


- ① Luminosity Spectrum Reconstruction at Linear Colliders
Authors: Stéphane Poss, André Sailer
<https://arxiv.org/abs/1309.0372>
- ② Errors on top quark mass provided by Pablo Gomis
Link to his Presentation

BackUp 1: χ^2 for larger range for Θ




BackUp 2: χ^2 for parameter b

Chi²/ndf of Fit




BackUp 3: Luminosity: 50vs80

Lumi




BackUp 4: Luminosity: 80vs50

Lumi




BackUp 5: Luminosity: 200vs280

Lumi




BackUp 6: Luminosity: 220vs260

Lumi




BackUp 7: Luminosity: 260vs220

Lumi



BackUp 8: Luminosity: 260vs250

Lumi



BackUp 9: Luminosity: 220vs230

Lumi

