



Contribution ID: 21

Type: **not specified**

Amplitude Analysis of D decays to three hadrons in the LHCb experiment

Wednesday, May 9, 2018 2:30 PM (15 minutes)

The study of a heavy meson decaying to three lighter mesons can be described, in general, as quasi-two-body processes through the production of resonance intermediate states. To understand the dynamics of these processes, a full amplitude analysis of the corresponding Dalitz Plot (the two-dimensional representation of the decay phase space) is necessary. The most naïve way to describe it is to use the so-called Isobar Model where the total amplitude is written as a coherent sum of the individual resonance amplitudes, typically described as a product of the resonance propagator, angular functions and form factors. However, the Isobar Model turns out to be inadequate when dealing with broad scalar states and other approaches such as a model-independent partial wave analysis, K matrix, are interesting alternatives.

We show the current status of the amplitude analyses of the D decays into three mesons being pursued by the Rio Charm Group at the LHCb experiment.

Authors: CORREA DOS REIS, Alberto (CBPF - Brazilian Center for Physics Research (BR)); GOBEL BURLAMAQUI DE MELLO, Carla (Pontifical Catholic University of Rio de Janeiro (BR)); POLYCARPO MACEDO, Erica (Federal University of Rio de Janeiro (BR)); GONCALVES ABRANTES, Fernanda (Pontifical Catholic University of Rio de Janeiro (BR)); BAPTISTA, Juan (Universidade do Estado do Rio de Janeiro (BR)); AMATO, Sandra (Univ. Federal do Rio de Janeiro (BR))

Presenter: GONCALVES ABRANTES, Fernanda (Pontifical Catholic University of Rio de Janeiro (BR))

Session Classification: Short Communications