

# Higgs Physics

Paolo Francavilla - Istituto Nazionale di Fisica Nucleare, Pisa  
24/7/2018 - HASCO 2018 - Göttingen




Istituto Nazionale di Fisica Nucleare

# Outline

---


- **Particles in the Standard Model and their masses**
- **Condensate and the spontaneous breaking of symmetry**
- **Masses and the condensate**
- **Higgs boson**
- **Discovering the Higgs boson at the LHC**
- **Selected news from Run2**
- **Higgs Properties**
- **Bonus: Some important questions...**
- **Bonus: Back to the future**

# Particles in the Standard Model and their masses



# Particles in the Standard Model

## Standard Model of Elementary Particles



Imagine we see this for the first time  
What we would learn?

- Some organised way to map particles;
- Some coherency in spin and charge, with very well defined values;
- Mass is a mess...

Which is the allowed range for the mass of a fundamental particle?

---



Photons have 0 mass\*  $\Rightarrow$  Our scale starts from 0

Which is the allowed range for the mass of a fundamental particle?

---



Photons have 0 mass\*  $\Rightarrow$  Our scale starts from 0

Which is the allowed range for the mass of a fundamental particle?

---











Approaching the Plank mass, the gravitational effects  
will be more and more important  
Black Holes, so an object with different properties(?!)

1% of the mass of a mosquito

# Question time!!!








We have 16 particles (+ Higgs boson),  
between 0 and  $22 \mu\text{g}$ . Where are they?






# Question time!!!

We have 16 particles (+ Higgs boson),  
between 0 and 22  $\mu\text{g}$ . Where are they?




# Why?

known particles are in the range  $0-10^{-17} m_p$



# Why?

Maybe other particles out here,  
maybe we will find them in the coming years,  
maybe they are beyond the LHC reach



# What is special with these particles and their masses?


---

- We can write a version of the SM with all the fermions and vector bosons (without the Higgs boson/mechanism) in which all the particles are massless.
- This theory has all the nice features a theory could need (but one)
- and in part that could explain why they have a mass so small (it is like a small correction from 0)
- BUT:
  - it is not describing nature because particles have masses...

# Going back: what is mass?

---

empty box




>

=

<

?


box filled with photons





# Going back: what is mass?

**A system of massless constituents could have a mass**  
**In this case, it is an emergent phenomena**

empty box



box filled with photons




$$m = E/c^2$$


*which, by the way, is how Einstein wrote it in his paper*

# In the real World...


box filled with photons



neutron



Mass of the quarks is



of the mass of  
protons/neutrons

**~99% of our mass dynamically emerging from  
strong interactions**

# Particles in the Standard Model and their masses


- Compared to the Planck mass, the elementary particles are almost massless
- The mass can be an emergent property of a system
- It may be related to some underlying dynamics



# Condensate and the spontaneous breaking of symmetry

Use over time for: condensate

a relatively recent concept...




# Fields

- Fields characterises the behaviour of a quantity in that moment in that place
- They can vary from place to place, and they can evolve in time

2 m above gnd Temperature (C)


Wed 18.07.2018 11:00 CEST



- They could effect how things are moving, for example electric field
- Space can be filled with fields

# Fields


- Fields characterises the behaviour of a quantity in that moment in that place
- They can vary from place to place, and they can evolve in time



- They could effect how things are moving, for example electric field
- Space can be filled with fields

# Vacuum


- Ordinarily we think that fields are zero in empty space
- Is there a requirement in physics that says that this should be the case?
- What if our universe was filled with electric field?
  - i.e. capacitors placed infinitely far from us?
  - The electric field would just be there
  - We would experience charged particles moving in some peculiar way, but that would be just a fact of nature



# Energy of a field and vacuum

---


- In general, fields cost energy
- Space without an electric field has 0 energy
- Energy density of the electric field:
  - $u \propto E^2$
- From the Quantum Mechanical point of view, the vacuum is a state: the state of lowest energy...
- and the quanta of vibration of a field are particles



# Multiple fields

---

- Let's imagine we have 2 dim field :  
 $\Phi = (\phi_1, \phi_2)$




- The energy would depend on both the components
- No matter in which direction we displace the field, it costs energy
- If we have a paraboloid, to minimise the energy, the components of the field would be at the bottom of the “potential”.
- Must all the fields respect this parabolic shape?

# Paraboloid potential


---

- If the potential is symmetric, we could “move” the field in a circle.
- This motion of the field is very similar to angular momentum
  - It is not in space, it is in the internal field space
- But as the angular momentum, this is quantised.
- This corresponds to the quantisation of charge for that field.
- What if the potential energy is not a paraboloid?



# Mexican hat potential

- Maximum at the top of the hat.  
So,  $\phi = (0,0)$  is not the stable equilibrium
  - un-stable equilibrium
- The brim of the hat is where a ball placed on the top of the hat is going to go.
- The lowest energy state for the field would not be at zero field, but it would be in the brim.
- Interesting!!! The zero energy of the field is not at  $\phi = (0,0)$ , but on the rim
- Interesting vacuum: the value of the field in each point in space is not zero.
- How would we notice if we have this configuration?
- It may effect other things, and indeed it does...





# Something interesting is happening to the “spin”

---

- In the paraboloid, we had to excite the field to get it in rotation
  - This means that creating charged particle costs some energy
- And in the mexican hat?
  - We do not have to ride up the side of the hat to make a rotation.
  - it is for free in terms of potential energy!!!
  - With this potential, the field could “spin” for free in terms of potential energy in each place of the space.
  - This again correspond to a charge.
  - The entire space would have charge density for essentially no potential energy
  - This is known as condensate in space of charge



# Properties of this condensate

- **What if we want to find the lowest energy that the vacuum can have?**

- Best bet: make the field not move with time

- Remember, ride up the potential cost energy,

- but there is still some “kinetic” energy in moving in circle.


- *So minimum is should be standing steel?*

- **Problem:** uncertainty principle

- We know where the field is in the potential, but for the uncertainty principle, large uncertainty on how fast it is moving around the circle.

- **This means that in a condensate, we cannot have empty space with no charge in it**

- **Empty space is filled with a totally uncertain amount of charge.**




$$\Delta x \Delta p \geq \frac{\hbar}{2}$$

# Condensates

---

- What does it mean that the empty space is filled with a totally uncertain amount of charge?
  - Equal probability for the charge in a place in space to be 0, 1, 2, -1, -2, etc...
- What happens if we have an extra charged particle, and we trough it in?
  - It is not changing the probability of having a certain amount of charge.
  - It is the same from where we stated with.
- What if we take one charge out of this thing?
  - Again same status as before.
- In a condensate we would not really realise if we are putting a charge in or we are taking a charge out.
- The real word is not like that with respect to electric charge.
  - But this is what happens in superconductors! They are exactly like this!
- So, in nature, there are regions where the charge is totally uncertain, and we have condensate!



# Condensate and the spontaneous breaking of symmetry

- The vacuum can be filled with a condensate
- The condensate has a totally uncertain amount of charge
- It is a very weird beast....

# Masses and the condensate



Dirac theory of fermions:  
few "naive" considerations

---

$$\psi_L \equiv \frac{1}{2}(1 - \gamma_5)\psi$$

$$\psi_R \equiv \frac{1}{2}(1 + \gamma_5)\psi$$

Dirac Lagrangian

$$\mathcal{L}_{\text{Dirac}} = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi$$

$$\psi = \psi_L + \psi_R$$

Spinor represented  
with the chiral spinors

For massless fermions, the Dirac Lagrangian becomes:

$$\mathcal{L}_{\text{Dirac}} = \bar{\psi}_L(i\cancel{\partial})\psi_L + \bar{\psi}_R(i\cancel{\partial})\psi_R$$

It seems like these are two independent degrees of freedom in  
the particle zoo...

# Dirac theory of fermions: few "naive" considerations

---

- For 0 mass fermions, the two chiral spinors have a well defined physical meaning:
- They are the two possible projections of the spin in the direction of motion of the fermion - helicity.  
(This is not true anymore for massive fermions)

Right Handed

Left Handed



# Dirac fermions and interactions

---

- Are the LH and RH fermions completely equivalent in terms of interactions?
  - NO!!!!
- LH are charged for weak interaction
- RH are neutrals for weak interaction\*
- Why? One of the big mysteries.....

\*Note: After EWSB, RH are coupling with the  $Z^0$  as a result of the mixing of  $(W^0, B) \rightarrow (Z^0, A)$



# Dirac theory of fermions: few "naive" considerations

---

$$\psi_L \equiv \frac{1}{2}(1 - \gamma_5)\psi$$

$$\psi_R \equiv \frac{1}{2}(1 + \gamma_5)\psi$$

## Dirac Lagrangian

$$\mathcal{L}_{\text{Dirac}} = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi$$

$$\psi = \psi_L + \psi_R$$


Spinor represented  
with the chiral spinors

For massive fermions, the Dirac Lagrangian becomes:

$$\mathcal{L}_{\text{Dirac}} = \bar{\psi}_L(i\partial\!\!\!/)\psi_L + \bar{\psi}_R(i\partial\!\!\!/)\psi_R - m(\psi_L\psi_R + \psi_R\psi_L)$$


- The mass is in practise introducing an interaction between LH and RH fermions.
- In fact, the rate of flipping from LH to RH and back to LH is proportional to the mass.
- BUT, we have to conserve charge...

# Following a fermion



- $m_2 < m_1$
- But what about the conservation of the weak charge?

# Following a fermion



- $m_2 < m_1$
- But what about the conservation of the weak charge?
- No! We need some “magic” to conserve the weak charge...
- Fermion must be massless, apart from some magic


# Putting things together..

---


- Mass is proportional to the rate of flipping between LH-RH
- Flipping from RH-LH is not conserving the weak charge
- If we have a condensate, empty space is filled with a totally uncertain amount of (weak) charge.
- Maybe the RH-LH flip is allowed if the (weak) charge is continuously exchanged with the empty space.
- We are starting to give mass to the particles.....

# Fermions and their masses: The chiral symmetry breaking

---




- New ingredient: a boson (not yet the Higgs) from the condensate
- Recall, we can add one more to the condensate, or take one out, without changing the state of the condensate.
- So, the LH fermion can emit one of these bosons which carries its (weak) charge out.
- Where does this boson go?
  - To the condensate!
- And the fermion can borrow a charge back from the condensate...





# And the bosons?

## The Brout-Englert-Higgs Mechanism

---



- How the V boson get a mass? Something very similar
- The V boson can interact with particles with (weak) charge, so it can interact with these new bosons.
- How?
- **The Brout-Englert-Higgs mechanism**
- NOTE: if there was a condensate of ordinary particles, this would have happen to the photon...



# And the bosons?

## The Brout-Englert-Higgs Mechanism

The diagram illustrates the Brout-Englert-Higgs mechanism in three stages from top to bottom:

- Top stage:** Shows a photon field  $A^\mu$  and a Z boson field  $A^\nu$  interacting with Higgs vacuum expectation values  $\langle \phi_1 \rangle$ . The interaction term is  $e^2 g_{\mu\nu}$ . A gold medal and the text "F. Englert Séminaire Poincaré 2014" are shown.
- Middle stage:** Shows the mixing of the photon and Z boson fields. The photon field  $A^\mu$  has a coupling  $e q_\mu$  and the Z boson field  $A^\nu$  has a coupling  $-e q_\nu$ . The propagator is  $1/q^2$ .
- Bottom stage:** Shows the resulting mass matrix  $D_{\mu\nu}$  and the Z boson mass  $M_V^2 = e^2 \langle \phi_1 \rangle^2$ .

On the right side, a diagram shows the resulting Z boson and photon fields. The Z boson is represented by a blue arrow and the photon by a green arrow. The mass  $m_V$  is indicated.

**Equations:**

$$\Pi_{\mu\nu} = \left( g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) e^2 \langle \phi_1 \rangle^2$$


$$D_{\mu\nu} = \frac{g_{\mu\nu} - q_\mu q_\nu / q^2}{q^2 - e^2 \langle \phi_1 \rangle^2}$$

$$M_V^2 = e^2 \langle \phi_1 \rangle^2$$

- How the Z boson gets its mass.
- The Z boson and photon are mixed bosons.
- How?
- So, the Z boson has a mass.
- This is the Brout-Englert-Higgs mechanism.
- NOTE: if there were more Higgs fields...

# Back to the mexican hat

- We have two modes for the field in this potential:



- **Rotating on the brim, with no cost in potential energy**
  - This is causing our condensate, and the bosons related to this mode are the bosons entering in the mechanism seen before
- **Oscillating up-down hill**
  - This costs energy -> create this boson cost energy
  - This mode is like an “sound” wave of the density of the condensate
- **THE HIGGS BOSON**



# Masses and the condensate

- Thanks to the exchange of charges with the condensate, we have a way to give mass to the elementary particles:
  - It works for fermions
  - It works for bosons
- The model predicts the existence of a sound wave of the condensate:  
The Higgs boson

Higgs boson

# THE HIGGS BOSON



# The Standard Model

Հիմնական հասկացությունները: Գրգռվածություն, Կապակցվածություն, Կապակցվածություն, Կապակցվածություն

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i \bar{\psi} \not{D} \psi + \text{h.c.} + \bar{\psi}_i \gamma_{ij} \psi_j \phi + \text{h.c.} + |D_\mu \phi|^2 - V(\phi)$$

- Kinematic term of the gauge bosons
- Kinematic term of the fermions, and interaction between fermions and gauge bosons
- Higgs-fermions interaction
- Kinematic term of the Higgs boson, and interaction with the gauge bosons
- Higgs potential and self interaction

# The Standard Model

Հիմնականորեն: ժամանակակից  
ֆիզիկայի մեծագույնը

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i \bar{\psi} \not{D} \psi + h.c.$$

- Kinematic term of the gauge bosons
- Kinematic term of the fermions, and interaction between fermions and gauge bosons

$$+ \bar{\psi}_i \gamma_{ij} \psi_j \phi + h.c. + |D_\mu \phi|^2 - V(\phi)$$

- Higgs-fermions interaction
- Kinematic term of the Higgs boson, and interaction with the gauge bosons
- Higgs potential and self interaction

# Higgs sector in the Standard Model

---

- Probably the less elegant sector
  - Largest number of parameters

$$+ \bar{\Psi}_i y_{ij} \Psi_j \phi + \text{h.c.}$$
$$+ |D_\mu \phi|^2 - V(\phi)$$

# Higgs sector in the Standard Model

---

- Probably the less elegant sector
  - Largest number of parameters
  
- However:
  - It solves the issue of how masses in the standard model
  - Predict the relation between the masses and couplings of the gauge bosons
  - Predict the existence of the Higgs boson


# Higgs sector in the Standard Model

---

- Probably the less elegant sector
  - Largest number of parameters

- It solves another big issue:  
The unitarity of the longitudinal vector boson scattering.

- However:
  - It solves the issue of how masses in the standard model
  - Predict the relation between the masses and couplings of the gauge bosons
  - Predict the existence of the Higgs boson




# Higgs sector in the Standard Model

- Probably the less elegant sector
  - Largest number of parameters

- It solves another big issue:  
The unitarity of the longitudinal vector boson scattering.

- However:
  - It solves the issue of how masses in the standard model
  - Predict the relation between the masses and couplings of the gauge bosons
  - Predict the existence of the Higgs boson






# Higgs sector in the Standard Model

- Probably the less elegant sector
  - Largest number of parameters
- However:
  - It solves the issue of how masses in the standard model
  - Predict the relation between the masses and couplings of the gauge bosons
  - Predict the existence of the Higgs boson

- It solves another big issue:  
The unitarity of the longitudinal vector boson scattering.



In absence of a Higgs boson with  $m_H < 1$  TeV, would imply a strong dynamics which could be produced in the WW process

# Higgs sector in SM Lagrangian $\mathcal{L}_{\text{Higgs}} = (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi)$



- In what we discussed, we had some simplifications, being a bit more precise :)
- A new SU(2) doublet of spin-0 particles is added to the lagrangian, and interact with the W and B bosons

$$D_\mu \Phi = (\partial_\mu + ig\sigma^a W_\mu^a / 2 + ig' Y B_\mu / 2) \Phi$$

- 4 new degrees of freedom: doublet+anti-particles
- It has a very specific potential

$$V(\Phi) = m^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2$$

couplings for the EW sector



$$\sqrt{(-m^2/2\lambda)} = v/2 > 0$$

# Consequences - the masses of the bosons

- Two massive charged vector bosons

$$m_W^2 = \frac{g^2 v^2}{4} \quad \text{Corresponding to the observed charged currents}$$

$$\frac{G_F}{\sqrt{2}} = \left( \frac{g}{2\sqrt{2}} \right)^2 \frac{1}{m_W^2} \quad \Rightarrow \quad v = \sqrt{\frac{1}{\sqrt{2}G_F}} \approx 246.22 \text{ GeV}$$

- One massless vector boson


$$m_\gamma = 0$$

- One massive neutral vector boson Z

$$m_Z^2 = (g^2 + g'^2)v^2/4$$

- One massive scalar particle: The Higgs Boson

$$m_H^2 = \frac{4\lambda(v)m_W^2}{g^2}$$




Whose mass is an unknown parameter of the theory as the quartic coupling  $\lambda$

# Higgs Boson couplings in the SM

---

- All the couplings of the Higgs boson to SM particles (except itself) known before the discovery



Is the H responsible for the fermion masses?  
For all the fermion masses?  
Why are the families so different?

# Higgs Boson couplings in the SM

- All the couplings of the Higgs boson to SM particles (except itself) known before the discovery

$\frac{m_f}{v}$ 
 $+ \bar{\Psi}_i y_{ij} \Psi_j \phi + h.c.$

$\frac{2m_V^2}{v}$ 
 $+ |D_\mu \phi|^2$ 
 This term could not exist without a vev

How do we proof there is a condensate?

$v H V^\mu V_\mu$

# Higgs Boson couplings in the SM


- All the couplings of the Higgs boson to SM particles (except itself) known before the discovery

$\frac{m_f}{v}$ 
 $+ \bar{\Psi}_i y_{ij} \Psi_j \phi + h.c.$

$\frac{2m_V^2}{v}$ 
 $+ |D_\mu \phi|^2$ 
 This term could not exist without a vev


$\frac{3m_H^2}{v}$ 
 $\frac{3m_H^2}{v^2}$ 
 $V(\phi)$

What do we know of the real shape of the potential?



# Higgs Couplings - measurements

## Where we are




Measurements of vector bosons and 3rd generation fermions

2nd and 1st generation?

Self couplings?


# (Main) Higgs boson production modes

**Production for  
80fb<sup>-1</sup> in Run2**



**Gluon Fusion**


~4M events produced






# (Main) Higgs boson production modes

**Production for  
80fb<sup>-1</sup> in Run2**




**Gluon Fusion**

~4M events produced




**Vector boson Fusion**

~0.3M events produced




# (Main) Higgs boson production modes

**Production for  
80fb<sup>-1</sup> in Run2**




**Gluon Fusion**

~4M events produced




**Vector boson Fusion**

~0.3M events produced




**W and Z Assoc. Prod.**

~0.2M events produced




# (Main) Higgs boson production modes

## Production for $80\text{fb}^{-1}$ in Run2




### Gluon Fusion

$\sim 4\text{M}$  events produced




### Vector boson Fusion

$\sim 0.3\text{M}$  events produced




### W and Z Assoc. Prod.

$\sim 0.2\text{M}$  events produced




### Top Assoc. Prod.

$\sim 40\text{k}$  events produced




# (Main) Higgs boson production modes

## Production for 80fb<sup>-1</sup> in Run2




### Gluon Fusion

~4M events produced




### Vector boson Fusion

~0.3M events produced




### W and Z Assoc. Prod.

~0.2M events produced



### Top Assoc. Prod.

~40k events produced



**Cross sections increase with center of mass energy.**  
**Note ttH!!**


# Higgs boson decay channels

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

## Third generation quarks

Decay mode	Expected BR for $m_H=125$ GeV
<b>bb</b>	57 %



$$\Gamma_{H \rightarrow bb} \propto \sum_{\text{colors}} \left| \begin{array}{c} \text{H} \\ \text{---} \\ \text{---} \end{array} \begin{array}{l} \nearrow f \\ \searrow \bar{f} \end{array} \right|^2$$


# Higgs boson decay channels

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

## Third generation quarks

Decay mode	Expected BR for $m_H=125$ GeV
<b>bb</b>	57 %



$$\Gamma_{H \rightarrow bb} \propto \sum_{\text{colors}} \left| \begin{array}{c} H \\ \vdots \\ \swarrow \quad \searrow \\ f \quad \bar{f} \end{array} \right|^2 \frac{m_f}{v}$$


# Higgs boson decay channels

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

## Third generation quarks

Decay mode	Expected BR for $m_H = 125$ GeV
<b>bb</b>	57 %



$$\Gamma_{H \rightarrow bb} \propto \sum_{\text{colors}} \left| \begin{array}{c} H \\ \vdots \\ \swarrow \quad \searrow \\ f \quad \bar{f} \end{array} \right|^2 \left( \frac{m_f}{v} \right)^2$$

$$\sim k N_c m_f^2(m_H)(1 + \Delta_{\text{QCD}})$$

$N_c = 3$


# Higgs boson decay channels

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

## Third generation quarks

Decay mode	Expected BR for $m_H = 125$ GeV
<b>bb</b>	57 %



$$\Gamma_{H \rightarrow bb} \propto \sum_{\text{colors}} \left| \begin{array}{c} H \\ \vdots \\ \swarrow \quad \searrow \\ f \quad \bar{f} \end{array} \right|^2 \frac{m_f}{v}$$

$$\sim k N_c \boxed{m_f^2(m_H)} (1 + \Delta_{\text{QCD}})$$

$\uparrow$  3  
 $\uparrow$  2.8 GeV




# Higgs boson decay channels

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

## Third generation quarks

Decay mode	Expected BR for $m_H=125$ GeV
<b>bb</b>	57 %



$$\Gamma_{H \rightarrow bb} \propto \sum_{\text{colors}} \left| \begin{array}{c} H \\ \vdots \\ \swarrow \quad \searrow \\ f \quad \bar{f} \end{array} \right|^2 \frac{m_f}{v}$$

QCD corr.

$$\sim k N_c m_f^2(m_H) (1 + \Delta_{\text{QCD}})$$

$\uparrow$  3                       $\uparrow$  2.8 GeV                       $\uparrow$  0.2


# Higgs boson decay channels

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

## Third generation quarks

Decay mode	Expected BR for $m_H = 125$ GeV
<b>bb</b>	57 %



$$\Gamma_{H \rightarrow bb} \propto \sum_{\text{colors}} \left| \begin{array}{c} H \\ \vdots \\ f \\ \swarrow \\ \bar{f} \end{array} \right|^2 \frac{m_f}{v}$$

QCD corr.

$$\sim k N_c m_f^2(m_H)(1 + \Delta_{\text{QCD}})$$

$8 \times 10^{-5} / \text{GeV}$  (points to  $\sim k$ )  
 $3$  (points to  $N_c$ )  
 $2.8 \text{ GeV}$  (points to  $m_f$ )  
 $0.2$  (points to  $\Delta_{\text{QCD}}$ )

Total width  $\Gamma_{\text{tot}}$ : 4.1 MeV


# Higgs boson decay channels

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

W bosons

Decay mode	Expected BR for $m_H=125$ GeV
bb	57 %
WW	22 %




$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

# Higgs boson decay channels

## Third generation fermions

Decay mode	Expected BR for $m_H=125$ GeV
$bb$	57 %
$WW$	22 %
$\tau\tau$	6.3 %




$$\Gamma_{H \rightarrow \tau\tau} \sim [8 \times 10^{-5} / \text{GeV}] m_f^2$$

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

# Higgs boson decay channels

Z bosons




Decay mode	Expected BR for $m_H=125$ GeV
$bb$	57 %
$WW$	22 %
$\tau\tau$	6.3 %
$ZZ$	3 %

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

# Higgs boson decay channels

## Second generation quarks




Decay mode	Expected BR for $m_H=125$ GeV
$bb$	57 %
$WW$	22 %
$\tau\tau$	6.3 %
$ZZ$	3 %
$cc$	3 %

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

# Higgs boson decay channels

Photons ?? How?





Decay mode	Expected BR for $m_H=125$ GeV
bb	57 %
WW	22 %
$\tau\tau$	6.3 %
ZZ	3 %
cc	3 %
$\gamma\gamma$	0.2 %

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

# Higgs boson decay channels

Photons ?? How?



Decay mode	Expected BR for $m_H=125$ GeV
$bb$	57 %
$WW$	22 %
$\tau\tau$	6.3 %
$ZZ$	3 %
$cc$	3 %
$\gamma\gamma$	0.2 %

Possible through loops  
 What if we have some new particle in the loop?





$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

# Higgs boson decay channels

Z $\gamma$ ?? How?




Decay mode	Expected BR for $m_H=125$ GeV
bb	57 %
WW	22 %
tau tau	6.3 %
ZZ	3 %
cc	3 %
gamma gamma	0.2 %
Z gamma	0.2 %

$$BR_i = \Gamma_i / \Gamma_{\text{tot}}$$

$$\Gamma_{\text{tot}} = \sum \Gamma_i$$

# Higgs boson decay channels

## Second generation fermions



Decay mode	Expected BR for $m_H=125$ GeV
$b\bar{b}$	57 %
$W^+W^-$	22 %
$\tau^+\tau^-$	6.3 %
$Z^+Z^-$	3 %
$c\bar{c}$	3 %
$\gamma\gamma$	0.2 %
$Z\gamma$	0.2 %
$\mu^+\mu^-$	0.02 %

# Higgs boson


- Couplings with the SM particles
- Production modes
- Decay Branching Ratio
  - NOTE: for fermions
  - $\Gamma_{H \rightarrow ff} \sim k N_c m_f^2 (m_H) (1 + \Delta_{\text{QCD}})$

And now.... some fresh air...

# Discovering the Higgs boson at the LHC




# Which are the needed performances for our detector?




- $\gamma\gamma$ : identification and measurement of **photons**
- $ZZ, WW$ : identifications and measurement of **muons, electrons**
- $WW, \tau\tau$ : measurement of **missing transverse energy** (requiring energy measurement up to very forward -  $|\eta| \sim 5$ )
- $bb, \tau\tau$ , efficient and pure **b-tagging** and  **$\tau$  identification**
- **VBF**: Capability to detect **forward jets** (for vector boson fusion processes)

# Concept




# Concept






# Concept



# Concept




# Concept

## Neutrino

No signal

We use the conservation of momentum on the transverse plane to "measure" the transverse missing momentum




# Concept

## Hadrons

(in  $b$ - or  $\tau$ -decays of the Higgs boson)

Hadronic shower  
in the calorimeter

+ Hadronic  
Calorimeter  
tracks




# Concept


## Hadrons (in $b$ - or $\tau$ -decays of the Higgs boson)

Hadronic shower  
in the calorimeter

+ Hadronic  
Calorimeter  
tracks





# ATLAS



# CMS





$$H \rightarrow \gamma\gamma$$



- Low S/B, but relatively high statistics for the signal  
Signal: ~300 events in Run1      Background: Largest contribution from jets
- Very simple selection, quality of detector response and performance play a crucial role
- Main production modes and decay mode occur through loop. Probe for new physics




$$H \rightarrow \gamma\gamma$$





- Low S/B, but relatively high statistics for the signal  
Signal: ~300 events in Run1      Background: Largest contribution from jets
- Very simple selection, quality of detector response and performance play a crucial role
- Main production modes and decay mode occur through loop. Probe for new physics



$$H \rightarrow ZZ^* \rightarrow 4l$$




- High S/B (2-10), but relatively low statistics  
Background:  $pp \rightarrow ZZ$  estimated by MC
- Typically one Z is on-mass shell
- Note:  $BR(H \rightarrow ZZ) \sim 3\%$ ,  $BR(Z \rightarrow ee) + BR(Z \rightarrow \mu\mu) \sim 6.7\%$





$$H \rightarrow ZZ^* \rightarrow 4l$$



- High S/B (2-10), but relatively low statistics  
 Background:  $pp \rightarrow ZZ$  estimated by MC
- Typically one Z is on-mass shell
- Note:  $BR(H \rightarrow ZZ) \sim 3\%$ ,  $BR(Z \rightarrow ee) + BR(Z \rightarrow \mu\mu) \sim 6.7\%$





$$H \rightarrow WW^* \rightarrow 2l2\nu$$





- Large signal event rate, but large background from top and WW
- Requires a very good understanding of the background in simulations and with control regions
- The presence of neutrinos spoil the mass resolution

$$H \rightarrow WW^* \rightarrow 2l2\nu$$



- Large signal event rate, but large background from top and WW
- Requires a very good understanding of the background in simulations and with control regions
- The presence of neutrinos spoil the mass resolution

# A new era - differential measurements



What if New physics exist, but just beyond the reach of the LHC?

We should start having “anomalies”

# Discovering the Higgs boson at the LHC


- Detector needs for the discovery
- Short review of  $H \rightarrow \gamma\gamma$
- Short review of  $H \rightarrow ZZ$
- Short review of  $H \rightarrow WW$
- Measurement of the differential cross sections

# Higgs Properties




# Higgs boson mass

- Higgs mass is the only parameter unconstrained by SM
- Crucial in SM prediction of production and decay modes
- Measurement based on  $H \rightarrow ZZ^* \rightarrow 4l$  and  $H \rightarrow \gamma\gamma$  final states, for which invariant mass can be reconstructed with high precision




[Phys. Rev. D 90, 052004 \(2014\)](#)



[Eur. Phys. J. C 74 \(2014\) 3076](#)


# Measurement of the Higgs boson mass - Systematic impact on the measurements

Largest impact from energy scale - as expected...




# Measurement of the Higgs boson mass

- Mass of Higgs boson measured with  $<0.2\%$  precision
- $M_H = 125.09 \pm 0.24 \text{ GeV} [ \pm 0.21 \text{ (stat.) } \pm 0.11 \text{ (syst.) } ]$
- Dominant systematics: energy or momentum scale and resolution for  $\gamma, e, \mu$



# Measurement of the Higgs boson mass

- Mass of  $H$
- $m_H = 125.09 \pm 0.24$  GeV
- Dominant



for  $\gamma, e, \mu$

ATLAS  
LHC F  
ATLAS H  
CMS H  
ATLAS H  
CMS H  
ATLAS+  
ATLAS+  
ATLAS+



) GeV

$m_H$  [GeV]

# Knowing the mass....

- SM predictions for production mode cross sections and decay BR fully determined

[CERN-2013-004, FERMILAB-CONF-13-667-T](#)



- Combining measurements and searches by ATLAS and CMS collaborations published in 17 individual publications in RUN1

# Analysis - Measurements in ATLAS and CMS

- Integrated luminosities per experiment:  **$\sim 5 \text{ fb}^{-1}$  at  $\sqrt{s} = 7 \text{ TeV}$   $\sim 20 \text{ fb}^{-1}$  at  $\sqrt{s} = 8 \text{ TeV}$**

- **Why combining?**

Doubling statistical power in measuring the Higgs boson production and decay rates

**Rule of thumbs: improving by  $1/\sqrt{2}$  the precision**

Channel	References for individual publications		Signal strength [ $\mu$ ] from results in this paper (Section 5.2)		Signal significance [ $\sigma$ ]	
	ATLAS	CMS	ATLAS	CMS	ATLAS	CMS
$H \rightarrow \gamma\gamma$	[91]	[92]	1.14 <sup>+0.27</sup> <sub>-0.25</sub> (+0.26) (-0.24)	1.11 <sup>+0.25</sup> <sub>-0.23</sub> (+0.23) (-0.21)	5.0 (4.6)	5.6 (5.1)
$H \rightarrow ZZ$	[93]	[94]	1.52 <sup>+0.40</sup> <sub>-0.34</sub> (+0.32) (-0.27)	1.04 <sup>+0.32</sup> <sub>-0.26</sub> (+0.30) (-0.25)	7.6 (5.6)	7.0 (6.8)
$H \rightarrow WW$	[95,96]	[97]	1.22 <sup>+0.23</sup> <sub>-0.21</sub> (+0.21) (-0.20)	0.90 <sup>+0.23</sup> <sub>-0.21</sub> (+0.23) (-0.20)	6.8 (5.8)	4.8 (5.6)
$H \rightarrow \tau\tau$	[98]	[99]	1.41 <sup>+0.40</sup> <sub>-0.36</sub> (+0.37) (-0.33)	0.88 <sup>+0.30</sup> <sub>-0.28</sub> (+0.31) (-0.29)	4.4 (3.3)	3.4 (3.7)
$H \rightarrow bb$	[100]	[101]	0.62 <sup>+0.37</sup> <sub>-0.37</sub> (+0.39) (-0.37)	0.81 <sup>+0.45</sup> <sub>-0.43</sub> (+0.45) (-0.43)	1.7 (2.7)	2.0 (2.5)
$H \rightarrow \mu\mu$	[102]	[103]	-0.6 <sup>+3.6</sup> <sub>-3.6</sub> (+3.6) (-3.6)	0.9 <sup>+3.6</sup> <sub>-3.5</sub> (+3.3) (-3.2)		
$ttH$ production	[77, 104, 105]	[107]	1.9 <sup>+0.8</sup> <sub>-0.7</sub> (+0.7) (-0.7)	2.9 <sup>+1.0</sup> <sub>-0.9</sub> (+0.9) (-0.8)	2.7 (1.6)	3.6 (1.3)

off-shell analyses not in combination

	Untagged	VBF	VH	ttH
$H \rightarrow \gamma\gamma$	✓	✓	✓	✓
$H \rightarrow ZZ \rightarrow 4l$	✓	✓	✓	✓
$H \rightarrow WW \rightarrow 2l2\nu$	✓	✓	✓	✓
$H \rightarrow \tau\tau$	✓	✓	✓	✓
$H \rightarrow bb$			✓	✓
$H \rightarrow \mu\mu$	✓	✓		
$H \rightarrow Z\gamma$				
$H \rightarrow \text{inv}$				

in ATLAS combination

in CMS combination



overwhelming multijet BKG  
not yet in combination  
extremely low  $\sigma_i \times B_{\mu\mu}$

# From single channel, to combined results

- To **enhance the sensitivity**, the experimental analysis uses **event categories(k)** also based on **multi variate techniques**
- Sensitivity to different production modes**

$$n_{\text{signal}}(k) = \mathcal{L}(k) \cdot \sum_i \sum_f \left\{ \sigma_i \cdot A_i^{f,SM}(k) \cdot \varepsilon_i^f(k) \cdot B^f \right\}$$

Inclusive SM cross-section for production mode  $i$  i.e. gluon-gluon fusion





# From single channel, to combined results

- To **enhance the sensitivity**, the experimental analysis uses **event categories(k)** also based on **multi variate techniques**
- Sensitivity to different production modes**

$$n_{\text{signal}}(k) = \mathcal{L}(k) \cdot \sum_i \sum_f \left\{ \sigma_i \cdot A_i^{f,SM}(k) \cdot \varepsilon_i^f(k) \cdot \boxed{\mathbf{B}^f} \right\}$$

Branching Fractions  
i.e.:  $H \rightarrow ZZ$





# From single channel, to combined results

- To **enhance the sensitivity**, the experimental analysis uses **event categories(k) also based on multi variate techniques**
- Sensitivity to different production modes**

$$n_{\text{signal}}(k) = \mathcal{L}(k) \cdot \sum_i \sum_f \left\{ \sigma_i \cdot \boxed{A_i^{f,SM}(k) \cdot \varepsilon_i^f(k)} \cdot B^f \right\}$$

Acceptances and efficiencies, from MC assuming SM


Production process	Event generator	
	ATLAS	CMS
<i>ggF</i>	POWHEG [79–83]	POWHEG
VBF	POWHEG	POWHEG
<i>WH</i>	PYTHIA8 [84]	PYTHIA6.4 [85]
<i>ZH</i> ( $qq \rightarrow ZH$ or $qg \rightarrow ZH$ )	PYTHIA8	PYTHIA6.4
<i>ggZH</i> ( $gg \rightarrow ZH$ )	POWHEG	See text
<i>ttH</i>	POWHEL [87]	PYTHIA6.4
<i>tHq</i> ( $qb \rightarrow tHq$ )	MADGRAPH [89]	AMC@NLO [78]
<i>tHW</i> ( $gb \rightarrow tHW$ )	AMC@NLO	AMC@NLO
<i>bbH</i>	PYTHIA8	PYTHIA6.4, AMC@NLO

# From single channel, to combined results

- To **enhance the sensitivity**, the experimental analysis uses **event categories(k)** also based on **multi variate techniques**
- Sensitivity to different production modes**

$$n_{\text{signal}}(k) = \mathcal{L}(k) \cdot \sum_i \sum_f \left\{ \sigma_i \cdot A_i^{f,SM}(k) \cdot \varepsilon_i^f(k) \cdot B^f \right\}$$

Example: ttH, H → multilepton



Category	Higgs boson decay mode			
	WW*	ττ	ZZ*	Other
2ℓ0τ <sub>had</sub>	80%	15%	3%	2%
3ℓ	74%	15%	7%	4%
2ℓ1τ <sub>had</sub>	35%	62%	2%	1%
4ℓ	69%	14%	14%	4%
1ℓ2τ <sub>had</sub>	4%	93%	0%	3%

# From single channel, to combined results

---

- To **enhance the sensitivity**, the experimental analysis uses **event categories(k) also based on multi variate techniques**
- **Sensitivity to different production modes**

$$n_{\text{signal}}(k) = \mathcal{L}(k) \cdot \sum_i \sum_f \left\{ \sigma_i \cdot A_i^{f,SM}(k) \cdot \varepsilon_i^f(k) \cdot B^f \right\}$$

Full combination: **~600 signal regions & control regions**

**Grand total of ~4200 nuisance parameters:**

related to (systematic) uncertainties

**Correlation scheme:** strategy of nuisance parameters a delicate and complicated task (would deserve a separate talk)

# From single channel, to combined results



---

- To **enhance the sensitivity**, the experimental analysis uses **event categories(k) also based on multi variate techniques**
- **Sensitivity to different production modes**

$$n_{\text{signal}}(k) = \mathcal{L}(k) \cdot \sum_i \sum_f \left\{ \sigma_i \cdot A_i^{f,SM}(k) \cdot \varepsilon_i^f(k) \cdot B^f \right\}$$

- **What to measure?**
- **To reduce as much as possible the assumptions on the SM nature of the Higgs boson, we can measure  $\sigma_i B^f$ .  
SM assumption only on  $A$   $\varepsilon$  and  $\sigma_i(7\text{TeV})/\sigma_i(8\text{TeV})$**

# Cross Sections times Branching Ratios



• As expected, correlations due to signal **mix** of **production modes** in the analysis categories:

- **ggF VS VBF** (in 2-jet selections) or **WH VS ZH** ( $V \rightarrow \text{hadrons}$ ) in  $H \rightarrow \gamma\gamma$ ;

and **decay modes**:

- **ττ VS WW** in **ttH** (in multileptons)

$$\sigma_i \cdot \mathbf{B}^f = \frac{\sigma_i(\vec{k}) \cdot \Gamma^f(\vec{k})}{\Gamma_H}$$


$$\kappa_j^2 = \sigma_j / \sigma_j^{\text{SM}} \quad \text{or} \quad \kappa_j^2 = \Gamma^j / \Gamma_{\text{SM}}^j$$

# Coupling modifiers

- $\sigma^{\text{SM}}_j$  and  $\Gamma_{\text{SM}}^j$  are calculated using the status of art theoretical SM predictions
- Higgs vertexes scales by a factor  $k$ 
  - Recover the SM if  $k=1$

## Example: $ggF \rightarrow H \rightarrow WW$ (or $ZZ$ )

Effective  $\kappa$



$$\sigma_{ggF} = \kappa_g^2 \sigma_{ggF}(\text{SM})$$

$$\Gamma_{W,Z} = \kappa_{W,Z}^2 \Gamma_{W,Z}(\text{SM})$$

- and  $\Gamma_H$ ?

- Option 1: assume only SM decay modes


$$\Gamma_H = \sum_j \mathbf{B}_{\text{SM}}^j \kappa_j^2 \cdot \Gamma_H^{\text{SM}}$$

- Option 2: allow for an additional branching fractions in BSM

$$\Gamma_H = \frac{\sum_j \mathbf{B}_{\text{SM}}^j \kappa_j^2 \cdot \Gamma_H^{\text{SM}}}{1 - \mathbf{B}_{\text{BSM}}} \quad 107$$

Resolving the loop, assuming only SM

or





$$\sigma_{ggF} = (1.06 \kappa_t^2 + 0.01 \kappa_b^2 - 0.07 \kappa_b \kappa_t) \sigma_{ggF}(\text{SM})$$

$$\Gamma_{W,Z} = \kappa_{W,Z}^2 \Gamma_{W,Z}(\text{SM})$$

# Resolving the loops and assuming coupling with only SM particles

- **Interferences** help to resolve the sign (*NB*:  $\kappa_\tau$  and  $\kappa_\mu$ )
- *NB*: in this fit model, low measured value of  **$\kappa_b$**  reduces total width  $\Gamma_H \Rightarrow$  all  $\kappa_i$  measured low

Production	Loops	Interference	Resolved scaling factor
$\sigma(ggF)$	✓	$t-b$	$1.06 \cdot \kappa_t^2 + 0.01 \cdot \kappa_b^2 - 0.07 \cdot \kappa_t \kappa_b$
$\sigma(\text{VBF})$	-	-	$0.74 \cdot \kappa_W^2 + 0.26 \cdot \kappa_Z^2$
$\sigma(\text{WH})$	-	-	$\kappa_W^2$
$\sigma(qq/qg \rightarrow ZH)$	-	-	$\kappa_Z^2$
$\sigma(gg \rightarrow ZH)$	✓	$t-Z$	$2.27 \cdot \kappa_Z^2 + 0.37 \cdot \kappa_t^2 - 1.64 \cdot \kappa_Z \kappa_t$
$\sigma(ttH)$	-	-	$\kappa_t^2$
$\sigma(gb \rightarrow tHW)$	-	$t-W$	$1.84 \cdot \kappa_t^2 + 1.57 \cdot \kappa_W^2 - 2.41 \cdot \kappa_t \kappa_W$
$\sigma(qq/qb \rightarrow tHq)$	-	$t-W$	$3.40 \cdot \kappa_t^2 + 3.56 \cdot \kappa_W^2 - 5.96 \cdot \kappa_t \kappa_W$
$\sigma(bbH)$	-	-	$\kappa_b^2$
Partial decay width			
$\Gamma^{ZZ}$	-	-	$\kappa_Z^2$
$\Gamma^{WW}$	-	-	$\kappa_W^2$
$\Gamma^{\gamma\gamma}$	✓	$t-W$	$1.59 \cdot \kappa_W^2 + 0.07 \cdot \kappa_t^2 - 0.66 \cdot \kappa_W \kappa_t$
$\Gamma^{\tau\tau}$	-	-	$\kappa_\tau^2$
$\Gamma^{bb}$	-	-	$\kappa_b^2$
$\Gamma^{\mu\mu}$	-	-	$\kappa_\mu^2$
Total width ( $B_{\text{BSM}} = 0$ )			
$\Gamma_H$	✓	-	$0.57 \cdot \kappa_b^2 + 0.22 \cdot \kappa_W^2 + 0.09 \cdot \kappa_g^2 + 0.06 \cdot \kappa_\tau^2 + 0.03 \cdot \kappa_Z^2 + 0.03 \cdot \kappa_c^2 + 0.0023 \cdot \kappa_\gamma^2 + 0.0016 \cdot \kappa_{(Z\gamma)}^2 + 0.0001 \cdot \kappa_s^2 + 0.00022 \cdot \kappa_\mu^2$



# Fermions and bosons

- Testing the intrinsic difference between couplings to

- W/Z: EW Symmetry Breaking

$$\kappa_Z = \kappa_W = \kappa_V$$


- fermions: Yukawa couplings

$$\kappa_t = \kappa_\tau = \kappa_b = \kappa_F$$

- Sensitivity to the relative sign between  $\kappa_V$  and  $\kappa_F$  through interference terms

- Large asymmetry between the positive and negative coupling ratios for  $H \rightarrow \gamma\gamma$

$$\Gamma^{\gamma\gamma}_{t-W} = 1.59 \cdot \kappa_W^2 + 0.07 \cdot \kappa_t^2 - 0.66 \cdot \kappa_W \kappa_t$$



- Fits on individual channels have slight preference for negative  $\kappa_F$
- Combined result converges to positive  $\kappa_F$



# Higgs Properties

- Higgs Mass
- Higgs production XS times BR
- Higgs couplings

## Two quarks for Muster Higgs yesterday!!!

Since the big discovery of 2012, the Large Hadron Collider at CERN has been accumulating data and making steady progress. Two recent results establish the origins of the mass of the two heaviest quarks





Selected news from Run2

▲ The visitor centre at the ALICE experiment on the CERN Large Hadron Collider Photograph: Jon Butterworth

Observed!





# The Big news (1): $\tau$ -Higgs interaction




## VBF

With two forward jets and a large rapidity gap between the jets



Analysis based on several channels depending on the decay mode of the  $\tau$ .



Background is Z production with two jets


# The Big news (2): Top -Higgs interaction

---


$$t \ t \ H$$

# The Big news (2): Top -Higgs interaction



---





# The Big news (2): Top -Higgs interaction



# The Big news (2): Top -Higgs interaction




# The Big news (2): Top -Higgs interaction







# The Big news (2): Top -Higgs interaction



Currently cleanest and most sensitive channel






# The Big news (2): Top -Higgs interaction




$ttH(bb)$




Very large backgrounds of top pair production associated with b jets














# The Big news (2): Top -Higgs interaction




# The Big news (2): Top -Higgs interaction



# The Big news (2): Top -Higgs interaction



# The Big news (2): Top -Higgs interaction





## Multi lepton channel

Large number of topologies  
intricate reducible backgrounds of jets faking leptons












The diagrams illustrate various event topologies in a multi-lepton channel. Each diagram shows a central vertex with multiple outgoing particles represented by colored cones. Labels include:

- top-quark** (red dashed circles)
- b-jet** (pink cones)
- light-jet** (blue cones)
- Higgs (bb)** (pink cones)
- lepton** (red lines)
- neutrino** (red lines)
- tau** (green cones)
- photon** (yellow cones)




# The Big news (2): Top -Higgs interaction

*it seems a bit complex...*



Observed!

# The Big news (2): Top -Higgs interaction





- CMS and ATLAS results presented at LHCP2018
- Both collaborations had to combine channels to obtain the sensitivity necessary to claim an observation
- All possible advanced tools were used
  - Multi Variate Analysis
  - Matrix Element Method
  - Status of the art theory predictions)



Observed!


# The Big news (2): Top -Higgs interaction

- Why important:
  - We have a proof that top and H interact...
  - ...and they interact strongly
  - Is this because of some new dynamics?
  - Is this strong coupling indicating something more than the SM?
  - We have an handle to know what happens in the gluon-gluon fusion loop.



# The Big news (3): Bottom -Higgs interaction


- ATLAS presented the  $H \rightarrow bb$  observation in ICHEP 2018
- In addition to probing coupling to b-quarks:
  - $H \rightarrow bb$  drives the uncertainty on the total decay width, and thus on measurement of absolute couplings
  - It also drives the indirect limit on “undetected/ invisible” decays




# The Big news (3): Bottom -Higgs interaction

G. Piacquadio - ICHEP 2018


0-lepton




1-lepton



2-lepton



- VH production most sensitive mode for  $H \rightarrow bb$  at the LHC
- 3 channels (0-, 1-, 2 charged leptons from  $V=W/Z$  boson)
- Select 2 b-tagged jets and  $p_T(V) > 75$  or 150 GeV
- Main discriminant variables  $m(bb)$ ,  $p_T(V)$  and  $\Delta R(bb)$  (combined into a Boosted Decision Tree)




Non-resonant backgrounds:  
 $t\bar{t}$ ,  
 single top  
 (NLO, PowHeg)

W+jets  
 Z+jets  
 (NLO for up to 2 extra jets, Sherpa 2.2.1)

Overall strategy:  
 normalization from data, shapes from MC

# The Big news (3): Bottom -Higgs interaction

G. Piacquadio - ICHEP 2018



Fit result with 79.8 fb<sup>-1</sup> of Run-2 data

$$\mu = \sigma_{\text{meas}} / \sigma_{\text{SM}} = 1.16^{+0.27}_{-0.25}$$

Significance: **4.9 $\sigma$**  (4.3 $\sigma$  expected)


Combination with Run-1:

$$\mu = 0.98 \pm 0.14(\text{stat.})^{+0.17}_{-0.16}(\text{syst.})$$

Significance: **4.9 $\sigma$**  (5.1 $\sigma$  expected)


● Detailed validation of analysis:

- Fit to diboson VZ, Z  $\rightarrow$  bb:  $\mu = 1.20^{+0.20}_{-0.18}$  (**9.6 $\sigma$** )
- m(bb) fit for VH, H  $\rightarrow$  bb:  $\mu = 1.06^{+0.36}_{-0.33}$  (**3.6 $\sigma$** )



# Yesterday in Higgs Hunting 2018 in Paris

## First comprehensive presentation of the analysis!



I. Luise  
Ph.D. Student


Observed!

# The Big news (3): Bottom -Higgs interaction AND VH production mode!!!

## H → bb combination

NEW

- Run-1+Run-2
  - VH, H → bb
  - VBF(+ggF), H → bb
  - ttH, H → bb



Significance:  
**5.4σ** observed  
(5.5σ expected)

**Observation of  
H → bb!!**

## VH combination

NEW

- Run-2
  - VH, H → bb
  - VH, H → γγ
  - VH, H → ZZ\*

Significance:  
**5.3σ** observed (4.8σ expected)

**Observation of  
VH production!!**



# Where we are today

---



## Production

## Decays


gluon fusion






vector boson fusion (VBF)




associated prod. with W/Z




associated prod. with tt



 = observed



# ... more rare options....

## Invisible decays





$Br_{inv} < 0.24$  (0.23)  
at 95% CL

## di-muons




Limits currently ~2 x SM

## Z-photon




Limits currently ~6 x SM


## Quarkonia-photon



Higgs ~400 x SM




Higgs ~200 x SM





Higgs ~50 x SM



# ...and the big challenge: Double Higgs production!




- The nature of the Higgs potential is one of the big open questions in EW symmetry breaking. In SM potential is determined by  $G_F$  and  $m_H$
- Direct measurement of Higgs self coupling is the big challenge in the EWSB.






- Destructive interference  $\rightarrow \sigma_{hh} \approx \frac{\sigma_T + \sigma_B}{2.5} \rightarrow \sigma_{hh} = 39.5 \text{ fb @ 14 TeV}$

- Deviations from SM because of BSM?
- Resonances?
- Which channels? which machines?



# ...and the big challenge: Double Higgs production!

- The nature of the Higgs potential is one of the big open question in EW symmetry breaking. In SM potential is determined by  $G_F$  and  $m_H$



**ATLAS  $bb\tau\tau$   $\sigma_{HH} \rightarrow bb\tau\tau < 13 \times SM$  (15 exp.)**

**CMS  $bbbb$  and full combination!  $\sigma_{HH} \text{ comb} < 22 \times SM$  (13 exp.)**

$$-11.8 \lambda_{SM} < \lambda < 18.8 \lambda_{SM} @ 95\% \text{ CL}$$



## Selected news from Run2

- Observation of  $H \rightarrow \tau\tau$
- Observation of  $pp \rightarrow ttH$
- Observation of  $H \rightarrow bb$
- Observation of  $pp \rightarrow VH$
- Beginning of the quest for  $pp \rightarrow HH$  and constrains on  $\lambda$

BONUS

Some important questions...

# Some more considerations: back to the potential, and the fate of the Universe



- What if the the EW minimum is a local and not the absolute minimum of the potential?
- Couplings use to run.... What about the self coupling  $\lambda$ ?
- It depends on the mass of the Higgs boson and of the Top quarks
- If it at some scale,  $\lambda$  changes sign, the vacuum is not stable... Maybe we are safe if some new dynamics enters. For the time being, the only new dynamics we know is related to gravity, and it has a scale of the order of the Plank mass  $10^{19}$  GeV.

# Some more considerations: back to the potential, and the fate of the Universe

*slide from  
R. Goncalo*


The universe seems to live  
near a critical condition

JHEP 1208 (2012) 098

Why?!

Explained by underlying theory?

Anthropic principle?




# Is the Standard Model complete?

---

- mmm... a lot of open question...
- Why do we observe matter and almost no antimatter, if we believe there is a symmetry between the two in the universe?
- What is the “dark matter” that we can’t see, but it has visible gravitational effect in the cosmos?
- Are quarks and leptons actually fundamental, or made up of more fundamental particles?
- Why are there exactly three generations of quarks and leptons?
- What is the explanation for the observed pattern for the particle masses?

no hint of new physics at the LHC yet...



# *Where is everybody?*








# Where is everybody?

Option 1: **New physics at TeV exist,**  
we (you) will discover it soon!



# Where is everybody?

Option 1: **New physics at TeV exist,**  
we (you) will discover it soon!





Option 2: New physics exist, **but just beyond the reach of the LHC.**  
But we should start having “**anomalies**”

# Where is everybody?

Option 1: **New physics at TeV exist**, we (you) will discover it soon!

Option 3: **No new physics at the TeV** scale we need to understand better the questions we are asking



Option 2: New physics exist, **but just beyond the reach of the LHC.**  
But we should start having “**anomalies**”

# An example: Models with two Higgs doublets


- Why just one doublet for the Higgs Sector?
- It is the more economical, but no limitation to the presence of other scalar doublets.
- Example: hMSSM  
(some configuration of SUSY models, with two Higgs doublets)

- This leads to 5 different Higgs Bosons

- CP even (scalar):  $h, H$
- CP odd (pseudoscalar):  $A$
- Charged :  $H^+ H^-$

- We can search these new particles...

- ...or we can check if they leave some anomaly in the couplings of the boson we observed...




Back to the future

# The reach of the LHC

---

*Energy*




*R&D  
new ideas*


*Data*  
NOTE: Higgs Discovery  
0.2M bosons H produced

# The reach of the LHC

---




# The reach of the LHC






# The reach of the LHC




# Future colliders




# Future colliders

HL-LHC = same energy, but collision rate 2÷3 time higher



# Future colliders

HE-LHC = 2 times the center of mass energy?



# Future colliders

FCC-ee ( ? ) **90–350 GeV**

→ FCC-hh ( ? ) **100 TeV**

**100 km**

CepC (2028 ?) **240 GeV**

→ SppC (2042 ?) **70 TeV**

**54 km**

**27 km**

HL-LHC (2026–2035)

**14 TeV**

**33 TeV**

**0.3 km**

CLIC ( ? ) **380 GeV – 3 TeV**

**48 km**

muon collider ( ? ) **125 GeV**

**11 km**

**31 km**

**50 km**

ILC ( ? ) **250–500 GeV**

# Ideas for new Future Colliders

		Begin -end data taking		L[ab-1] @ $\sqrt{s}$ [GeV]				L[ab-1] @ $\sqrt{s}$ [TeV]						Total L[ab-1] at $\sqrt{s}>240$ GeV	#years	#H events
				90	~240	350-380	500	1.5	3	7-14	27	70	100			
LHC	pp	2010	2023							0.3				0.3	13	<b>15M</b>
HL-LHC	pp	2026	2035							3				3	~10	<b>150M</b>
CepC	ee	2028?	2038?		5									5	~10	<b>1M</b>
ILC	ee	2030?	2050?		2	0.2	4							6.2	~20	<b>1.6M</b>
CLIC	ee	2035?	2055?			0.5		1.5	3					4	~20	<b>1.5-2M</b>
FCC-ee	ee	2039?	2055?	150	5	2								13	~15	<b>1-2M</b>
HE-LHC	pp	2040?	2060?								>10			>10	~20	<b>1B</b>
FCC-hh	pp	2043? (FCC-ee?)	2063?										40	40	~25	<b>40B</b>
SppC	pp	2045?	2060?									30		30	~10-15	<b>30B</b>

LHC→HL-LHC: 10 times more H (50-100x analysed data)

HL-LHC→HE-LHC: ~10 times more H (500-1000x analysed data)

HE-LHC→FCC-hh: ~40 times more H (20000-40000x analysed data)

Backup

Going beyond the SM



# BSM: Direct Searches


---

Direct searches of new physics  
gave non positive results so far  
Some examples

# BSM: Direct Searches

Direct searches of new physics gave non positive results so far  
Some examples

**Supersymmetry  
(MSSM)**



\*Observed limits at 95% C.L. - theory uncertainties not included  
Only a selection of available mass limits. Probe \*up to\* the quoted mass limit for  $m_{\text{LSP}} \approx 0$  GeV unless stated otherwise

# BSM: Direct Searches

Direct searches of new physics gave non positive results so far  
Some examples

**Supersymmetry (MSSM)**




**Top-partners**  
**Composite Higgs**  
**Extra-dimensions**  
**Excited Quark**

...



Selected CMS SUSY Results\* - SMS Interpretation

ICHEP '16 - Moriond '17



ATLAS Exotics Searches\* - 95% CL Upper Exclusion Limits

Status: July 2017

ATLAS Preliminary  
 $\int \mathcal{L} dt = (3.2 - 37.0)\text{ fb}^{-1}$   
 $\sqrt{s} = 8, 13\text{ TeV}$

Model	$\ell, \gamma$	Jets <sup>†</sup>	Emiss <sup>†</sup>	$[\mathcal{L} dt][\text{fb}^{-1}]$	Limit	Reference	
Extra dimensions	ADD $G_{KK} + g/q$	$0, e, \mu$	$1-4j$	Yes	36.1	$M_{\text{Pl}}$ 7.75 TeV	$n=2$ ATLAS-CONF-2017-060
	ADD non-resonant $\gamma\gamma$	$2\gamma$	-	-	36.7	$M_{\text{Pl}}$ 8.6 TeV	$n=3$ HLZ NLO CERN-EP-2017-132
	ADD QBH	-	$2j$	-	37.0	$M_{\text{Pl}}$ 8.9 TeV	$n=6$ 1703.09217
	ADD BH high $\sum p_T$	$\geq 1, e, \mu$	$\geq 2j$	-	3.2	$M_{\text{Pl}}$ 8.2 TeV	$n=6, M_D = 3\text{ TeV, rot BH}$ 1608.02265
	ADD BH multijet	-	$\geq 3j$	-	3.6	$M_{\text{Pl}}$ 9.55 TeV	$n=6, M_D = 3\text{ TeV, rot BH}$ 1512.02586
RS1 $G_{KK} \rightarrow \gamma\gamma$	$2\gamma$	-	-	36.7	$G_{KK}$ mass 4.1 TeV	$k/M_{\text{Pl}} = 0.1$ CERN-EP-2017-132	
Bulk RS $G_{KK} \rightarrow WW \rightarrow qq\bar{q}\bar{q}$	$1, e, \mu$	$1, j$	-	36.1	$G_{KK}$ mass 1.75 TeV	$k/M_{\text{Pl}} = 1.0$ ATLAS-CONF-2017-051	
ZUED / RPP	$1, e, \mu$	$\geq 2b, \geq 3j$	Yes	13.2	$KK$ mass 1.6 TeV	Tier (1,1), $\mathcal{B}(A^{(1,1)} \rightarrow t\bar{t}) = 1$ ATLAS-CONF-2016-104	
Gauge bosons	SSM $Z' \rightarrow \ell\ell$	$2, e, \mu$	-	-	36.1	$Z'$ mass 4.5 TeV	ATLAS-CONF-2017-027
	SSM $Z' \rightarrow \tau\tau$	$2, \tau$	-	-	36.1	$Z'$ mass 2.4 TeV	ATLAS-CONF-2017-050
	Leptophobic $Z' \rightarrow b\bar{b}$	-	$2b$	-	3.2	$Z'$ mass 1.5 TeV	1603.08791
	Leptophobic $Z' \rightarrow t\bar{t}$	$1, e, \mu$	$\geq 1b, \geq 1J/2j$	Yes	3.2	$Z'$ mass 2.0 TeV	$\Gamma/m = 3\%$ ATLAS-CONF-2016-014
	SSM $W' \rightarrow \ell\nu$	$1, e, \mu$	-	Yes	36.1	$W'$ mass 5.1 TeV	1706.04788
HVT $V' \rightarrow WW \rightarrow qq\bar{q}\bar{q}$ model B	$0, e, \mu$	$2, j$	-	36.7	$V'$ mass 3.5 TeV	CERN-EP-2017-147	
HVT $V' \rightarrow WH/ZH$ model B	multi-channel	-	-	36.1	$V'$ mass 2.93 TeV	ATLAS-CONF-2017-055	
LRSM $W'_\mu \rightarrow t\bar{b}$	$1, e, \mu$	$2b, 0-1j$	Yes	20.3	$W'$ mass 1.92 TeV	1410.4103	
LRSM $W'_\mu \rightarrow t\bar{b}$	$0, e, \mu$	$\geq 1b, 1, j$	-	20.3	$W'$ mass 1.76 TeV	1408.0886	
CI	CI $qqqq$	-	$2j$	-	37.0	$A$ 21.8 TeV $\eta_{LL}$	1703.09217
	CI $\ell\ell qq$	$2, e, \mu$	-	-	36.1	$A$ 40.1 TeV $\eta_{LL}$	ATLAS-CONF-2017-027
	CI $uu\tau\tau$	$2(SS) \geq 3, e, \mu$	$\geq 1b, \geq 1j$	Yes	20.3	$A$ 4.9 TeV	$ C_{\text{SM}}  = 1$ 1504.04605
DM	Axial-vector mediator (Dirac DM)	$0, e, \mu$	$1-4j$	Yes	36.1	$m_{\text{DM}}$ 1.5 TeV	$g_p = 0.25, g_s = 1.0, m(\chi) < 400\text{ GeV}$ ATLAS-CONF-2017-060
	Vector mediator (Dirac DM)	$0, e, \mu, 1\gamma$	$\leq 1j$	Yes	36.1	$m_{\text{DM}}$ 1.2 TeV	$g_p = 0.25, g_s = 1.0, m(\chi) < 480\text{ GeV}$ 1704.03848
	$VV_{JK}$ EFT (Dirac DM)	$0, e, \mu$	$1, j, \leq 1j$	Yes	3.2	$M_c$ 700 GeV	$m(\chi) < 150\text{ GeV}$ 1608.02372
LO	Scalar LQ 1 <sup>st</sup> gen	$2, e$	$\geq 2j$	-	3.2	LQ mass 1.1 TeV	$\beta = 1$ 1605.06035
	Scalar LQ 2 <sup>nd</sup> gen	$2, \mu$	$\geq 2j$	-	3.2	LQ mass 1.05 TeV	$\beta = 1$ 1605.06035
	Scalar LQ 3 <sup>rd</sup> gen	$1, e, \mu$	$\geq 1b, \geq 3j$	Yes	20.3	LQ mass 640 GeV	$\beta = 0$ 1508.04735
Heavy quarks	VLO $TT \rightarrow Ht + X$	$0$ or $1, e, \mu$	$\geq 2b, \geq 3j$	Yes	13.2	$T$ mass 1.2 TeV	$\mathcal{B}(T \rightarrow Ht) = 1$ ATLAS-CONF-2016-104
	VLO $TT \rightarrow Zt + X$	$1, e, \mu$	$\geq 1b, \geq 3j$	Yes	36.1	$T$ mass 1.16 TeV	$\mathcal{B}(T \rightarrow Zt) = 1$ 1705.10751
	VLO $TT \rightarrow Wb + X$	$1, e, \mu$	$\geq 1b, \geq 1J/2j$	Yes	36.1	$T$ mass 1.35 TeV	$\mathcal{B}(T \rightarrow Wb) = 1$ CERN-EP-2017-094
	VLO $BB \rightarrow Hb + X$	$1, e, \mu$	$\geq 2b, \geq 3j$	Yes	20.3	$B$ mass 700 GeV	$\mathcal{B}(B \rightarrow Hb) = 1$ 1505.04306
	VLO $BB \rightarrow Zb + X$	$2j \geq 3, e, \mu$	$\geq 2j \geq 1b$	-	20.3	$B$ mass 790 GeV	$\mathcal{B}(B \rightarrow Zb) = 1$ 1409.5500
	VLO $BB \rightarrow Wt + X$	$1, e, \mu$	$\geq 1b, \geq 1J/2j$	Yes	36.1	$B$ mass 1.25 TeV	$\mathcal{B}(B \rightarrow Wt) = 1$ CERN-EP-2017-094
Excited fermions	VLO $QQ \rightarrow WqWq$	$1, e, \mu$	$\geq 4j$	Yes	20.3	$Q$ mass 690 GeV	1509.04261
	Excited quark $q^* \rightarrow qg$	-	$2j$	-	37.0	$q^*$ mass 6.0 TeV	only $u'$ and $d'$ , $\Lambda = m(q')$ 1703.09127
	Excited quark $q^* \rightarrow q\gamma$	$1\gamma$	$1j$	-	36.7	$q^*$ mass 5.3 TeV	only $u'$ and $d'$ , $\Lambda = m(q')$ CERN-EP-2017-148
	Excited quark $b^* \rightarrow bg$	-	$1b, 1j$	-	13.3	$b^*$ mass 2.3 TeV	ATLAS-CONF-2016-060
	Excited quark $b^* \rightarrow Wt$	$1$ or $2, e, \mu$	$1b, 2-0j$	Yes	20.3	$b^*$ mass 1.5 TeV	$f_u = f_b = f_s = 1$ 1510.02664
Other	Excited lepton $\ell^*$	$3, e, \mu, \tau$	-	-	20.3	$\ell^*$ mass 3.0 TeV	$\Lambda = 3.0\text{ TeV}$ 1411.2921
	Excited lepton $\nu^*$	$3, e, \mu, \tau$	-	-	20.3	$\nu^*$ mass 1.6 TeV	$\Lambda = 1.6\text{ TeV}$ 1411.2921
	LRSM Majorana $\nu$	$2, e, \mu$	$2j$	-	20.3	$\nu^c$ mass 2.0 TeV	$m(W_\mu) = 2.4\text{ TeV, no mixing}$ 1506.06020
	Higgs triplet $H^{\pm\pm} \rightarrow \ell\ell$	$2, 3, 4, e, \mu$ (SS)	-	-	36.1	$H^{\pm\pm}$ mass 870 GeV	DY production, $\mathcal{B}(H^{\pm\pm} \rightarrow \ell\tau) = 1$ ATLAS-CONF-2017-053
	Higgs triplet $H^{\pm\pm} \rightarrow \ell\tau$	$3, e, \mu, \tau$	-	-	20.3	$H^{\pm\pm}$ mass 400 GeV	1411.2921
Monotop (non-res prod)	$1, e, \mu$	$1b$	Yes	20.3	spin-1 invisible particle mass 637 GeV	$A_{\text{non-res}} = 0.2$ 1410.5404	
Multi-charged particles	-	-	-	20.3	multi-charged particle mass 785 GeV	DY production, $ q  = 5e$ 1504.04188	
Magnetic monopoles	-	-	-	7.0	monopole mass 1.34 TeV	DY production, $ g  = 1g_D, \text{spin } 1/2$ 1509.08059	

\*Only a selection of the available mass limits on new states or phenomena is shown.

<sup>†</sup>Small-radius (large-radius) jets are denoted by the letter J (j).

# BSM: Direct Searches


Direct searches of new physics gave non positive results so far  
Some examples

**Supersymmetry (MSSM)**



**Top-partners**  
**Composite Higgs**  
**Extra-dimensions**  
**Excited Quark**

...



ATLAS Exotics Searches\* - 95% CL Upper Exclusion Limits

Status: July 2017

ATLAS Preliminary  
 $\int \mathcal{L} dt = (3.2 - 37.0)\text{fb}^{-1}$   
 $\sqrt{s} = 8, 13\text{TeV}$

Search	Model	Upper Exclusion Limit	Notes
ADD $G_{KK} + g/q$	$0 < \mu, 1-4j$	Yes 36.1	$M_{\text{Pl}}$ 7.75 TeV, $n=2$
ADD OBH	$\geq 2j$	37.0	$M_{\text{BH}}$ 8.9 TeV, $n=6$
ADD BH high $\sum p_T$	$\geq 1 e, \mu, \geq 2j$	3.2	$M_{\text{BH}}$ 8.2 TeV, $n=6, M_D = 3\text{TeV}$ , rot BH
ADD BH multijet	$\geq 3j$	3.6	$M_{\text{BH}}$ 9.55 TeV, $n=6, M_D = 3\text{TeV}$ , rot BH
Bulk RS $G_{KK} \rightarrow WW \rightarrow qq\bar{q}$	$1 e, \mu, 1j$	Yes 36.1	$G_{KK}$ mass 1.75 TeV, $x/M_D = 1.0$
ZUED / RPP	$\geq 2b, \geq 3j$	Yes 13.2	$KK$ mass 1.6 TeV, $\text{Tier}(1,1), \theta(A^{1,3} \rightarrow \tau\tau) = 1$
SSM $Z' \rightarrow \ell\ell$	$2 e, \mu$	36.1	$Z'$ mass 4.5 TeV
SSM $Z' \rightarrow \tau\tau$	$2\tau$	36.1	$Z'$ mass 2.4 TeV
Leptophobic $Z' \rightarrow b\bar{b}$	$\geq 2b$	3.2	$Z'$ mass 1.5 TeV
Leptophobic $Z' \rightarrow \tau\tau$	$1 e, \mu, \geq 1b, \geq 1J/2j$	Yes 3.2	$Z'$ mass 2.0 TeV
SSM $W' \rightarrow \ell\nu$	$1 e, \mu$	Yes 36.1	$W'$ mass 5.1 TeV
HVT $V' \rightarrow WW \rightarrow qq\bar{q}$ model B	$0 e, \mu, 2j$	36.7	$V'$ mass 3.5 TeV, $g_V = 3$

Examples: models with  
**Quantum Black Holes**  
excluded up to masses  
of the BH  $\sim 10\text{TeV}$

**10 TeV**

\*Only a selection of the available mass limits on new states or phenomena is shown.  
†Small-radius (large-radius) jets are denoted by the letter (J).

# BSM: Direct Searches

Direct searches of new physics gave non positive results so far  
Some examples

**Supersymmetry (MSSM)**




**Top-partners**  
**Composite Higgs**  
**Extra-dimensions**  
**Excited Quark**

...



Selected CMS SUSY Results\* - SMS Interpretation

ICHEP '16 - Moriond '17



ATLAS Exotics Searches\* - 95% CL Upper Exclusion Limits

Status: July 2017

Model	$\ell, \gamma$	Jets <sup>†</sup>	Emiss <sup>†</sup>	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Limit	Reference
Extra dimensions	ADD $G_{KK} + g/q$	$0, e, \mu$	$1-4j$	Yes	36.1	$M_{KK}$ 7.75 TeV
	ADD non-resonant $\gamma\gamma$	$2\gamma$	-	-	36.7	$M_{KK}$ 8.6 TeV
	ADD QBH	-	$2j$	-	37.0	$M_{KK}$ 8.9 TeV
	ADD BH high $\sum p_T$	$\geq 1, e, \mu$	$\geq 2j$	-	3.2	$M_{KK}$ 8.2 TeV
	ADD BH multijet	-	$\geq 3j$	-	3.6	$M_{KK}$ 9.55 TeV
Gauge bosons	SSM $Z' \rightarrow \ell\ell$	$2, e, \mu$	-	-	36.1	$Z'$ mass 4.5 TeV
	SSM $Z' \rightarrow \tau\tau$	$2\tau$	-	-	36.1	$Z'$ mass 2.4 TeV
	Leptophobic $Z' \rightarrow bb$	-	$2b$	-	3.2	$Z'$ mass 1.9 TeV
	Leptophobic $Z' \rightarrow tt$	$1, e, \mu$	$\geq 1b, \geq 1J/2j$	Yes	3.2	$Z'$ mass 2.0 TeV
	SSM $W' \rightarrow \ell\nu$	$1, e, \mu$	-	Yes	36.1	$W'$ mass 5.1 TeV
CI	HVT $V' \rightarrow W\gamma$	$0, e, \mu$	$2j$	-	36.7	$V'$ mass 3.5 TeV
	HVT $V' \rightarrow WH/ZH$ model B	multi-channel	-	-	36.1	$V'$ mass 2.93 TeV
	LRSM $W'_\mu \rightarrow tb$	$1, e, \mu$	$2b, 0-1j$	Yes	20.3	$W'$ mass 2 TeV
	LRSM $W'_\mu \rightarrow tb$	$0, e, \mu$	$\geq 1b, 1j$	-	20.3	$W'$ mass 1 TeV
	CI $qqqq$	-	$2j$	-	37.0	$A$ 21.8 TeV
DM	Axial-vector mediator (Dirac DM)	$0, e, \mu$	$1-4j$	Yes	36.1	$m_{DM}$ 1.5 TeV
	Vector mediator (Dirac DM)	$0, e, \mu, 1\gamma$	$\leq 1j$	Yes	36.1	$m_{DM}$ 1.2 TeV
	VV <sub>KK</sub> EFT (Dirac DM)	$0, e, \mu$	$1j, \leq 1j$	Yes	3.2	$M_{KK}$ 700 GeV
	Scalar LQ 1 <sup>st</sup> gen	$2, e$	$\geq 2j$	-	3.2	LQ mass 1.1 TeV
	Scalar LQ 2 <sup>nd</sup> gen	$2, \mu$	$\geq 2j$	-	3.2	LQ mass 1.05 TeV
Heavy quarks	Scalar LQ 3 <sup>rd</sup> gen	$1, e, \mu$	$\geq 1b, \geq 3j$	Yes	20.3	LQ mass 640 GeV
	VLO $TT \rightarrow Ht + X$	$0$ or $1, e, \mu$	$\geq 2b, \geq 3j$	Yes	13.2	$T$ mass 1.2 TeV
	VLO $TT \rightarrow Zt + X$	$1, e, \mu$	$\geq 1b, \geq 3j$	Yes	36.1	$T$ mass 1.16 TeV
	VLO $TT \rightarrow Wb + X$	$1, e, \mu$	$\geq 1b, \geq 1J/2j$	Yes	36.1	$T$ mass 1.35 TeV
	VLO $BB \rightarrow Hb + X$	$1, e, \mu$	$\geq 2b, \geq 3j$	Yes	20.3	$B$ mass 700 GeV
Excited fermions	VLO $BB \rightarrow Zb + X$	$2j \geq 3, e, \mu$	$\geq 2j \geq 1b$	-	20.3	$B$ mass 790 GeV
	VLO $BB \rightarrow Wt + X$	$1, e, \mu$	$\geq 1b, \geq 1J/2j$	Yes	36.1	$B$ mass 1.25 TeV
	VLO $QQ \rightarrow WqWq$	$1, e, \mu$	$\geq 4j$	Yes	20.3	$Q$ mass 690 GeV
	Excited quark $q^* \rightarrow qg$	-	$2j$	-	37.0	$q^*$ mass 6.0 TeV
	Excited quark $q^* \rightarrow q\gamma$	$1\gamma$	$1j$	-	36.7	$q^*$ mass 5.3 TeV
Other	Excited quark $b^* \rightarrow bg$	-	$1b, 1j$	-	13.3	$b^*$ mass 2.3 TeV
	Excited quark $b^* \rightarrow Wt$	$1$ or $2, e, \mu$	$1b, 2-0j$	Yes	20.3	$b^*$ mass 1.5 TeV
	Excited lepton $\ell^*$	$3, e, \mu, \tau$	-	-	20.3	$\ell^*$ mass 3.0 TeV
	Excited lepton $\nu^*$	$3, e, \mu, \tau$	-	-	20.3	$\nu^*$ mass 1 TeV
	LRSM Majorana $\nu$	$2, e, \mu$	$2j$	-	20.3	$\nu$ mass 0 TeV
Higgs triplet $H^{\pm\pm} \rightarrow \ell\ell$	$2, 3, 4, e, \mu$ (SS)	-	-	36.1	$H^{\pm\pm}$ mass 870 GeV	
Higgs triplet $H^{\pm\pm} \rightarrow \ell\tau$	$3, e, \mu, \tau$	-	-	20.3	$H^{\pm\pm}$ mass 400 GeV	
Monotop (non-res prod)	$1, e, \mu$	$1b$	Yes	20.3	spin-1 invisible particle mass 637 GeV	
Multi-charged particles	-	-	-	20.3	multi-charged particle mass 785 GeV	
Magnetic monopoles	-	-	-	7.0	monopole mass 1.34 T	

\*Only a selection of the available mass limits on new states or phenomena is shown.

†Small-radius (large-radius) jets are denoted by the letter J (j).

1 TeV


# BSM: Direct Searches

Direct searches of new physics gave non positive results so far  
Some examples

**Supersymmetry (MSSM)**

**Top-partners**  
**Composite Higgs**  
**Extra-dimensions**  
**Excited Quark**

...



1 TeV


# An example: Models with two Higgs doublets

## Another possible benchmark with two Higgs doublets

Deviations in couplings up to 10%


We can expect deviations, even if we do not see directly the new particles

Higgs sector is new, and weird.... we must do our best to measure all the properties in the best way we can




# Energy, intensity, and accuracy frontiers

R. Torre HL/HE LHC workshop



## New physics: a pictorial representation






# High $p_T$ Higgs


## Higgs at high $p_T$

- Large statistics of Higgs at high  $p_T$ :  
 $10^6$   $H$  with  $p_T > 1.5$  TeV and  
 $10$   $H$  with  $p_T > 8$  TeV ( $20 \text{ ab}^{-1}$ )
- For  $p_T > 0.8$  TeV,  $ttH > gg \rightarrow H$   
 For  $p_T > 1.8$  TeV,  $VBF > gg \rightarrow H$
- Background and systematics considerations can be very different from LHC
- At high  $p_T$  better discriminating power  
 $H \rightarrow bb$  with jet sub-structure
- Test of Higgs couplings at high energy G.Giudice - ICFA2017



**10% precision at  $E = \text{TeV}$  probes  
 New Physics as much 0.1%  
 precision in Higgs decays**


## New physics: a pictorial representation



# Higgs couplings

- LHC coupling measurements at the order of 10-20%
- Projections for HL-LHC: O(5%)
  - based on Run1 experience
  - iper-conservative?
- $e^+e^-$  colliders can provide model independent measurements for the Higgs couplings O(1%)
- What can the HL/HE-LHC program say about coupling with first and second generation?
  - If SM,  $H\mu\mu$  can be observed in HL-LHC
  - Which options for  $Hcc$  at LHC?
    - Direct searches? Wh asymmetries?
- and self-coupling?

Peskin @ ICFA2017



(HL- LHC measurements are model dependent)

Tenchini @ FCC week 2017

in %	HL-LHC	FCC-ee
$g_{HZ}$	<b>2-4</b>	<b>0.21</b>
$g_{HW}$	<b>2-5</b>	<b>0.43</b>
$g_{Hb}$	<b>5-7</b>	<b>0.64</b>
$g_{Hc}$	-	<b>1.0</b>
$g_{Hg}$	<b>3-5</b>	<b>1.2</b>
$g_{H\tau}$	<b>5-8</b>	<b>0.81</b>
$g_{H\mu}$	<b>5</b>	<b>8.8</b>
$g_{H\gamma}$	<b>2-5</b>	<b>2.1</b>
$\Gamma_H$	<b>5-8%</b>	<b>1.5</b>

# up/down-type fermion and lepton/quark asymmetries

## Asymmetries in Higgs couplings

- between up-type and down-type fermion
- between lepton and quark

predicted by several BSM physics models

Parameterise model in terms of ratios of coupling strength modifiers

$$\lambda_{du} = \kappa_d / \kappa_u$$



$$\lambda_{Vu} = \kappa_V / \kappa_u$$

$$\kappa_{uu} = \kappa_u \cdot \kappa_u / \kappa_H$$

$$\lambda_{\ell q} = \kappa_\ell / \kappa_q$$


$$\lambda_{Vq} = \kappa_V / \kappa_q$$

$$\kappa_{qq} = \kappa_q \cdot \kappa_q / \kappa_H$$




# Effective couplings and BSM BR

- What if we have new particles in the loops?
- Specific fit not resolving the loops, but use effective couplings  $\kappa_g$  and  $\kappa_\gamma$




Production	Loops	Interference	Effective scaling factor
$\sigma(ggF)$	✓	$t-b$	$\kappa_g^2$
Partial decay width			
$\Gamma_{\gamma\gamma}$	✓	$t-W$	$\kappa_\gamma^2$

- Fix all tree-level Higgs couplings to SM ( $\kappa_W, \kappa_Z, \kappa_b, \kappa_t, \kappa_\mu, \kappa_\tau = 1$ ) and  $B_{BSM} = 0$




# Effective couplings and BSM BR

- What if we have new particles in the loops?
- Specific fit not resolving the loops, but use effective couplings  $\kappa_g$  and  $\kappa_\gamma$




Production	Loops	Interference	Effective scaling factor
$\sigma(ggF)$	✓	$t$ - $b$	$\kappa_g^2$
Partial decay width			
$\Gamma_{\gamma\gamma}$	✓	$t$ - $W$	$\kappa_\gamma^2$




# Effective couplings and BSM BR


- What if we have new particles in the loops?
- Specific fit not resolving the loops, but use effective couplings  $\kappa_g$  and  $\kappa_\gamma$
- And if the Higgs boson decays in some other mode we did not detect yet?
  - Constrain  $B_{\text{BSM}} \geq 0$  and  $|\kappa_V| \leq 1$



Production	Loops	Interference	Effective scaling factor
$\sigma(ggF)$	✓	$t$ - $b$	$\kappa_g^2$
Partial decay width			
$\Gamma_{\gamma\gamma}$	✓	$t$ - $W$	$\kappa_\gamma^2$




# Ratios of production Cross Sections and BR



- Measuring ratios of production cross sections and BR
 
$$\sigma_i \cdot B^f = \sigma(gg \rightarrow H \rightarrow ZZ) \cdot \left( \frac{\sigma_i}{\sigma_{ggF}} \right) \cdot \left( \frac{B^f}{B^{ZZ}} \right)$$
- No additional SM assumption on these measurements
- p-value(SM) = 16% ( $\sim 1\sigma$ )


# Ratios of production Cross Sections and BR



- Measuring ratios of production cross sections and BR
 
$$\sigma_i \cdot B^f = \sigma(gg \rightarrow H \rightarrow ZZ) \cdot \left( \frac{\sigma_i}{\sigma_{ggF}} \right) \cdot \left( \frac{B^f}{B^{ZZ}} \right)$$
- No additional SM assumption on these measurements
- p-value(SM) = 16% ( $\sim 1\sigma$ )
- $\sigma_{ZH}/\sigma_{ggF} \sim 3$ , mainly due to ZH,  $H \rightarrow WW$




# Ratios of production Cross Sections and BR



- Measuring ratios of production cross sections and BR  

$$\sigma_i \cdot B^f = \sigma(gg \rightarrow H \rightarrow ZZ) \cdot \left( \frac{\sigma_i}{\sigma_{ggF}} \right) \cdot \left( \frac{B^f}{B^{ZZ}} \right)$$
- No additional SM assumption on these measurements
- p-value(SM) = 16% ( $\sim 1\sigma$ )
- $\sigma_{ZH}/\sigma_{ggF} \sim 3$ , mainly due to ZH,  $H \rightarrow WW$
- $\sigma_{ttH}/\sigma_{ggF} \sim 3\sigma$  excess with respect to SM due to ttH,  $H \rightarrow$  multi lepton: WW/ $\tau\tau$ /(ZZ)

# Ratios of production Cross Sections and BR



- Measuring ratios of production cross sections and BR
- $$\sigma_i \cdot B^f = \sigma(gg \rightarrow H \rightarrow ZZ) \cdot \left( \frac{\sigma_i}{\sigma_{ggF}} \right) \cdot \left( \frac{B^f}{B^{ZZ}} \right)$$
- No additional SM assumption on these measurements


• p-value(SM) = 16% ( $\sim 1\sigma$ )

•  $\sigma_{ZH}/\sigma_{ggF} \sim 3$ , mainly due to ZH,  $H \rightarrow WW$

•  $\sigma_{ttH}/\sigma_{ggF} \sim 3\sigma$  excess with respect to SM due to ttH,  $H \rightarrow$  multi lepton: WW/ $\tau\tau$ /(ZZ)

High ZH,  $H \rightarrow WW$ ,  
High ttH,  $H \rightarrow$  multi lept  
Low ZH,  $H \rightarrow bb$   
contribute to...

•  $B^{bb}/B^{ZZ}$ :  
deficit  $\sim 2.5\sigma$   
with respect to SM



# Signal strengths at the end of Run1

- Measurements of signal strengths  $\mu$  for each production mode and for each decay mode by fixing the relative  $B^f$  or the  $\sigma_i$  to the SM prediction.

$$n_{\text{signal}}(k) = \mathcal{L}(k) \cdot \sum_i \sum_j \mu_i \mu^f \left\{ \sigma_i^{\text{SM}} \cdot A_i^{f,\text{SM}}(k) \cdot \epsilon_i^f(k) \cdot B_{\text{SM}}^f \right\}$$



Production process	Measured significance ( $\sigma$ )	Expected significance ( $\sigma$ )
VBF	5.4	4.6
WH	2.4	2.7
ZH	2.3	2.9
VH	3.5	4.2
ttH	4.4	2.0
Decay channel		
$H \rightarrow \tau\tau$	5.5	5.0
$H \rightarrow bb$	2.6	3.7

- By fixing all the  $B^f$  and  $\sigma_i$  to the SM prediction, and allowing for only one global signal strength, one gets:


$$\mu = 1.09_{-0.10}^{+0.11} = 1.09_{-0.07}^{+0.07} (\text{stat})_{-0.04}^{+0.04} (\text{expt})_{-0.03}^{+0.03} (\text{thbgd})_{-0.06}^{+0.07} (\text{thsig}),$$

Assuming SM Decay BR

Assuming SM Production XS



# Physics backgrounds at hadron colliders



LHC experiments need to be able to select every second the  $O(200)$  most interesting events out of  $10^9$  collisions that took place


jets

high- $p_T$  jets

W production


H production

# Verifying the SM, Preparing the ground for a discovery




# Verifying the SM, Preparing the ground for a discovery

~ Production Probability




# Verifying the SM, Preparing the ground for a discovery

~ Production Probability



# Verifying the SM, Preparing the ground for a discovery


~ Production Probability






# Verifying the SM, Preparing the ground for a discovery

~ Production Probability




# Verifying the SM, Preparing the ground for a discovery

~ Production Probability




# Verifying the SM, Preparing the ground for a discovery

~ Production Probability




# Verifying the SM, Preparing the ground for a discovery

~ Production Probability




# Combination of the results

- $p_0$ : probability that the data could come from a model with no Higgs boson.
- Very high standards:  
**Evidence benchmark**  $\rightarrow p_0 = 0.00135$  (3 Gaussian standard deviations)  
**Discovery benchmark**  $\rightarrow p_0 = 2.6 \times 10^{-7}$  (5 Gaussian standard deviations)




**5.9 s.d.** at  $m_H = 126.5 \text{ GeV}$



**5.0 s.d.** at  $m_H = 125.5 \text{ GeV}$


(The analysis of more data since July 4, 2012 has further strengthened the significance)

# The Big news (1): Top -Higgs interaction




$$H \rightarrow ZZ^* \rightarrow 4\ell$$

$$H \rightarrow \gamma\gamma$$



**ATLAS 4.1 $\sigma$  (3.7 $\sigma$  exp.) (80fb<sup>-1</sup>)**



**CMS 1.4 $\sigma$  (1.5 $\sigma$  exp.)**



$$H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$$


$$H \rightarrow \tau\tau$$

(multi-leptons)



$$H \rightarrow b\bar{b}$$


[arXiv:1804.03682]  
[Phys. Rev. D 97 (2018) 072016]



(CMS: incl. all-hadronic channel)

# Some more considerations: hierarchy and naturalness

- Plank Scale
  - $\sim 10^{19}$  GeV
- Scale at which  $\lambda$  becomes negative : vacuum instability (with current measured  $m_{\text{top}}$  and  $m_H$ )
  - $\sim 10^{16}$  GeV
- Mass of the particles (other than Higgs): 0-100 GeV
  - They should be 0 for gauge symmetries , with a “small” correction from the Higgs condensate
- But why the condensate is at 250 GeV and the mass of the Higgs is at 125 GeV?
  - Just by chance?



$$m_H^2 = m_{H0}^2 - \frac{3}{8\pi^2} y_t^2 \Lambda^2 + \frac{1}{16\pi^2} g^2 \Lambda^2 + \frac{1}{16\pi^2} \lambda^2 \Lambda^2$$

*Quantum correction to  $m_H$ ,  
assuming as cut off scale  
due to new physics  $\Lambda$*


Putting numbers, one gets:

$$(125 \text{ GeV})^2 = m_{H0}^2 + [-(2 \text{ TeV})^2 + (700 \text{ GeV})^2 + (500 \text{ GeV})^2] \left( \frac{\Lambda}{10 \text{ TeV}} \right)^2$$

# Some more considerations: hierarchy and naturalness

- Planck scale
- Scalar masses
- Masses
- But and

$$m_H^2 \sim (125 \text{ GeV})^2$$



$$(125 \text{ GeV})^2 = m_{H_0}^2 + [-(2 \text{ TeV})^2 + (700 \text{ GeV})^2 + (500 \text{ GeV})^2] \left( \frac{\Lambda}{10 \text{ TeV}} \right)^2$$

Definition of naturalness: less than 90% cancellation:

$$\Lambda_t < 3 \text{ TeV}$$

$\Rightarrow$  top partners must be "light"

$$(125 \text{ GeV})^2 = m_{H_0}^2 + [-(2 \text{ TeV})^2 + (700 \text{ GeV})^2 + (500 \text{ GeV})^2] \left( \frac{\Lambda}{10 \text{ TeV}} \right)^2$$

Quantum  
assumptions  
due to



# Object Resolution

