

Production of muons from open heavy-flavour hadron decays in heavy-ion collisions with ALICE

Zuman Zhang for the ALICE Collaboration


Institute of Particle Physics, CCNU, Wuhan, China

Key Laboratory of Quark & Lepton Physics, MoE, China

Laboratoire de Physique de Clermont, CNRS/IN2P3, Clermont-Ferrand, France

Outline

- Physics motivation
- Open heavy-flavour measurements with the ALICE muon spectrometer
- Analysis procedure
- Results
- Summary and outlook



**QGP-France, 2-5 July 2018,
Etretat, France**


Charm and beauty quarks: **sensitive probes of the medium properties**

Open heavy flavours in nucleus-nucleus (AA) collisions probe

- ❑ In-medium parton energy loss: gluon radiation and elastic collisions
- ❑ Heavy-quark participation in the collective expansion

Also needed:

- ❑ **proton-proton (pp) collisions**
reference, tests of pQCD-based predictions
- ❑ **pA collisions**
reference, cold nuclear matter effects




Observable

- ✓ The nuclear modification factor, R_{AA} ,
sensitive to the medium effects

$$R_{AA}(p_T) = \frac{1}{\langle T_{AA} \rangle} \times \frac{dN_{AA}/dp_T}{d\sigma_{pp}/dp_T} = \frac{QCD \text{ Medium}}{QCD \text{ Vacuum}}$$

- ✓ If no nuclear effects: $R_{AA} = 1$
- ✓ Effects of the hot and dense medium produced in the collision breakup binary scaling: $R_{AA} \neq 1$

ALICE Run-2 setup




Data samples and muon selections

	pp collisions at $\sqrt{s} = 5.02$ TeV	Xe-Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV	Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV
Data	LHC15n, collected in 2015	LHC17n, collected in 2017	LHC15o, collected in 2015
L_{int}	MSL(MSH): ≈ 53.7 (104.4) nb^{-1}	MSL: ≈ 0.34 μb^{-1}	MSL(MSH): ≈ 21.9 (202.3) μb^{-1}

MSL (MSH) triggers: p_T threshold $\sim 0.5, 1$ (4.2) GeV/c

Muon track selection

- **Acceptance & geometrical cuts**
select tracks in the spectrometer acceptance
- **p_T cut at 2 GeV/c**
reject μ from secondary π, K
- **Muon tracking tracks matched with muon trigger tracks**
reject hadrons crossing the front absorber
- **$p \times \text{DCA}$ (Dist. of Closest Approach) in 6σ**
reject beam-gas interactions & particles produced in the absorber



background

$\mu^\pm \leftarrow b, c$ studies

▪ Remaining background

- $\mu \leftarrow$ primary π, K decays (main contribution at low p_T)
- $\mu \leftarrow W, Z/\gamma^*$ decays (main contribution at high p_T)
- $\mu \leftarrow J/\psi$ decays (dominates at $p_T \sim 5 \text{ GeV}/c$)

Analysis procedure in pp collisions



$$R_{AA}(p_T) = \frac{1}{\langle T_{AA} \rangle} \cdot \frac{dN_{AA}^{\text{inclusive } \mu} / dp_T - dN_{AA}^{\text{bkg}} / dp_T}{d\sigma_{pp}^{\mu \leftarrow \text{HF}} / dp_T}$$

- Get inclusive muons with muon-triggered data, after muon event and track selection. Then **normalize inclusive muons to minimum-bias events** and **apply acceptance x efficiency correction**
- **Background: $\mu \leftarrow \pi/K$** (dominates at low p_T , max. ~40% at 2 GeV/c)
 - ✓ Inputs: π/K spectra at mid-rapidity at 5.02 TeV, extrapolated to higher p_T via power-law fit
 - ✓ Then, get π/K spectra in 4π with p_T -dependent rapidity shape in Monte-Carlo via:

$$\frac{1}{N_{\text{ev}}} \frac{d^2 N_{pp}^{K/\pi}}{dp_T dy} = \frac{1}{N_{\text{ev}}} \frac{dN_{pp}^{K/\pi}}{dp_T} \Big|_{|y| < 0.8} \times F(p_T, y)$$

- ✓ Produce the π/K decay muon background in Monte-Carlo with fast simulation with parametrized muon front absorber response
- **Background: $\mu \leftarrow W, Z/\gamma^*$** (dominates at high p_T , max. ~13% in [18,20] GeV/c)
 - ✓ Obtained from pp collisions by (POWHEG) simulation with CT10 PDF
- **Background: $\mu \leftarrow J/\psi$** (dominates at $p_T \sim 5$ GeV/c, small compared to $\mu \leftarrow \pi/K$)
 - ✓ Use J/ψ extrapolated p_T and y distributions as inputs, convert J/ψ spectra to muons with fast simulation, max. ~4% contribution at $p_T \sim 5$ GeV/c

$$R_{AA}(p_T) = \frac{1}{\langle T_{AA} \rangle} \cdot \frac{dN_{AA}^{\text{inclusive } \mu} / dp_T - dN_{AA}^{\text{bkg}} / dp_T}{d\sigma_{pp}^{\mu \leftarrow \text{HF}} / dp_T}$$

- Get inclusive muons with muon-triggered data, after muon event and track selection. Then **normalize inclusive muons to minimum-bias events and apply acceptance x efficiency correction**
- **Background: $\mu \leftarrow \pi/K$ (dominates at low p_T)**
 - ✓ Inputs: π/K spectra at mid-rapidity, extrapolated to higher p_T
 - ✓ Then, get π/K spectra in 4π with p_T -dependent rapidity shape in Monte-Carlo via:

$$\begin{aligned} \frac{d^2 N_{AA}^{K/\pi}}{dp_T dy} &= n_y \times R_{AA} \times F(p_T, y) \times \frac{dN_{pp}^{K/\pi}}{dp_T} \Big|_{|y| < 0.8} \\ &= n_y \times F(p_T, y) \times \frac{dN_{AA}^{K/\pi}}{dp_T} \Big|_{|y| < 0.8} \end{aligned}$$

$n_y (= 1)$: quenching factor, systematic uncertainty varying n_y within 0.5-1.5




$F(p_T, y)$: p_T -dependent y extrapolation factor, from pp simulations with Monte-Carlo event generators

- ✓ Produce the π/K decay muon background in Monte-Carlo with fast simulation with parametrized muon front absorber response
- Background: $\mu \leftarrow J/\psi$ (dominates at $p_T \sim 5 \text{ GeV}/c$, small compared to $\mu \leftarrow \pi/K$)
- Background: $\mu \leftarrow W, Z/\gamma^*$ (dominates at high p_T , obtained from pp, pn, np and nn collisions by (POWHEG) simulation)




ALICE

pp results:
differential production cross sections of
muons from heavy-flavour hadron decays




Run:282016
Timestamp:2017-11-11 21:38:31(UTC)
Colliding system:p-p
Energy: 5.02 TeV

p_T -differential production cross section (1/2)




ALI-PREL-152053




- ❑ Measurement over a **wide p_T range**: $2 < p_T < 20$ GeV/c in pp collisions at 5.02 TeV
- ❑ Data in **agreement with FONLL predictions** within uncertainties although at the upper edge of FONLL predictions
- ❑ Quite precise measurement, strong constraints on pQCD-based calculations
- ❑ The **reference** for Pb-Pb measurements at $\sqrt{s_{NN}} = 5.02$ TeV
- ❑ Systematic uncertainties **reduced by about a factor two**, compared to Run-1 results

p_T -differential production cross section (2/2)



ALI-PREL-152040




- $\sqrt{s} = 5.02$ TeV measurement in $3 < p_T < 8$ GeV/c scaled to $\sqrt{s} = 5.44$ TeV with FONLL
- Data in **agreement with FONLL predictions**, strong constraints on pQCD-based calculations
- The **reference** for Xe-Xe measurements at $\sqrt{s_{NN}} = 5.44$ TeV



Xe-Xe and Pb-Pb results:
nuclear modification factor vs p_T and
centrality of muons from heavy-flavour
hadron decays

Run:280235
Timestamp:2017-10-13 00:31:48(UTC)
Colliding system:Xe-Xe
Energy: 5.44 TeV




ALI-PREL-152081

- Measurement in $3 < p_T < 8$ GeV/c for 10% most central collisions ($3 < p_T < 7$ GeV/c for 10-20%, 20-40%, 40-60%) with muon-triggered events
- Increasing normalized p_T -differential yields from peripheral to central collisions


p_T -differential R_{AA} of $\mu^\pm \leftarrow b, c$

$$R_{AA}(p_T) = \frac{1}{\langle T_{AA} \rangle} \times \frac{dN_{AA}/dp_T}{d\sigma_{pp}/dp_T}$$

$\langle T_{AA} \rangle$ values, see backup



ALI-PREL-152146




ALI-PREL-152163

- p_T -differential R_{AA} of heavy-flavour decay muons in different centrality classes
- Increasing suppression from peripheral to central collisions


p_T -differential R_{AA} of $\mu^\pm \leftarrow b, c$ in Pb-Pb collisions

Centrality: 0-10%, 5.02 vs 2.76 TeV

Vitev: Phys. Rev. C 80 (2009) 054902
TAMU: Phys. Lett. B 735 (2014) 445






ALI-PREL-116429



ALI-PREL-116437


- Similar suppression at 5.02 TeV and at 2.76 TeV for central collisions within uncertainties
- Better precision in Run 2 ($\sqrt{s_{NN}} = 5.02$ TeV)
- R_{AA} measurements at $\sqrt{s_{NN}} = 5.02$ TeV provide new constraints on energy loss models

p_T -differential R_{AA} of $\mu^\pm \leftarrow b$ in Pb-Pb collisions: comparison with MC@HQ + EPOS2




□ R_{AA} measurements at $\sqrt{s_{NN}} = 5.02$ TeV provide new constraints on energy loss models

R_{AA} : $\sqrt{s_{NN}} = 5.44$ TeV in Xe-Xe collisions vs $\sqrt{s_{NN}} = 5.02$ TeV in Pb-Pb collisions



ALI-PREL-152237




ALI-PREL-152264

- Similar heavy-flavour hadron decay muon R_{AA} observed in 0-10% Xe-Xe and 10-20% Pb-Pb collisions at similar charged-particle multiplicity
- ✓ Possible interplay of geometry and path-length dependence


Comparison of Xe-Xe and Pb-Pb results with p-Pb measurements

Centrality: 0-10%, Xe-Xe vs 0-10%, Pb-Pb



ALI-PREL-152237

p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV




ALI-PUB-127462

Phys. Lett B 770 (2017) 459

- R_{pPb} : consistent with unity within uncertainties over the whole p_T range
- The suppression observed in central Xe-Xe and Pb-Pb collisions results from final-state effects related to parton energy loss



Comparison with measurements of $e \leftarrow c, b$ (mid-rapidity)



ALI-PREL-148699

- Compatible results within uncertainties for heavy-flavour decay electrons ($|y| < 0.8$) and heavy-flavour decay muons ($2.5 < y < 4$) R_{AA} in 0-10% and 20-40% centrality classes
- Indication that heavy quarks suffer a strong interaction in a wide rapidity interval

Comparison with model calculations



- Centrality: 20-40%, models describe the measured R_{AA} within uncertainties;
- Centrality: 10-20%, models overestimate the measured R_{AA}
- R_{AA} measurements have the potential to constrain energy loss models

PHSD model: arXiv:1803.02698 and Phys.Rev. C96 (2017) no.1, 014905

Conclusion

- ❑ Measurements in pp collisions, over **a wider p_T range**, extended to $p_T = 20$ GeV/c
 - Precise reference for the R_{AA} computation
 - Described by pQCD-based calculations (FONLL)

- ❑ Measurements in Xe-Xe collisions, **new system**, w.r.t Pb-Pb collisions
 - **Strong suppression**, a factor ~ 3 for $6 < p_T < 8$ GeV/c in 0-10% centrality class
 - The measured suppression is **due to final-state effects** ($R_{pPb} \sim 1$)
 - Results compatible within uncertainties with those obtained at $\sqrt{s_{NN}} = 5.02$ TeV Pb-Pb collisions and with mid-rapidity electrons from heavy-flavour hadron decays
 - R_{AA} measurements have the potential to constrain energy loss models

More to come soon

- Elliptic flow, v_2 , measurement in Run-2
- More differential measurements with the next Pb-Pb run scheduled end of 2018



Thank you for your attention

Backup



ALICE

Charm and beauty quarks: **sensitive probes of the medium properties**

Open heavy flavours in nucleus-nucleus (A-A) collisions probe

- ❑ In-medium parton energy loss
- ❑ Heavy-quark participation in the collective expansion

Also needed:


- ❑ **proton-proton (pp) collisions**
reference, tests of pQCD-based predictions
- ❑ **p-A collisions**
reference, cold nuclear matter effects

Observable

- ✓ The nuclear modification factor, R_{AA} , sensitive to the medium effects



$$R_{AA}(p_T) = \frac{1}{\langle T_{AA} \rangle} \times \frac{dN_{AA}/dp_T}{d\sigma_{pp}/dp_T} = \frac{QCD \text{ Medium}}{QCD \text{ Vacuum}}$$

- ✓ If no nuclear effects: $R_{AA} = 1$



Run-2 provides measurements of heavy-flavour hadron decay muons in a wide p_T range with improved precision which bring new constraints on in-medium energy loss

Comparison with models



□ Elliptic flow v_2 : complementary measurement to R_{AA} sensitive to:

- Low p_T : collective motion
- High p_T : path-length dependence of parton energy loss

□ Positive v_2 measured with a significance $> 3\sigma$ for $3 < p_T < 5$ GeV/c in 20-40%

□ Confirmation of significant interaction of heavy quarks with the medium

□ Simultaneous description of R_{AA} in central collisions and v_2 in semi-central collisions is challenging

□ R_{AA} and v_2 measurements starts to provide constraints on energy loss models

□ Similar picture for heavy-flavour decay electrons and D mesons






ALICE

Analysis strategy in Pb-Pb collisions (2/5)

Muons from K/ π decays

Inputs, from QM2017 approval



Get K/ π spectra **at mid-rapidity** in Pb-Pb collisions at 5.02 TeV and do p_T extrapolation

✓ p_T extrapolation with power-law fit



(1) $p_T < 12$ GeV/c : Gaussian fit with gRandom->Gaus(μ, σ)

μ : center value of K/ π spectra; σ : stat. and syst. uncertainties of data

(2) $p_T > 12$ GeV/c

μ and σ from K/ π spectra in p_T -extrapolated intervals

R_{AA} in Xe-Xe collisions at 5.44 TeV



ALI-PREL-152264

T_{AA} values in Xe-Xe and Pb-Pb collisions



Xe-Xe

Cent (%)	Npart	RMS	Sys	Ncoll	RMS	Sys	TAA	RMS	Sys
0-10%	221.2	19	2.2	843.1	1.4e+02	70	12.33	2	1
10-20 %	164.8	18	2.8	510.6	86	51	7.465	1.3	0.74
20-30 %	118.4	14	3.8	302.8	58	40	4.426	0.85	0.59
30-40 %	82.21	11	3.9	171.3	38	27	2.505	0.56	0.4
40-50 %	54.56	8.8	3.6	91.81	24	16	1.342	0.35	0.24
50-60 %	34.06	6.5	3	46.04	14	8.8	0.6731	0.2	0.13
60-70 %	19.72	4.7	2.1	21.65	7.6	4.1	0.3166	0.11	0.061
70-80 %	10.5	3.1	1.1	9.515	3.9	1.6	0.1391	0.056	0.024
80-90 %	5.127	1.9	0.46	3.838	1.9	0.5	0.05611	0.028	0.0074
90-100 %	2.488	0.8	0.12	1.449	0.73	0.1	0.02118	0.011	0.0015

Pb-Pb

Cent	bmin [fm]	bmax [fm]	Npart	RMS	Sys	Ncoll	RMS	Sys	TAA	RMS	Sys
00 - 10 %	0.00	4.96	359	31.2	3.0	1636	246	170	23.4	3.51	0.78
10 - 20 %	4.96	7.01	263	27.1	3.6	1001	154	97	14.3	2.2	0.46
20 - 30 %	7.01	8.59	188	22.5	3.0	601	106	54	8.59	1.52	0.27
30 - 40 %	8.59	9.92	131	19.1	2.3	344	74.7	29	4.92	1.07	0.16
40 - 50 %	9.92	11.1	86.3	16.3	1.7	183	50.8	14	2.61	0.726	0.1
50 - 60 %	11.1	12.1	53.6	13.6	1.2	89.8	32.4	6	1.28	0.463	0.063
60 - 70 %	12.1	13.1	30.4	10.8	0.76	39.8	19.1	2.4	0.569	0.273	0.032
70 - 80 %	13.1	14.0	15.6	7.83	0.45	16.2	10.5	0.92	0.232	0.15	0.015
80 - 90 %	14.0	15.0	7.59	4.89	0.19	6.57	5.27	0.3	0.0923	0.0753	0.007
90 - 100 %	15.0	19.6	3.77	2.5	0.079	2.66	2.41	0.088	0.0378	0.0344	0.0033