

Measurement of the Effective Weak-mixing Angle $\sin^2 \theta_{eff}^l$ in $p\bar{p} \rightarrow Z/\gamma^* \rightarrow l^+l^-$

Chen Wang

University of Science and Technology of China

2018.04.25

➤ Overview

➤ 2017, $8.6fb^{-1}$ RunIIb
Muon Channel
Precision:0.00064
Best muon channel to date

➤ 2011, $1.1fb^{-1}$ RunIIa
+ $3.9fb^{-1}$ RunIIb
Electron Channel
Precision:0.0010

➤ D0 combination
Precision:0.00040
Best single experiment to date
Best light-quark measurement



➤ 2008, $1.1fb^{-1}$ RunIIa
Electron Channel
Precision:0.0019
First hadron measurement

➤ 2015, $1.1fb^{-1}$ RunIIa
+ $8.6fb^{-1}$ RunIIb
Electron Channel
Precision:0.00047
Best single channel to date
First time close to LEP/SLD

➤ All D0 RunII weak-mixing angle measurements

➤ Strategy

- Measured from raw A_{FB}
- Statistical dominated
- PDF uncertainty
- Systematic uncertainty: Lepton energy calibration

➤ Measured from raw $A_{FB}(M_{ll})$

- Not from unfolded $A_{FB}(M_{ll})$
- Same sensitivity
- Extra systematic uncertainty from the response matrix

➤ $1.1fb^{-1}$ electron channel measurement (2008)

➤ First hadron measurement

- Tevatron RunIIa, 35K events
- Measured by observing $A_{FB}(M_{ll})$
- PDF:CTEQ6L
- Simple higher-order correction(ZGRAD2)

$$\begin{aligned}\sin^2 \theta_{eff}^l &= 0.2326 \pm 0.0019 \\ &= 0.2326 \pm 0.0018(stat.) \pm 0.0003(syst.) \pm 0.0005(PDF)\end{aligned}$$

	$1.1fb^{-1}$ results	Theoretical expectation for $10fb^{-1}$ *
σ_{stat}	0.0018	0.0005
σ_{syst}	0.0003	0.00007
σ_{PDF}	0.0005	0.00007
σ_{Total}	0.0019	~0.0005

*arXiv:hep-ex/0011009

➤ $5fb^{-1}$ electron channel measurement (2011)

➤ Mid-term estimation

- Tevatron RunIIa+part of RunIIb, 160K events
- Measured by observing $A_{FB}(M_{ll})$
- PDF:CTEQ6L

$$\begin{aligned}\sin^2 \theta_{eff}^l &= 0.2309 \pm 0.0010 \\ &= 0.2309 \pm 0.0008(stat.) \pm 0.00029(syst.) \pm 0.00048(PDF)\end{aligned}$$



	$5fb^{-1}$ results	Theoretical expectation for $10fb^{-1}$	Expectation for $10fb^{-1}$ using $5fb^{-1}$ results
σ_{stat}	0.0008	0.0005	> 0.0006
σ_{syst}	0.00029	0.00007	0.0003
σ_{PDF}	0.00048	0.00007	~ 0.00048
σ_{Total}	0.00048	~ 0.0005	~ 0.00085

➤ Systematic uncertainty

➤ Lepton calibration

- Affects $A_{FB}(M_{ll})$ same with $\sin^2 \theta_W$ does
- Very sensitive



$$\frac{\delta M}{M} / \sigma_{\sin^2 \theta_W} \sim 0.01\% / 0.00003$$



➤ Lepton calibration

$$\frac{\delta M}{M} / \sigma_{\sin^2 \theta_W} \sim 0.01\% / 0.00003$$

- A mass calibration
 - Only mass scale affects $A_{FB}(M_{ll})$ at Tevatron
 - At LHC, lepton energy scale directly affects $A_{FB}(M_{ll})$
- In LHC, E_l becomes more important
 - Quark direction, according to Z boson boost(p_Z)
 - May change the direction/the sign of p_Z



➤ $9.7fb^{-1}$ electron channel measurement (2015)

- First high precision measurement
- Best electron channel measurement to date
 - Tevatron RunII, ~560K events
 - Measured by observing $A_{FB}(M_{ll})$
 - PDF: NNPDF2.3/3.0

$$\begin{aligned}\sin^2 \theta_{eff}^l &= 0.23137 \pm 0.00047 \\ &= 0.23137 \pm 0.00043(stat.) \pm 0.00009(syst.) \pm 0.00017(PDF)\end{aligned}$$

	$9.7fb^{-1}$ results	Theoretical expectation for $10fb^{-1}$	Expectation for $10fb^{-1}$ using $5fb^{-1}$ results
σ_{stat}	0.00043	0.0005	> 0.0006
σ_{syst}	0.00009	0.00007	0.0003
σ_{PDF}	0.00017	0.00007	~0.0002
σ_{Total}	0.00047	~0.0005	~0.00085

➤ $8.6fb^{-1}$ muon channel measurement (2017)



➤ Best muon channel measurement to date

- Tevatron RunIIb, ~480K events
- Measured by observing $A_{FB}(M_{ll})$
- PDF: NNPDF3.0

$$\begin{aligned}\sin^2 \theta_{eff}^l &= 0.23016 \pm 0.00064 \\ &= 0.23016 \pm 0.00059(stat.) \pm 0.00006(syst.) \pm 0.00024(PDF)\end{aligned}$$

➤ Originally not in the plan

- Charge-dependent muon momentum scale



➤ DZero combination (2017)

$$\begin{aligned}\sin^2 \theta_{eff}^l &= 0.23095 \pm 0.00040 \\ &= 0.23095 \pm 0.00035(stat.) \pm 0.00007(syst.) \pm 0.00019(PDF)\end{aligned}$$

➤ Higher order correction

➤ Straight forward high order corrections

- ZFITTER-based form factor calculation
- ResBos vs. PYTHIA comparison

	$\Delta \sin^2 \theta_{eff}^l$
Different effective coupling for u and d quarks	+0.00008
Mass-scale dependence and complex calculation	+0.00014
Total	+0.00022

$$\Delta \sin^2 \theta_W = +0.00022 \pm 0.00004(\text{Dominated by } m_t)$$

➤ Summary

$$\begin{aligned}\sin^2 \theta_{eff}^l &= 0.23095 \pm 0.00040 \\ &= 0.23095 \pm 0.00035(stat.) \pm 0.00007(syst.) \pm 0.00019(PDF)\end{aligned}$$

- Best single experiment to date
- Best light-quark measurement

- High precision calibration
 - Both statistical uncertainty and systematic uncertainty
 - Only need mass scale at Tevatron

- Some systematic uncertainty will become more important in LHC

- Higher order correction
 - Straight forward higher order corrections
 - Uncertainty dominated by m_t