ACAT 2019



Contribution ID: 422

Type: Oral

## Charged Particle Tracking as a QUBO problem solved with Quantum Annealing-Inspired Optimization

Monday, 11 March 2019 17:10 (20 minutes)

With the upgrade of the LHC to high luminosity, an increased rate of collisions will place a higher computational burden on track reconstruction algorithms. Typical algorithms such as the Kalman Filter and Houghlike Transformation scale worse than quadratically. However, the energy function of a traditional method for tracking, the geometric Denby-Peterson (Hopfield) network method, can be described as a quadratic unconstrained binary optimization (QUBO) problem. Quantum annealers have shown shown promise in their ability to solve QUBO problems despite being NP-hard. We present a novel approach for track reconstruction by applying a quantum annealing-inspired algorithm to the Denby-Peterson method. We propose additional techniques to divide an LHC event into disjoint subgraphs in order to allow the problem to be embeddable on existing quantum annealing hardware, using multiple anneals to fit tracks to a single event. To accommodate this dimension reduction, we use Bayesian methods and further algorithms to pre- and post-process the data. Results on the TrackML dataset are presented, demonstrating the successful application of quantum annealing-inspired algorithms to the track reconstruction problem.

**Primary authors:** ZLOKAPA, Alexander (California Institute of Technology (US)); Dr VLIMANT, Jean-Roch (California Institute of Technology (US)); ANAND, Abhishek (California Institute of Technology); Prof. LIDAR, Daniel (USC); SPIROPULU, Maria (California Institute of Technology); Dr JOB, Joshua (USC)

Presenter: Dr VLIMANT, Jean-Roch (California Institute of Technology (US))

Session Classification: Track 2: Data Analysis - Algorithms and Tools

Track Classification: Track 2: Data Analysis - Algorithms and Tools