

Algorithmics of Diffraction

Mikael Mieskolainen

Diffraction and low- x 2018, Reggio Calabria, Italy

31/08/18





"A process or **set of rules** to be followed in calculations or other problem-solving operations, especially by a computer."

Ok, rules

Propagator (non-rel.) in space-time

$$\int_{\text{all paths}} \mathcal{D}x e^{iS[x(t)]}$$

Pretty cool algorithm, even if computationally hard in practise.

Basic principles:

Keep the measurement as a **fiducial measurement** (no geometrical extrapolation), minimize model \otimes data cocktails, and factorize fits out of the measurement.

What is a strict "fiducial" observable? A strict fiducial is expressed in terms of **final state** observables; pseudorapidity, transverse momentum etc.

For example, a cut on central system rapidity alone, is not, a strict fiducial cut. Think about system $y = P_t = 0$ case and arbitrary decays going outside central detector geometry. Same story for diffractive forward systems with not-measured $\xi \simeq M^2/s$ etc.

Let us slice the (pseudo)rapidity axis on N discrete intervals, count Bernoulli observables on each

Partial cross sections ($\#2^N$) \sim

$$\frac{1}{2s} \sum_M \frac{1}{\text{sym}(M)} \int_{\Omega_M} d\Pi_M \delta^{(4)} \left(p_1 + p_2 - \sum_M p_i \right) |\mathcal{M}_{2 \rightarrow M}|^2 \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}^{\otimes N} \begin{pmatrix} 1 \\ \mathcal{I}\{\Pi_M; \Xi_1\} \end{pmatrix}^{\otimes} \begin{pmatrix} 1 \\ \mathcal{I}\{\Pi_M; \Xi_2\} \end{pmatrix}^{\otimes} \cdots \otimes \begin{pmatrix} 1 \\ \mathcal{I}\{\Pi_M; \Xi_N\} \end{pmatrix},$$

where the acceptance function $\mathcal{I} : \Pi_M \rightarrow \{0, 1\}$, Π_M is a set of final state kinematical variables and Ξ_i is the i -th fiducial acceptance domain parametrization. The expression above is a 2^N -vector, essentially a multivariate polynomial expression / N -point interval correlation functional.


For some more information, see **Appendix** or: [<MM, Inverse Mathematics for QCD Diffraction, Bad Honnef, 25/09/17>](#)


A synthetic Monte Carlo example


3 rapidity slices giving us Bernoulli combinations:
 $\langle 000 \rangle, \langle 001 \rangle, \langle 010 \rangle, \dots, \langle 111 \rangle$


Particles drawn uniformly over rapidity, with fluctuating number of particles per interval $\sim \text{Poisson}(\langle N_{ch}/\Delta\eta \rangle)$ with transverse momentum $p_t \sim p_t \exp(-bp_t^2)$ (Gaussian $p_x, p_y \rightarrow$ Rayleigh p_t).

Next we vary smoothly the p_t cutoff (normalized by $\langle p_t \rangle$) for **four different particle densities per discrete rapidity interval** $\Delta\eta$, and see how event topology properties change.









A p_t -cutoff dependent "flow structure" emerges

This toy demonstration shows that without explicit p_t -cutoff descriptions (experimental characterization, detector simulations, artificial rapidity gap data unfoldings/inversion/correction), observables relying on rapidity gap structure are **not inherently stable** at all \Rightarrow needs to be taken into account in every measurement relying on rapidity gaps.

OK, the topic is known but experimental or theoretical methods to solve it are not often emphasized.

Graniitti - a new Monte Carlo event generator and analyzer for semi-exclusive *pp*-diffraction¹

To be available:

github.com/mieskolainen/graniitti, C++17, MIT license

¹Graniitti is granite in Finnish, a felsic intrusive igneous rock.

Some useful papers regarding the soft models involved:

Y. I. Azimov, V. A. Khoze, E. M. Levin, and M. G. Ryskin, Sov.J.Nucl.Phys. 21, 215 (1975)

A. Kaidalov, V. A. Khoze, A. D. Martin, and M. Ryskin, Eur. Phys. J. C21, 521-529 (2001).

V. Khoze, F. Krauss, A. Martin, M. Ryskin, and K. Zapp, Eur. Phys. J. C69, 85-93 (2010).

P. Lebiedowicz, A. Szczurek, Phys.Rev.D81:036003, (2010).

L. Harland-Lang, V. A. Khoze, M. G. Ryskin, Eur. Phys. J. C, 74(4), 2848 (2014).

⋮

Synthesis \Leftrightarrow Analysis dual

A technical tool born out of need to understand and analyze the data in detail, with maximal flexibility.

Other (exclusive) generators on market: FPMC, Superchic, Dime, GenEx, ...

Input and Processes

Compilation by `make`, **C++17** compiler + HepMC3 and LHAPDF6 required, ROOT for fit/analysis tools (optional)

♠ Steered by `.json` object description ascii files

♠ Processes:

- Pomeron-Pomeron ($\pi^+\pi^-$, K^+K^- , $p\bar{p}$, $\rho^0\rho^0$, 4π , $4K \dots$)
- Gamma-Pomeron (highly simplified)
- Gamma-Gamma (EPA) ($\ell^+\ell^-$, W^+W^- , $m\bar{m} \dots$)
- 'Durham-QCD' (main implementation done, interfacing to MadGraph color trace basis TBD)

Kinematics

♠ **Exact kinematics** for all processes: $2 \rightarrow 2$, $2 \rightarrow 3$, $2 \rightarrow 4 \dots$
and $2 \rightarrow 3 \oplus 1 \rightarrow N$ via (well known) phase space factorization relations.

♠ **Forward proton low-mass excitation**, also with exact kinematics

♠ **Fully multithreaded** (max. CPU utilization) VEGAS Monte Carlo implementation; C++ `<threads>`, `<future>`

♠ Arbitrary length decay trees according to phase space

Amplitudes

- ♠ Elastic pp -scattering via typical (single channel) **eikonalized pomeron** (Fourier-Bessel transform + exponentiation + inverse FBT), eikonalized SD and DD (triple Pomeron) at "skeleton kinematics level"
- ♠ Soft Regge central production continuum amplitudes (2,4,6 central final states) + **interfering resonances**
- ♠ Helicity amplitudes via MadGraph 5 export → standard (LO) SM + arbitrary BSM (via UFO) models




Figure: Elastic scattering test example.

Differential Screening/Absorption

♠ **Loop screening amplitude** driven by the same eikonized pomeron as used for elastic scattering. This has been 'around' since the 70's. However, only few event generators include this: QGSJet (+more complex AGK), Dime/Superchic, at least.

If the subprocess amplitude includes helicity amplitudes, each of them is screened.

♠ Numerical 2D-loop integral in \vec{k}_t -space, event-by-event. Speed vs. accuracy tradeoff. Example: screening loop $|k_t|$ and ϕ integrated separately, or via 2D-cubature polynomials (under testing)

♠ Phenomenologically of high (extreme) importance for cross section normalization and transverse plane observables, such as $\Delta\phi_{pp}$.

Spin (polarization) Systematics

- ♠ Synthesis: **Arbitrary helicity amplitudes** (Jacob-Wick style) for low-mass resonances decaying to pseudoscalar pairs, parametrized via von Neumann density matrices
- ♠ Analysis: Complete spherical harmonics expansion differentially in system (M_X, p_T) in typical Lorentz frames such as Collins-Soper, Helicity, Gottfried-Jackson ("pseudo" and forward-proton spanned)
- ♠ Analysis: Calculation of spherical moment mixing matrices ← induced by limited acceptance
- ♠ Analysis: Spin density matrix (von Neumann) fits




Figure: Continuum K^+K^- and $|\eta(K^\pm)| < 0.9$: Integrated decay angles $(\cos\theta, \phi)$ in Collins-Soper, Helicity, Lab, Gottfried-Jackson, Pseudo-Gottfried-Jackson and Non-Rotated Rest Frame.

A tool designed to study polarization
'angular momentum mixing problems' -
non-trivial, significant effects

1. Induced by limited η -acceptance of central final states
2. Induced by central system transverse momentum
3. Related with 2., using a Lorentz frame with spin-quantization (z-axis) spanned by not using forward protons \Rightarrow unknown event-by-event rotation (typical problem, measurements rarely fully exclusive)

Autofit

♠ **Fit machinery** to fit eikonal model parameters by running the generator + Non-convex minimization program (Minuit, by default) ← INPUT elastic $d\sigma(\sqrt{s})/dt$ for different \sqrt{s}

♠ Soft Pomeron central production parameters (resonance couplings) fitted by running the full MC machinery ← INPUT for example: invariant mass M_X spectrum, system $p_T(M_X)$, forward proton $\Delta\phi_{pp}(M_X)$, spherical harmonic expansions $H(M_X, p_T)$, or complete multidimensional fiducial efficiency corrected representations of data

Fast Fiducial Observables

- ♠ Ultrafast fiducial observable studies for different model scenarios via ROOT libraries (optional)
- ♠ Interface directly to RIVET via HepMC

Output and Automated testing

HepMC3 event output, and a converter to LHEF (.xml)

♠ The measured fiducial cross sections of different processes are added as "bootstrap unit tests" such that generator functionality (code non-degeneracy etc.) is guaranteed

Further Technical Developments

- ♠ 'NeuroJacobian', deep learn optimal Jacobian mapping for MC integration (instead of VEGAS dim-by-dim factorization assumption)
- ♠ Higher level of automation of feeding in process amplitudes
- ♠ Analysis techniques: statistical separation of exclusive, semi-exclusive production (system P_t dependence, exponential vs powerlaw tail), deeper understanding of spin polarization issues . . .

Conclusions

A set of new algorithmic tools developed, many of them already applied to the LHC data. Physics goes forward.

Appendix

Proton-proton diffraction

Pomeron physics.

You can think also in terms of wee partons, soft color dipoles, pomeron parton (ladder) structure etc., unfortunately there are yet no truly solid experimental constraints from the LHC data for **inclusive inelastic diffraction**. Basic Regge domain features, however, are observed in data.

Essential **fluctuating** degrees of freedom: rapidity (predominantly low- x), p_t , multiplicity and multidimensional correlations over the full range of acceptance.

\Rightarrow N -dimensional observables

Basic questions of soft diffraction

- **Unitarity**, asymptotic energy behavior of total cross sections
- Transition between "different" Pomerons: soft ... hard \rightarrow Pomeron intercept $1 + \Delta_P$ (\rightsquigarrow s evolution) and slope $\alpha'_P \sim$ "t-cone behavior" functional behavior
- $p \rightarrow N^*$ **Good-Walker spectrum** of low-mass dissociation, relativistic wavefunction and "atmosphere" of proton
- Gluonia/glueballs/soft central diffractive production
- Regge/QCD factorization properties
- Pomeron via AdS space ...

- + Correlations and fluctuations
- ⋮

Ultimately, the goal here is have a "unified" approach for interpreting the data.

Vector space view to the soft pp Diffraction

So, usually the experimental definition when talking about soft diffraction goes through large rapidity gaps $\Delta y \gtrsim 3$ and

$$\cancel{\sigma_{inel}^{pp} \equiv \sigma_{SDL} + \sigma_{SDR} + \sigma_{DD} + \sigma_{CD} + \sigma_{ND}}$$

The decomposition above is experimentally well posed only in limited phase space.


So, instead, let us start with $n = 2^N - 1$ partial cross sections

$$\sigma_{inel}^{pp} \equiv \sigma_1 + \sigma_2 + \sigma_3 + \cdots + \sigma_n, \quad (1)$$

where each subcomponent corresponds to one particular **final state topology class** over rapidity.

"Slice the (pseudo)rapidity space into N intervals"

Example: Geom.-kinem. ALICE phase-space span at Run 2
 Not all subdetectors shown ($\sim \#20$). Very good (η, p_{\perp}) coverage for diffractive physics.



One practical and important open problem:

How do you characterize (η, p_t) acceptance of forward scintillators and other low granularity counters **without** relying on MC generator ⊛ GEANT?

Applications

- ★ A **machinery for the (multi)-rapidity gap** measurements and correlation structure
- ★ A framework for **generalized studies of Regge factorization** at the LHC. Not just simplified SD,DD type, but more general
- ★ Framework to study **AGK type shadowing, and beyond**, by comparing the differential distributions within each vector combination
- ★ An attempt to **re-define the soft diffraction** observables more precisely, also introducing a hierarchy of vector observables for minbias Monte Carlo tuning
- ★ A new framework for **extracting single diffraction (SD), double diffraction (DD)** ... type component cross sections using N -dimensional Monte Carlo model "templates", which can be tuned to data

With connections to

[E. Onofri, G. Veneziano, J. Wosiek, *Commun. Math. Phys.* (2007)],
"We show how a recently proposed supersymmetric quantum mechanics model leads to non-trivial results/conjectures on the combinatorics of binary necklaces and linear-feedback shift- registers."

[H. Fu, R. Sasaki, *J. Math. Phys.* 38 (1997)], *"Following the relationship between probability distribution and coherent states, for example the well known Poisson distribution and the ordinary coherent states and relatively less known one of the binomial distribution and the $su(2)$ coherent states."*

[D. Spector, *Commun. Math. Phys.* (1990)], *"We show that the Möbius inversion function of number theory can be interpreted as the operator $(-1)^F$ in quantum field theory."*

Algebraic representations

The *probability vector* \mathbf{p} (2^N -dim), the components of *ordinary moments* m_k and the components of *central moments* δ_k below are defined using the Kronecker (tensor) products

$$\mathbf{p} = \left\langle \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}^{\otimes N} \begin{pmatrix} 1 \\ X_N \end{pmatrix} \otimes \begin{pmatrix} 1 \\ X_{N-1} \end{pmatrix} \otimes \cdots \otimes \begin{pmatrix} 1 \\ X_1 \end{pmatrix} \right\rangle$$

$$m_k = \left\langle \prod_{i=1}^N X_i^{k_i} \right\rangle = \left\langle \begin{pmatrix} 1 \\ X_N \end{pmatrix} \otimes \begin{pmatrix} 1 \\ X_{N-1} \end{pmatrix} \otimes \cdots \otimes \begin{pmatrix} 1 \\ X_1 \end{pmatrix} \right\rangle_k$$

$$\delta_k = \left\langle \prod_{i=1}^N (X_i - \langle X_i \rangle)^{k_i} \right\rangle = \left\langle \begin{pmatrix} 1 & -\langle X_N \rangle \\ X_N & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & -\langle X_{N-1} \rangle \\ X_{N-1} & 1 \end{pmatrix} \otimes \cdots \otimes \begin{pmatrix} 1 & -\langle X_1 \rangle \\ X_1 & 1 \end{pmatrix} \right\rangle_k,$$

where we use $k = 1 + \sum_{i=1}^N k_i 2^{i-1}$ (little endian binary expansion), $1 \leq k \leq 2^N$ and $k_i \in \{0, 1\}$. The central moments describe the correlations ($\# 2^N - N - 1$) between any 2 or more subspaces (rapidity slices). X_i are the corresponding random variables.

[Teugels, Jozef L. "Some representations of the multivariate Bernoulli and binomial distributions." *Journal of multivariate analysis* 32.2 (1990): 256-268.]

Diffraction analysis technique++

To summarize, we utilize different detector combinations over $\eta \rightarrow$ vector signals \rightarrow partial cross sections + multidimensional model fitting to extract σ_{SD}, σ_{DD} etc.

This latest vector space combinatorial construction goes beyond multidimensional fitting, and is compatible with discussion about multigaps, gap destruction and rescattering and short/long range y -correlations:




Figure: (a) Multigap event, (b) Gap destruction, (c) Correlation coeff. R_2
Figure from: [Khoze, Martin, Ryskin, Shuvaev, J. Phys. G: Nucl. Part. Phys. 36 (2009) 093001]

AGK Cutting Rules

Field theory Combinatorics

The total cross section for exchange of μ Pomerons, σ_μ^{tot} , partial cross section $\sigma_\mu^{(\nu)}$ of a final state with a number of ν cut Pomerons and their ratio

$$\frac{\sigma_\mu^{(\nu)}}{\sigma_\mu^{tot}} = (-1)^{\mu-\nu} \frac{\mu!}{\nu!(\mu-\nu)!} (2^{\mu-1} - \delta_{0\nu}), \quad (2)$$

[Abramovski, Gribov, Kancheli, Sov. J. Nucl. Phys. 18, 308 (1974)], [E. Levin, hep-ph/9503399]

$\mu \setminus \nu$	0	1	2	3	4	5	6
1	0	1	0	0	0	0	0
2	1	-4	2	0	0	0	0
3	-3	12	-12	4	0	0	0
4	7	-32	48	-32	8	0	0
5	-15	80	-160	160	-80	16	0
6	31	-192	480	-640	480	-192	32

Table: AGK factors for $\mu = 1, 2, \dots, 6$ exchanged Pomerons. Summing over μ requires some explicit (Regge/Eikonal etc.) model in addition to these.

"Super-Eikonals"

Combinatorial (de)-compounding or pileup inversion

Poisson \otimes Multinomial Vector Model

$$\begin{aligned}\hat{y}_i &= \frac{1}{1 - e^{-\mu}} \sum_{k=1}^{\infty} \frac{\mu^k}{k!} e^{-\mu} W_{ik}, \quad i = 1, \dots, 2^N - 1 = n \\ &= \frac{e^{-\mu}}{1 - e^{-\mu}} \sum_{k=1}^{\infty} \frac{\mu^k}{k!} \left\{ \sum_{\Omega_{ik}} \frac{k!}{\prod_{j=1}^n x_j!} \prod_{j=1}^n p_j^{x_j} \right\}\end{aligned}\quad (3)$$

The multinomial term and its values of $x_j \in \mathbb{N}$ are evaluated over all valid combinations for probabilities y_i from the set of n -tuples Ω_{ik} , that is, those which are allowed by poset combinatorics:

$$\Omega_{ik} = \left\{ (x_1, \dots, x_j, \dots, x_n) \mid \bigvee_j x_j \mathbf{c}_j = \mathbf{c}_i \text{ and } \sum_j x_j = k \right\}, \quad (4)$$

where \bigvee operator takes care of "summing" the binary vectors \mathbf{c}_j of multiplicity x_j and thus evaluating the "pileup" compositions.

The idea in a nutshell: **We measure probabilities \mathbf{y} , and want to solve \mathbf{p}**

Solution based on the principle of inclusion-exclusion

General math framework: Incidence algebras [Gian-Carlo Rota, MIT, 60's]

The principle of inclusion-exclusion is the Möbius inversion for subsets. Now let different rapidity slices and their signals be represented with subsets $D_1, D_2, \dots, D_N \subset D$. Then

$$P\left(\bigcup_{i=1}^N D_i\right) = \sum_{k=1}^N \left((-1)^{k-1} \sum_{I \subset \{1, \dots, N\}, |I|=k} P(D_I) \right). \quad (5)$$

One can wrap that thing above into a matrix. Notice the $(-1)^{k-1}$ factor, that gives the essential structure.

Uniform (max entropy) input $\mathbf{p} = \mathbf{1}$ case, $N = 3$




Figure: A solution. On x-axis the Poisson μ and on y-axis the components of the vector \mathbf{y} .

Starting with very elementary definitions, interesting distributions emerge from combinatorics.

Alternating sign inverse solution for $N = 3$

$$\mathbf{p} = \frac{1}{\mu} \begin{pmatrix} \ln(e_{-}^{\mu} y_1 + 1) \\ \ln(e_{-}^{\mu} y_2 + 1) \\ - \sum_{c=1,2} \ln(e_{-}^{\mu} y_c + 1) + \ln(1 + \sum_{c=1,2,3} e_{-}^{\mu} y_c) \\ \ln(e_{-}^{\mu} y_4 + 1) \\ - \sum_{c=1,4} \ln(e_{-}^{\mu} y_c + 1) + \ln(1 + \sum_{c=1,4,5} e_{-}^{\mu} y_c) \\ - \sum_{c=2,4} \ln(e_{-}^{\mu} y_c + 1) + \ln(1 + \sum_{c=2,4,6} e_{-}^{\mu} y_c) \\ \mu + \sum_{c=1,2,4} \ln(e_{-}^{\mu} y_c + 1) - \ln(1 + \sum_{c=1,2,3} e_{-}^{\mu} y_c) \quad \dots \\ - \ln(1 + \sum_{c=1,4,5} e_{-}^{\mu} y_c) - \ln(1 + \sum_{c=2,4,6} e_{-}^{\mu} y_c) \end{pmatrix},$$

where by conservation of probability we chose to fix $y_7 = 1 - \sum_{c=1}^6 y_c$
and for saving ink we set $e_{-}^{\mu} \equiv e^{\mu} - 1$.




Figure: Poisson model \otimes Dirichlet distribution drawn probabilities as a statistical mixing operator (matrix) $S : \mathbf{p} \mapsto \mathbf{y}$, $N = 6$. Fractal structure, due to the Boolean vector space, is the Sierpinski triangle. (Dark blue = 0 ... Yellow = 1)




Figure: Hidden polynomial structure, $N = 8$.

Short summary

The vector space measurement model allows a mathematically self consistent way to do combinatorial analysis of soft diffraction, plus also to **extract** $\sigma_{SD}, \sigma_{DD}, \sigma_{ND}$ etc. via multidimensional Bayesian/Frequentist fitting (given the MC model).

AGK cutting rules can be incorporated into the **combinatorics inversion** framework. Leading the way to completely new analyses of, e.g., gap survival $S^2(\Omega)$ discussion. This framework works directly for pile-up inversion of gap topologies (multiple pp interactions per bunch crossing).

The vector space itself can be studied in the context of kinematics, diffraction models and Regge theory, together with tools from combinatorics and algebraic geometry (technically the structure is **Grassmannian**).

Recursive Inverse of Stochastic Autoconvolution

The first solution with fully non-linear uncertainty estimation



Recursion, M.C. Escher

The problem?

Think about having a superposition of final state multiplicities (= **autoconvolution**²), let's say, in proton-proton collisions

Main problem is limited statistics in steeply falling tails → huge oscillations, naive (textbook³) solutions fail miserably

²sum of random variables is equivalent to a convolution of their densities

³inverting stochastic autoconvolution is not usual textbook material




Figure: Poissonian superposition with different Poisson mean values μ , with x a random variable \sim Negative Binomial Distribution.

Forward problem

The autoconvoluted distribution of $Y \sim g_Y$ is now written formally as a Poisson probabilities weighted infinite series⁴

$$\begin{aligned} g_Y(y) &= P_1 f_X(y) + P_2 [f_X \circledast f_X](y) + P_3 [[f_X \circledast f_X] \circledast f_X](y) + \dots \\ &= \frac{1}{1 - e^{-\mu}} \sum_{K=1}^{\infty} \frac{\mu^K}{K!} e^{-\mu} f_X^{\circledast K}(y) \end{aligned} \quad (6)$$

where the *convolution power* \circledast^K is defined recursively as $f^{\circledast K} = f^{\circledast(K-1)} \circledast f$ and $f^{\circledast 1} = f$.

We do need not to limit ourself to the Poisson compound sum, but take that as an example

⁴We have removed the unobservable case $K = 0$ which gives $Y = 0$ and renormalized the remaining Poisson probabilities $P_K, K = 1, 2, 3, \dots$ to sum to one.

A spectral solution to the forward problem via the characteristic function

In the spectral domain, the characteristic function (CHF) φ_X is defined as

$$\varphi_X(t) = \mathbb{E}[e^{itX}] = \int_{\mathbb{R}} e^{itX} f_X(x) dx \quad (7)$$

and for the compound Poisson case you end up with

$$\varphi_g(t) \equiv \varphi_{Y|K>0}(t) = \frac{e^{-\mu} (e^{\mu\varphi_f(t)} - 1)}{1 - e^{-\mu}} = \frac{1}{e^{\mu} - 1} (e^{\mu\varphi_f(t)} - 1).$$

The main thing is that you want to find out $\varphi_f(t)$.

Inverse solution in a nutshell

To find out $\hat{f}(x)$, **use recursion**. First estimate $\hat{f}^0 = g(x)$.

Take Fast Fourier Transform (**FFT**) of $\hat{f}^k(x)$ to get $\hat{\varphi}_f^k(t)$, use the spectral map to get $\hat{\varphi}_g(t)$ and construct corresponding AC operator, take IFFT of AC operator, map $g(x) \rightarrow \hat{f}^{k+1}(x)$ in original domain with **Max Entropy** inversion + regularization, use **Efron's statistical Bootstrap** to estimate uncertainty, and add one so-called bias subtraction iteration around it:

"Bias subtraction" \curvearrowright

"Daughter Bootstrap" \curvearrowright

Fast Fourier Transform & Max Entropy recursion \circlearrowright




Figure: Inverse solution with algorithmic uncertainty estimation (blue band 95CL).




Figure: Data driven regularization parameter λ selection as an equilibrium between "backprojection" error $\chi_{\hat{g}}^2$ and smoothness $\|\nabla_x \hat{f} / \hat{f}\|^2$.

DeepEfficiency - Deep neural network⁵ based optimal algorithm for performing multidimensional detector efficiency inversion

Code to be available: github.com/mieskolainen/deepefficiency, MIT license

⁵TensorFlow driven, which is a "symbolic math" library from Google Brain.

Essentially, a 'Detector Matrix' in Higher Dimensions - utilizing all observable degrees of freedom

For what?

- Maximally MC generator independent efficiency corrections of, for example, two-body low mass central systems ($\pi^+\pi^-$, K^+K^- , $p\bar{p}$...). Correct all observables simultaneously!

Cannot I just divide histograms and correct for efficiency by that way, or construct some unfolding matrix observable by observable?

- Yes, if your physics is well simulated (say \sim QED process) and/or you're happy with MC generator biased results.

Work in progress