

# The CMS Pixel Detector for the High Luminosity LHC

*Tuesday 19 February 2019 17:20 (20 minutes)*

The High Luminosity Large Hadron Collider (HL-LHC) at CERN is expected to collide protons at a centre-of-mass energy of 14 TeV and to reach the unprecedented peak instantaneous luminosity of  $5-7.5 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$  with an average number of pileup events of 140-200. This will allow the ATLAS and CMS experiments to collect integrated luminosities up to 3000-4500  $\text{fb}^{-1}$  during the project lifetime. To cope with this extreme scenario the CMS detector will be substantially upgraded before starting the HL-LHC, a plan known as CMS Phase-2 upgrade. The entire CMS silicon pixel detector will be replaced and the new detector will feature increased radiation hardness, higher granularity and capability to handle higher data rate and longer trigger latency. In this talk the Phase-2 upgrade of the CMS silicon pixel detector will be reviewed, focusing on the features of the detector layout and on developments of new pixel devices.

**Authors:** SGUAZZONI, Giacomo (INFN (IT)); CMS TRACKER COLLABORATION

**Presenter:** SGUAZZONI, Giacomo (INFN (IT))

**Session Classification:** Semiconductor Detectors

**Track Classification:** Semiconductor Detectors