


Towards EeV Astronomy

catching the sources of ultra-high-energy cosmic rays

Exciting times!



And we still don't know the origin of UHECRs

A UHECR journey

Source?

- particle injection?
- acceleration? shocks?
- reconnection?...

Outflow

- structure?
- B?
- size?

Cosmic backgrounds

interactions on CMB, UV/opt/IR photons

cosmogenic neutrino and gamma-ray production

Intergalactic magnetic fields

magnetic deflection
temporal & angular spread/shifts

Backgrounds

- radiative? baryonic?
- evolution, density?
- magnetic field: deflections?

associated neutrino and gamma-ray production

Observables

UHECR

- mass
- spectrum
- anisotropy

neutrinos


- flavors
- spectrum
- anisotropy
- time variabilities

multi-wavelength photons


- spectral features
- time variabilities
- angular spread
- source distribution

GW

- spectrum
- arrival directions
- time



Current multi-messenger data: useful to understand UHECRs?



Cosmic backgrounds

interactions on CMB, UV/opt/
IR photons

*cosmogenic neutrino and
gamma-ray production*

Backgrounds

- radiative? baryonic?
- evolution, density?
- magnetic field: deflections?

*associated neutrino and
gamma-ray production*

Secondaries take up 5-10% of parent cosmic-ray energy

$$E_\nu \sim 5\% E_{\text{CR}}/A$$


$$E_\gamma \sim 10\% E_{\text{CR}}/A$$

$$E_{\text{CR}} > 10^{18} \text{ eV}$$

$$E_\nu > 10^{16} \text{ eV}$$


IceCube neutrinos do not directly probe UHECRs

Actually, none of the current multi-messenger data
(except UHECR data) can directly probe UHECRs
... but they help :-)




Learning from UHECR data

Energy spectrum



Verzi et al. (2017)


Arrival directions



$E > 10^{19}$ eV

Auger & TA combined analysis

Aab et al. (2014)



Auger Coll. ICRC 2017

Mass composition


most UHECRs have
rigidity $E/eZ \sim 3\text{-}10$ EV

deflections depend on rigidity


UHECRs and intergalactic magnetic fields

KK & Olinto 2011

few
observations
large
uncertainties!



Alves Batista et al. 2017



cumulative filling factors

salient structures of the IGMF


(scattering centers) determine the
deflection of UHECRs

$$\alpha \sim 2^\circ Z \left(\frac{E}{60 \text{ EeV}} \right)^{-1} \left(\frac{\tau}{3} \right)^{1/2} \left(\frac{r_i}{2 \text{ Mpc}} \right)^{1/2} \left(\frac{B_i}{10 \text{ nG}} \right) \left(\frac{\lambda_i}{0.1 \text{ Mpc}} \right)^{1/2}$$

number size strength coherence
length

KK & Lemoine 2008b

Galactic and Intergalactic magnetic fields



BLUR (+shift) from IGMF

blur: controlled by statistics


IMAGINE

Interstellar MAGnetic field INference Engine
arXiv: 1805.02496v1

a publicly available Bayesian platform that employs robust statistical methods to explore the multi-dimensional likelihood space using any number of modular inputs

GMF IGMF \longleftrightarrow UHECR arrival directions

Mollerach & Roulet 2017



SHIFTS (+blur) from GMF

shifts: knowledge on GMF will help

What can we do with
rigidities $E/eZ \sim 10$ EV
and deflections $\sim 10^\circ$?


Learning from large scale anisotropies

Galactic or extragalactic?

UHECR source(s) in our Galaxy imply high level of dipole amplitude


esp. for light mass composition @ 8 EeV

e.g., Calvez et al. 2010
Giacinti et al. 2011
Eichler et al. 2016



Auger Coll. Science 2017
Ahlers (2018) 1805.08220v1

di Matteo & Tinyakov 2018




dipole expected from LSS

also Globus & Piran 2018 —> dipole from LSS 5% @[4-8] EeV

Auger confirms that UHECRs are of extragalactic origin

5.2 sigma dipole of 6.5% observed at $E > 8$ EeV

dipole direction, amplitude and light composition at EeV energies in tension with source inside Galaxy



Learning from intermediate scale anisotropies


population of sources?

proton fraction

intermediate scale anisotropy
should be detected @ $E > 50$ EeV
with proton fraction $> 50\%$
and > 2000 events

Oikonomou, KK, Abdalla 2014

Rouillé d'Orfeuil et al. 2014




Auger reports anisotropy compatible
with starburst galaxy distribution

> 5000 events above 20 EeV
starburst model fits the data better than the hypothesis
of isotropy with a stat. significance of 4.0σ


There is a sizable fraction of particles
with rigidity $E/eZ > 10$ EV!

Auger Coll. arXiv:1801.06160




Learning from UHECR data

Energy spectrum



Verzi et al. (2017)


Arrival directions



$E > 10^{19} \text{ eV}$

Auger & TA combined analysis

Aab et al. (2014)




Auger Coll. ICRC 2017

Mass composition

a combined fit to the data?

Information from UHECR spectra and composition



UHECR parameters

- A flux normalisation
- α injection spectral index in $E^{-\alpha}$
- R_{\max} (max. rigidity \sim max. proton energy)
- composition
- source evolution e.g., SFR/AGN or in $(1+z)^m$

Alves Batista, de Almeida, Lago, KK, submitted

- if emissivity evolution free parameter \rightarrow best fit $m = -1.5$
- Negative source evolution:
 - e.g., tidal disruption events
 - cosmic variance local dominant of sources
- very hard spectral indices difficult to reconcile with most particle acceleration models. $\alpha > \sim 1$ favored in theory.

phenomenologically
reasonable models with
good deviances





Table 1. Best-fit parameters for specific spectral indices.

m	α	$\log(R_{\max}/\text{V})$	f_p	f_{He}	f_N	f_{Si}	f_{Fe}	D
-1.5	+1.00	18.7	0.0003	0.0002	0.8867	0.1128	0.0000	1.46
SFR	+0.80	18.6	0.0764	0.1802	0.6652	0.0781	0.0001	1.63
AGN	+0.80	18.6	0.1687	0.1488	0.6116	0.0709	0.0000	1.59
GRB	+0.80	18.6	0.1362	0.1842	0.6059	0.0738	0.0000	1.60

Learning from secondary neutrinos?

Alves Batista, de Almeida, Lago, KK, submitted
 GRAND Science & Design, in prep
 KK, Allard, Olinto 2010
 Van Vliet et al. arXiv:1707.04511



standard

99% C.L. fit to Auger among generic SFR, AGN, GRB source evolutions

pessimistic

90% C.L. fit to Auger data with $(1+z)^m$ source evolution with $m < 0$

most pessimistic!

adding IGMF \rightarrow harder $\alpha \rightarrow$ increases neutrino flux
 alleviating simplifying assumption \rightarrow increases neutrino flux

low rigidities

$R_{\max} \sim 10^{18.1-18.8} \text{ V}$

R_{\max}

below or above pion prod. threshold

very hard

$-1.5 < \alpha < +1.2$

spectral index

flux of secondary protons
 $E^{-\alpha}$

source evolution history

composition

intermediate dominated

p 8%, He 18%, N 67%,
 Fe 0.01%

UHECR flux normalisation

Learning from multi-wavelength observations

luminosity budget

condition for acceleration

$$t_{\text{acc}} \lesssim t_{\text{dyn}}$$

$$t_{\text{acc}} = \mathcal{A} t_L$$


depends on
acc. mechanism
and environment

$$\mathcal{A} \gg 1$$

$$\mathcal{A} \sim 1 \text{ at best}$$

Larmor
time

$$t_{\text{dyn}} \sim R / \beta_w \Gamma_w c$$



outflow magnetic luminosity

$$L_B \equiv \Gamma_w R^2 B^2 / 2 > 10^{45} Z^{-2} E_{20}^2 \text{ erg s}^{-1}$$

lower bound of the bolometric luminosity of source

Lemoine & Waxman 2009


Learning from multi-wavelength observations + UHECR anisotropy

source bolometric luminosity $> 10^{45} Z^{-2} E_{20}^2 \text{ erg s}^{-1}$ *Lemoine & Waxman 2009*

level of clustering in the sky in Auger data

➤ apparent number density of sources @ given energy and angular deflection α *Abreu et al. 2013*

UHECRs cannot be dominantly protons from steady sources



Fang & KK, 2016


possibilities for
heavy/intermediate composition
from steady sources e.g.,
BH jets: *Fang & Murase 2016*
radio galaxies: *Eichmann et al. 2018*

Learning from multi-wavelength observations

luminosity budget

source bolometric luminosity $> 10^{45} Z^{-2} E_{20}^2 \text{ erg s}^{-1}$ *Lemoine & Waxman 2009*

many transient sources could make it *Guépin & KK 2016*



"Hillas plot for transients"

Learning from secondary neutrinos?

a general criterion for transients

Guépin & KK 2016

transient source

observed

- photon flux Φ
- bolometric luminosity L_{bol}
- time variability t_{var}

choose

- Lorentz factor Γ



setting all parameters to **worst case scenario** for neutrino production

experimental
detection
limit

baryon loading
to be chosen
(theoretically ok
+ conservative)

minimal photon flux required for detection $\Phi_{\gamma, \min} = \frac{8}{3} \frac{4\pi\beta^2 c^2 \Gamma^4 \mathcal{S}_{\text{exp}}}{\langle \sigma_{p\gamma} \kappa_{p\gamma} \rangle} \eta_p^{-1} L_{\text{bol}}^{-1} (1+z)^{-1}$

$\simeq 2 \text{ Jy } \eta_p^{-1} \Gamma^4 L_{\text{bol}, 52}^{-1} (1+z)^{-1} .$



Neutrino flares and TXS 0506+56

Guépin & KK 2016

photon flux **needed**
for detection with GRAND


Class	$E_{\nu, \max}$ (GeV)	ϵ_{γ} (eV)	$\eta_p \Phi_{\gamma, \min}$ (ph cm ⁻² s ⁻¹)	$D_{L, \max}$ [z_{\max}]
Blazar flares	10^{10}	0.1	10^3	[1.2]
LL GRBs*	10^9	0.1	10^3	18 Mpc
TDEs	10^9	10^4	10^3	25 Mpc
SLSNe	10^9	10^{-3}	10^2	7.9 Mpc
SNe*	10^9	10^{-2}	10^4	79 kpc

$L_{\text{pk}} = 1.7 \times 10^{44}$ erg/s
 $t_{\text{rise}} = 3 \times 10^5$ s,
 $D = 66$ Mpc.

Si on considère le ca
 $\beta^{-2} \eta_p \Phi_{\gamma}$
cm⁻² s⁻¹.

Pour cette source Φ_{γ}
80 ph cm⁻² s⁻¹.


side view




IC170922A:
290 TeV neutrino

*IceCube Coll.
Science (2018)*

13 ± 5 above the background of atmospheric neutrinos, 3.5σ




top view



Neutrino flares and TXS 0506+56


10 public alerts and 41 archival events
Post-trials p-value for association: 3.0σ

IceCube Collaboration, Fermi-LAT, MAGIC, AGILE, ASAS-SN, HAWC, H.E.S.S., INTEGRAL, et al., Science (2018)



criterion from Guépin & KK 2016
Blazar flare from **TXS 0506+056**
coincident with IceCube 170922A?

$\phi_{\min} > 10^3 \text{ ph cm}^{-2} \text{ s}^{-1}$
 $\phi_{\text{obs}}(1 \text{ eV}) \sim 10^2 \text{ ph cm}^{-2} \text{ s}^{-1}$
source not excluded!




$z \sim 0.337$ *Paiano et al. 2018*


Candidate Neutrino Source: TXS 0506+056

Keivani et al. 2018

leptonic model



hadronic model



leptonic with radiatively subdominant hadronic component

detection proba. with IC in real time during 6-month flare = 1-2%

lucky?

but the 2014-2015 neutrino flares require higher rates

($L \sim 10^{47}$ erg/s over 158 days $\sim 4 \times$ average gamma-ray luminosity)

cascade implies high X-ray level to match observed neutrino flux

Gao et al. 2018

Cerruti et al. 2018

Zhang, Fang & Li 2018

Gokus et al. 2018

Sahakyan 2018

Multi-zone or more complicated models?

- Additional photomeson production by external radiation fields
 - hadronuclear production (e.g., jet-cloud interaction)
- More parameters introduced, the setup is ad-hoc


Murase,
Oikonomou,
Petropoulou 2018

Connection with GW observations

NS mergers as producers of UHECR and neutrinos

Decoene, Fang, Guépin, KK, Metzger, in prep.

Rosswog 2013





Particle acceleration:
disk wind? corona? wind/ejecta interface?

Kimura et al. 2015 (accretion disk)

Levinson 2000 (electric field)


Riquelme et al. 2012, Hoshino 2013, 2015 (magnetic reconnection)

Lynn et al. 2014 (stochastic)




Catching the sources of UHECRs

real-time EeV multi-messenger astronomy is the way




all connected after all
with EeV neutrinos as a
principal ingredient




Astrophysical Multimessenger Observatory Network

Proposal for
Institute for Multi-Messenger Astrophysics
arXiv:1807.04780



Can we hope to detect very high-energy neutrino sources?


Neutrinos don't have a horizon: won't we be polluted by background neutrinos?



boxes for experiments assuming neutrino flux: $10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1}$

YES if

- ▶ good angular resolution ($<$ fraction of degree)
- ▶ number of detected events > 100 s




What we can aim to do with future observatories

cosmogenic:
guaranteed

direct from source:
likely more abundant

pessimistic scenarios
of cosmogenic neutrinos = good!

low background for source neutrinos
talk by Heinze Tuesday PM




detect the
first EeV
neutrinos


detect EeV neutrino **point sources**

100s of events
~0.3° angular resolution

detect **cosmogenic** neutrinos





GRAND



<http://grand.cnrs.fr/>

The Giant Radio Array for Neutrino Detection







The GRAND Concept

radio detection: a mature and autonomous technique
AERA, LOFAR, CODALEMA/EXTASIS, Tunka-Rex, TREND

radio antennas cheap and robust: ideal for giant arrays

Inclined showers with mountain targets







The GRAND Concept

200,000 radio antennas over 200,000 km²
~20 hotspots of 10k antennas
in favorable locations in China & around the world

- ✓ Radio environment: radio quiet
- ✓ Physical environment: mountains
- ✓ Access
- ✓ Installation and Maintenance
- ✓ Other issues (e.g., political)



✱ A staged approach with self-standing pathfinders







GRANDProto300: a self-standing pathfinder

Site: 8 sites surveyed in China, 6 with excellent electromagnetic conditions


Layout: 300 antennas, 200km², 1km step size with denser infield
→ Erange = 10^{16.5}-10¹⁸eV



HorizonAntenna, successfully tested in the field (August 2018)

Electronics:

50-200MHz analog filtering,
500MSPS sampling
FPGA+CPU
Bullet WiFi data transfert



500MSPS
Quad ADC
2x 15GS/s serial output

GRAND Technical Challenges


- How to collect data?
 - Optimised trigger (machine learning (?), see Führer et al. ARENA2018) to improve selection @ antenna level
 - Optimised informations to be transmitted to central DAQ
- How to identify air showers out of the ultra dominant background ?
 - Specific signatures of air shower radio signals vs background transients demonstrated (TREND offline selection algorithm: 1 event out 10^8 pass & final sample background contamination $< 20\%$)
 - Improved setup (GRANDproto35, being deployed) should lead to even better performances
 - Deep learning techniques
- How well can we reconstruct the primary particle information
 - Simulations promising (similar performances as for standard showers) + deep learning technique

Need for an experimental setup to test and optimize techniques




- How to deploy and run 200,000 units over 200,000km²?
- How much will it cost? Who will pay for it?

go for industrial approach!
answers to be studied at
later stage




Simulated performances


exposure for Hotspot1
10'000 km²



~0.1-0.3° angular resolution for GP300
also achievable for Hotspot1




V. Decoene (IAP)



GRAND full sensitivity ($E > 10^{17}$ eV)
~4x10⁻¹⁰ GeV cm⁻² s⁻¹ sr⁻¹

X_{max} resolution:
< 40 g/cm² achievable for
 $E > 10^{19}$ eV
with GP300 & further stages


C. Guépin (IAP)



A rich science case

UHECR, hadronic physics

- 20 times the exposure of Auger!
- GRANDProto300: transition from Galactic/extragalactic
- hadronic physics: muon discrepancy, UHECR mass composition, p-air cross-section



neutrino physics

- neutrino cross-section measurements
- spectral, angular distortions
- flavor ratios

UHE gamma rays

- competitive with Auger at GRANDProto300 stage

radio-astronomy in a novel way

- unphased integration of signals: an almost full-sky survey of radio signals
- can detect FRBs and Giant Radio pulses of the Crab already at the GRANDProto300 stage

Jaime Álvarez-Muñiz¹, Rafael Alves Batista^{2,3}, Aswathi Balagopal V.⁴, Julien Bolmont⁵, Mauricio Bustamante^{6,7,8,†},
Washington Carvalho Jr.⁹, Didier Charrier¹⁰, Ismaël Cognard^{11,12}, Valentin Decoene¹³, Peter B. Denton⁶,
Sijbrand De Jong^{14,15}, Krijn D. De Vries¹⁶, Ralph Engel¹⁷, Ke Fang^{18,19,20}, Chad Finley^{21,22}, Stefano Gabici²³,
QuanBu Gou²⁴, Junhua Gu²⁵, Claire Guépin¹³, Hongbo Hu²⁴, Yan Huang²⁵, Kumiko Kotera^{13,*}, Sandra Le Coz²⁵,
Jean-Philippe Lenain⁵, Guoliang Lü²⁶, Olivier Martineau-Huynh^{5,25,*}, Miguel Mostafá^{27,28,29}, Fabrice Mottez³⁰,
Kohta Murase^{27,28,29}, Valentin Niess³¹, Foteini Oikonomou^{32,27,28,29}, Tanguy Pierog¹⁷, Xiangli Qian³³, Bo Qin²⁵,
Duan Ran²⁵, Nicolas Renault-Tinacci¹³, Markus Roth¹⁷, Frank G. Schröder^{34,17}, Fabian Schüssler³⁵, Cyril Tasse³⁶,
Charles Timmermans^{14,15}, Matías Tueros³⁷, Xiangping Wu^{38,25,*}, Philippe Zarka³⁹, Andreas Zech³⁰,
B. Theodore Zhang^{40,41}, Jianli Zhang²⁵, Yi Zhang²⁴, Qian Zheng^{42,24}, Anne Zilles¹³

~50 collaborators from 10 countries

*France (15), China (7), USA (6), Netherlands (2), Germany (2),
Copenhagen (2), Spain (2), Brazil (2), Belgium, Argentina, Sweden*



GRAND Workshop,
IAP, August 2018

electronics: Nikhef/Radboud U.

antenna design: Subatech (design),
Electronics University of XiAn (production)

simulations: IAP, VUB

particle detectors: Penn State U.


computing resources: KIT




Giant Radio Array for Neutrino Detection

<https://arxiv.org/abs/1810.09994>

Science and Design
White Paper



<http://grand.cnrs.fr/>



join us and bring your ideas!

<http://grand.cnrs.fr/>