

Overview of the MIXMAX Project

George Savvidy

Demokritos National Research Center
Athens, Greece

MIXMAX Consortium Meeting

3-4 July 2018
Athens

Publications and defended PhDs

Implementation of the MIXMAX random number generators

$A(N,s)$ and $A(N,s,m)$ Family of MIXMAX operators, C++ Code

Spectrum, Entropy and Spectral test of the MIXMAX Generator

1. G. Savvidy and Natalia Savvidi,
On the Monte Carlo simulation of physical systems
J.Comput.Phys. **97** (1991) 566; Preprint EFI, 1986
2. K. Savvidy, The MIXMAX random number generator
Comput.Phys.Commun. **196** (2015) 161
3. G. Savvidy, Anosov C-systems and Random Number Generators
Theor.Math.Phys. 2016; **arXiv:1507.06348**
4. K. Savvidy and G. Savvidy
Spectrum and Entropy of C-systems. MIXMAX Generator,
Chaos, Solitons and Fractals, **91** (2016) 33-38
5. G. Savvidy and K. Savvidy,
Hyperbolic Anosov C-systems.
Exponential Decay of Correlation Functions,
Chaos Solitons Fractals **107** (2018) 244-250; **arXiv:1702.03574**

- 6.N.Martirosyan, K.Savvidy and George Savvidy,
Spectral Test of the MIXMAX Random Number Generator
ArXiv **1806.05243** (2018)
7. S.Konitopoulos and K.Savvidy,
A Priori tests for the MIXMAX random number generator
ArXiv **1804.01563** (2018)
- 8.A.Poghosyan, H.Babujian and G. Savvidy,
Artin Billiard, Exponential Decay of Correlation Functions
ArXiv **1802.09998** (2018)
- 9.H.Poghosyan, K.Savvidy and G. Savvidy,
Classical Limit Theorems and high entropy
MIXMAX random number generator
ArXiv **1708.04129** (2018)

Defended PhD thesis's

1. Narek Martirosyan 2017
2. Asmik Poghosyan 2017
3. Hayk Poghosyan 2018

1. HEPFORGE.ORG, <http://mixmax.hepforge.org>;
<http://www.inp.demokritos.gr/savvidy/mixmax.php>
2. CLHEP/Geant4, Release 2.3.1.1, on November 10th, 2015
<http://proj-clhep.web.cern.ch/proj-clhep/>
3. PYTHIA8
http://home.thep.lu.se/~torbjorn/doxygen/MixMax_8h_source.html
4. CMS Experiment
<https://home.cern/about/experiments/cms>
5. ROOT, Release 6.04/06 on 2015-10-13,
https://root.cern.ch/doc/master/mixmax_8h_source.html
6. GSL-GNU Scientific Library,
<https://www.gnu.org/software/gsl/>

Family of operators $A(N,s)$ parametrised by the integers N and s



$$A(N, s) = \begin{pmatrix} 1 & 1 & 1 & 1 & \dots & 1 & 1 \\ 1 & 2 & 1 & 1 & \dots & 1 & 1 \\ 1 & 3+s & 2 & 1 & \dots & 1 & 1 \\ 1 & 4 & 3 & 2 & \dots & 1 & 1 \\ & & & \dots & & & \\ 1 & N & N-1 & N-2 & \dots & 3 & 2 \end{pmatrix} \quad (1)$$

The matrix is of the size $N \times N$

Its entries are all integers $A_{ij} \in \mathbb{Z}$

Det $A = 1$

The spectrum and the value of the Kolmogorov entropy?



Eigenvalue Distribution of $A(N,s)$ and of $A^{-1}(N,s)$
all of them are lying outside of the unit circle

Size N	Magic s	Entropy	Period $\approx \log_{10}(q)$
256	-1	164.5	4682
256	487013230256099064	193.6	4682

Table: Properties of operators A(N,s) for different special s .

Size N	Magic s	Entropy (lower bound)	Period $\approx \log_{10}(q)$
7307	0	4502.1	134158
20693	0	12749.5	379963
25087	0	15456.9	460649
28883	1	17795.7	530355
40045	-3	24673.0	735321
44851	-3	27634.1	823572

Table: Table of properties of the operator $A(N,s)$ for large matrix size N . The third column is the value of the Kolmogorov entropy. All these generators pass tests in the BigCrush suite. For the largest of them the period approaches a *million digits*.


$A(N,s,m)$

A three-parameter family of C-operators $A(N, s, m)$, where m is some integer:

$$\begin{pmatrix} 1 & 1 & 1 & 1 & \dots & 1 & 1 \\ 1 & 2 & 1 & 1 & \dots & 1 & 1 \\ 1 & m+2+s & 2 & 1 & \dots & 1 & 1 \\ 1 & 2m+2 & m+2 & 2 & \dots & 1 & 1 \\ 1 & 3m+2 & 2m+2 & m+2 & \dots & 1 & 1 \\ & & & \dots & & & \\ 1 & (N-2)m+2 & (N-3)m+2 & (N-4)m+2 & \dots & m+2 & 2 \end{pmatrix}$$

Size N	Magic m	Magic s	Entropy	Period $\approx \log_{10}(q)$
8	$m = 2^{53} + 1$	s=0	220.4	129
17	$m = 2^{36} + 1$	s=0	374.3	294
40	$m = 2^{42} + 1$	s=0	1106.3	716
60	$m = 2^{52} + 1$	s=0	2090.5	1083
96	$m = 2^{55} + 1$	s=0	3583.6	1745
120	$m = 2^{51} + 1$	s=1	4171.4	2185
240	$m = 2^{51} + 1$	s=487013230256099140	8418.8	4389

Table: Table of three-parameter MIXMAX generators A(N,s,m). These generators have an advantage of having a very high quality sequence for moderate and small N . In particular, the smallest generator we tested, $N = 8$, passes all tests in the BigCrush suite.



The distribution of the eigenvalues of the $A(N, s, m)$ for
 $N = 240, s = 487013230256099140, m = 2^{51} + 1$.

mixmax realese 110 *speed = 4 · 10⁻⁹second*
mixmax.c
mixmax.h
driver main.c
driver verification.c
mixmax skip N240.c
mixmax skip N17.c
-- -- -- > README.pdf

mixmax gsl driver.c

driver testU01.c

Dmitri Anosov, in his fundamental work on *hyperbolic dynamical C-systems* pointed out that the basic property of the geodesic flow on closed Riemannian manifolds V^n of negative curvature is a *uniform instability of all its trajectories*.

In physical terms that means that *in the neighbourhood of every fixed trajectory the trajectories behave similarly to the trajectories in the neighbourhood of a saddle point*.

The hyperbolic instability of the dynamical system $\{T^t\}$ which is defined by the equations ($w \in W^m$)


$$\dot{w} = f(w) \quad (2)$$

takes place for all solutions $\delta w \equiv \omega$ of the deviation equation

$$\dot{\omega} = \left. \frac{\partial f}{\partial w} \right|_{w(t)=T^t w} \omega \quad (3)$$

in the neighbourhood of each phase trajectory $w(t) = T^t w$.

The Anosov C-systems are genuine hyperbolic systems



the behaviour of all nearby trajectories is exponentially unstable

The C-condition requires that the tangent space R_w^m at each point w of the m -dimensional phase space W^m of the dynamical system $\{T^n\}$ should be decomposable into a direct sum of the two linear spaces X_w^k and Y_w^l with the following properties:

C1.

$$R_w^m = X_w^k \oplus Y_w^l$$

and they are such that $\{\tilde{T}^n\}$

C2.

$$\begin{aligned} a) |\tilde{T}^n \xi| &\leq a |\xi| e^{-cn} \text{ for } n \geq 0; |\tilde{T}^n \xi| \geq b |\xi| e^{-cn} \text{ for } n \leq 0, \xi \in X_w^k \\ b) |\tilde{T}^n \eta| &\geq b |\eta| e^{cn} \text{ for } n \geq 0; |\tilde{T}^n \eta| \leq a |\eta| e^{cn} \text{ for } n \leq 0, \eta \in Y_w^l, \end{aligned}$$

where the constants a, b and c are positive and are the same for all $w \in W^m$ and all $\xi \in X_w^k, \eta \in Y_w^l$. The length $|\dots|$ of the tangent vectors ξ and η is defined by the Riemannian metric ds on W^m .

The contracting and expanding foliations Σ_w^k and Σ_w^l




Figure: At each point w of the C-system the tangent space R_w^m is decomposable into a direct sum of two linear spaces Y_w^l and X_w^k . The expanding and contracting geodesic flows are γ^+ and γ^- . The expanding and contracting invariant foliations Σ_w^l and Σ_w^k are *transversal* to the geodesic flows and their corresponding tangent spaces are Y_w^l and X_w^k .

Important Example of C-system: *Torus Automorphisms*

Consider linear automorphisms of the unit hypercube in Euclidean space R^N with coordinates (u_1, \dots, u_N) where $u \in [0, 1)$

$$u_i^{(k+1)} = \sum_{j=1}^N A_{ij} u_j^{(k)}, \quad \text{mod } 1, \quad k = 0, 1, 2, \dots \quad (7)$$

- The dynamical system defined by the integer matrix A has determinant equal to one $\text{Det}A = 1$.


Important Example of C-system: *Torus Automorphisms*


Consider linear automorphisms of the unit hypercube in Euclidean space R^N with coordinates (u_1, \dots, u_N) where $u \in [0, 1)$

$$u_i^{(k+1)} = \sum_{j=1}^N A_{ij} u_j^{(k)}, \quad \text{mod } 1, \quad k = 0, 1, 2, \dots \quad (7)$$

- ▶ The dynamical system defined by the integer matrix A has determinant equal to one $\text{Det}A = 1$.
- ▶ The Anosov hyperbolicity C-condition: the matrix A has no eigenvalues on the unit circle. Thus the spectrum $\Lambda = \lambda_1, \dots, \lambda_N$ fulfils the two conditions:

$$1) \text{Det}A = \lambda_1 \lambda_2 \dots \lambda_N = 1, \quad 2) |\lambda_i| \neq 1. \quad (8)$$





The Kolmogorov entropy of a Anosov C-system is:

$$h(A) = \sum_{|\lambda_\beta| > 1} \ln |\lambda_\beta|. \quad (9)$$

The entropy $h(A)$ depends on the spectrum of the operator A .

This allows to characterise and compare the chaotic properties of dynamical C-systems quantitatively → computing and comparing their entropies.

The decorrelation time τ_0 of the dynamical system A can be expressed in terms of its entropy

$$\tau_0 = 1/h(A) \quad (10)$$

The relaxation time τ , at which the initial states occupying the volume δv_0 will spread over the whole phase space, is

$$\tau = \tau_0 \ln(1/\delta v_0) \quad (11)$$

These important characteristic time scales should fulfil the following fundamental relation

$$\tau_0 \leq T \leq \tau \quad (12)$$

where $T = 1$ is a time of one iteration.

Thus there are three characteristic time scales associated with the C-system:

$$\left(\begin{array}{c} \textit{Decorrelation} \\ \textit{time} \\ \tau_0 = \frac{\pi}{4pN^2} \end{array} \right) < \left(\begin{array}{c} \textit{Interaction} \\ \textit{time} \\ t_{int} = n = 1 \end{array} \right) < \left(\begin{array}{c} \textit{Stationary} \\ \textit{distribution time} \\ \tau = \frac{1}{h(T)} \ln \frac{1}{\delta v_0} \end{array} \right).$$

The generator $N = 256$ has the entropy $h(T) = 194$, therefore the characteristic time scales for this generator are

$$\begin{pmatrix} \text{Decorrelation} \\ \text{time} \\ \tau_0 = 0.000012 \end{pmatrix} < \begin{pmatrix} \text{Interaction} \\ \text{time} \\ t_{int} = 1 \end{pmatrix} < \begin{pmatrix} \text{Stationary} \\ \text{distribution time} \\ \tau = 95 \end{pmatrix}.$$

The MIXMAX generator $N = 240$ has the entropy $h(T) = 8679$, therefore the characteristic time scales for this generator are

$$\left(\begin{array}{c} \text{Decorrelation} \\ \text{time} \\ \tau_0 = 0.000004 \end{array} \right) < \left(\begin{array}{c} \text{Interaction} \\ \text{time} \\ t_{int} = 1 \end{array} \right) < \left(\begin{array}{c} \text{Stationary} \\ \text{distribution time} \\ \tau = 1.17 \end{array} \right).$$

Both generators have very short decorrelation time. The second generator $N = 240$ has much bigger entropy and therefore its relaxation time τ is much smaller, of order 1.17, and is close to the interaction time.

A strong instability of trajectories of a dynamical C-system leads to the appearance of statistical properties in its behaviour. As a result the time average of the function $f(w)$ on W^m

$$S_N(w) = \frac{1}{N} \sum_{n=0}^{N-1} f(A^n w) \quad (13)$$

behaves as a superposition of quantities which are statistically weakly dependent. Therefore for the C-systems on a torus it was demonstrated by Leonov that the *deviation of the time averages (13) from the phase space averages*

$$\langle f \rangle = \int_{W^m} f(w) dw \quad (14)$$

multiplied by \sqrt{N} have at large $N \rightarrow \infty$ the Gaussian distribution:

$$\lim_{N \rightarrow \infty} \mu \left\{ w : \sqrt{N} \left(S_N - \langle f \rangle \right) < z \right\} = \frac{1}{\sqrt{2\pi}\sigma_f} \int_{-\infty}^z e^{-\frac{y^2}{2\sigma_f^2}} dy.$$

The deviation of the time average S_N from its limit value $\langle f \rangle$ is asymptotically Gaussian! The quantity $\sqrt{N} \left(S_N(w) - \langle f \rangle \right)$ converges in distribution to the normal random variable with standard deviation σ_f

$$\sigma_f^2 = \sum_{n=-\infty}^{+\infty} [\langle f(w) f(A^n w) \rangle - \langle f(w) \rangle^2]. \quad (15)$$

We were able to express it in terms of entropy

$$\sigma_f^2 = \sum_{n=-\infty}^{+\infty} \frac{M^2}{128\pi^4} e^{-4nh(T)} = \frac{M^2}{128\pi^4} \frac{e^{4h(A)} + 1}{e^{4h(A)} - 1} \quad (16)$$

Spectral Test of the MIXMAX Random Number Generators

The test is aimed at answering the question of distribution of the generated pseudo-random vectors in dimensions d that are larger than the genuine dimension of a generator $d > N$.

The resolution of the MIXMAX generators in $d \leq N$ is 2^{-61} .

To perform the spectral test one should find the shortest wave vector on the lattice generated in $d = rN$ where $r = 2, 3, \dots$:

$$l_r = \frac{1}{\lambda_{min}}, \quad \text{where} \quad \lambda_{min} = \min_{\mathbf{k}} |\mathbf{k}|.$$

The generator $N = 17$, $m = 2^{36} + 1$ and $s = 0$ has spectral index

$$l_2 = 1.49 \cdot 10^{-8}.$$

This is the main advantage of the MIXMAX generators with parameter m in the region between 2^{24} and 2^{36} . At $r = 3$ the spectral index is $l_3 = 5 \cdot 10^{-4}$.

N	m	s	Entropy	$\log_{10}(q)$	spectral index
8	$m = 2^{36} + 1$	s=0	149.7	129	$1.49 \cdot 10^{-8}$
17	$m = 2^{36} + 1$	s=0	374.3	294	$1.49 \cdot 10^{-8}$
240	$m = 2^{32} + 1$	s=271828282	5445.7	4389	$7.6 \cdot 10^{-10}$

Table: Table of the new three-parameter MIXMAX generators $A(N, s, m)$ in (2).

Conclusion

Use MIXMAX for your Monte-Carlo simulations !

it will provide a fast convergence!

This work was supported in part by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No 644121.

Thank you!