

LHC detectors CMS present performances

*7th ENHEP School on High Energy Physics
26-31 January 2019*

Ain Shams University Cairo - Egypt

Ludwik Dobrzynski

Laboratoire Leprince Ringuet - Ecole polytechnique - CNRS - IN2P3

LHC detectors

CMS present performances

- ◆ *Introduction*
- ◆ *Physics objectives*
- ◆ *Hadron collider detectors*
- ◆ *Detector upgrades for future searches*
- ◆ *Conclusion*

7th ENHEP School on High Energy Physics

26-31 January 2019


Ain Shams University Cairo - Egypt

Ludwik Dobrzynski

Laboratoire Leprince Ringuet - Ecole polytechnique - CNRS - IN2P3

LHC

*Explore of a new energy frontier
in Proton-proton collisions at $E_{CM} = 13 \text{ TeV}$
and Pb-Pb collision at 5 TeV*



LHC



*Explore of a new energy frontier
in Proton-proton collisions at $E_{CM} = 13 \text{ TeV}$
and Pb-Pb collision at 5 TeV*



*LHC ring:
27 km circumference*

LHC

*Explore of a new energy frontier
in Proton-proton collisions at $E_{CM} = 13 \text{ TeV}$
and Pb-Pb collision at 5 TeV*





LHC Physics capabilities and Objectives



- *In pp mode, the physics potential comes both from the *greatly increased energy* and the *greatly increased luminosity*, offering the possibility to access processes that up till now have been too rare to be studied.*
- *In PbPb mode, annual data collection rates are comparable to **RHIC** but there is a **25-fold** increase in centre-of-mass energy.*



LHC Physics capabilities and Objectives



- *In pp mode, the physics potential comes both from the **greatly increased energy** and the **greatly increased luminosity**, offering the possibility to access processes that up till now have been too rare to be studied.*
- *In PbPb mode, annual data collection rates are comparable to **RHIC** but there is a **25-fold increase in centre-of-mass energy**.*
- *The four major LHC experiments together cover very different areas of physics.*



- *In pp mode, the physics potential comes both from the greatly increased energy and the greatly increased luminosity, offering the possibility to access processes that up till now have been too rare to be studied.*
- *In PbPb mode, annual data collection rates are comparable to RHIC but there is a 25-fold increase in centre-of-mass energy.*
- *The four major LHC experiments together cover very different areas of physics.*
 - *ALICE is designed for Pb-Pb collisions having very high multiplicities. In Pb-Pb mode the LHC can deliver 8kHz of interactions, while in pp the maximum interaction rate the experiment can handle is about 100 kHz.*



- *In pp mode, the physics potential comes both from the greatly increased energy and the greatly increased luminosity, offering the possibility to access processes that up till now have been too rare to be studied.*
- *In PbPb mode, annual data collection rates are comparable to RHIC but there is a 25-fold increase in centre-of-mass energy.*
- *The four major LHC experiments together cover very different areas of physics.*
 - *ALICE is designed for Pb-Pb collisions having very high multiplicities. In Pb-Pb mode the LHC can deliver 8kHz of interactions, while in pp the maximum interaction rate the experiment can handle is about 100 kHz.*
 - *ATLAS and CMS are looking for rare processes in pp interactions, for which they need the highest possible luminosity.*



- *In pp mode, the physics potential comes both from the **greatly increased energy** and the **greatly increased luminosity**, offering the possibility to access processes that up till now have been too rare to be studied.*
- *In PbPb mode, annual data collection rates are comparable to **RHIC** but there is a **25-fold increase in centre-of-mass energy**.*
- *The four major LHC experiments together cover very different areas of physics.*
 - *ALICE is designed for Pb-Pb collisions having very high multiplicities. In Pb-Pb mode the LHC can deliver 8kHz of interactions, while in pp the maximum interaction rate the experiment can handle is about 100 kHz.*
 - *ATLAS and CMS are looking for rare processes in pp interactions, for which they need the highest possible luminosity.*
 - *LHCb has been optimized for beauty decays requiring a very high “level 1” trigger rate (around 1 MHz). By using the trigger to select interesting decay modes, this rate is reduced to a final level trigger rate of around 200 Hz*


First LHC's achievement

LIVE

Nobelpriset 2013

The Nobel Prize 2013

The Nobel Prize in Physics 2013




Evolution of the signal
for the new particle in
2011 and 2012

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/Hig13002TWiki>

#NobelPrize

Nobelprize.org



*The Universe is very big and often
beyond the reach of our minds and
instruments*

Big ideas and powerful instruments have
enabled revolutionary progress

The image is a vertical composition. The top half shows a dense, tangled network of purple and orange filaments, representing a quark-gluon plasma. This network narrows into a thin, bright yellow and red neck, which then expands into a large, smooth, reddish-orange cone. The bottom half of the image shows a vast field of galaxies, including several prominent spiral galaxies with bright yellow cores, set against a dark background with scattered stars and smaller galaxy clusters. The overall effect is one of expansion and evolution from a microscopic state to a macroscopic cosmic structure.

this very big idea

connections between quarks
& the cosmos

How we come there

- a jiffy* after the beginning: tremendous burst of expansion (inflation) that smoothed spacetime, created hot quark soup, and turned subatomic quantum fluctuations into seeds for galaxies
- until 0.00001 sec: quark soup era during which ordinary matter and dark matter arose
- 0.00001 to 300 sec: neutrons and protons, then nuclei of the lightest elements were created
- 100,000 years to 5 billion years: gravity of dark matter builds cosmic structure from quantum seeds
- 5 billion years on: Dark energy takes over and speeds up the expansion

*a jiffy = 10^{-40} ish sec

How we come there

- a jiffy* after the beginning: tremendous burst of expansion (inflation) that smoothed spacetime, **created hot quark soup**, and turned subatomic quantum fluctuations into seeds for galaxies
- until 0.00001 sec: quark soup era during which ordinary matter and dark matter arose
- 0.00001 to 300 sec: neutrons and protons, then nuclei of the lightest elements were created
- 100,000 years to 5 billion years: gravity of dark matter builds cosmic structure from quantum seeds
- 5 billion years on: Dark energy takes over and speeds up the expansion

*a jiffy = 10^{-40} ish sec

How we come there

- a jiffy* after the beginning: tremendous burst of expansion (inflation) that smoothed spacetime, **created hot quark soup**, and turned subatomic **quantum fluctuations** into seeds for galaxies
- until 0.00001 sec: quark soup era during which ordinary matter and dark matter arose
- 0.00001 to 300 sec: neutrons and protons, then nuclei of the lightest elements were created
- 100,000 years to 5 billion years: gravity of dark matter builds cosmic structure from quantum seeds
- 5 billion years on: Dark energy takes over and speeds up the expansion

*a jiffy = 10^{-40} ish sec

How we come there

- a jiffy* after the beginning: tremendous burst of expansion (inflation) that smoothed spacetime, **created hot quark soup**, and turned subatomic **quantum fluctuations** into seeds for galaxies
- until 0.00001 sec: **quark soup era** during which ordinary matter and dark matter arose
- 0.00001 to 300 sec: neutrons and protons, then nuclei of the lightest elements were created
- 100,000 years to 5 billion years: gravity of dark matter builds cosmic structure from quantum seeds
- 5 billion years on: Dark energy takes over and speeds up the expansion

*a jiffy = 10^{-40} ish sec

How we come there

- a jiffy* after the beginning: tremendous burst of expansion (inflation) that smoothed spacetime, **created hot quark soup**, and turned subatomic **quantum fluctuations** into seeds for galaxies
- until 0.00001 sec: **quark soup era** during which ordinary matter and dark matter arose
- 0.00001 to 300 sec: **neutrons and protons**, then nuclei of the lightest elements were created
- 100,000 years to 5 billion years: gravity of dark matter builds cosmic structure from quantum seeds
- 5 billion years on: Dark energy takes over and speeds up the expansion

*a jiffy = 10^{-40} ish sec

How we come there

- a jiffy* after the beginning: tremendous burst of expansion (inflation) that smoothed spacetime, **created hot quark soup**, and turned subatomic **quantum fluctuations** into seeds for galaxies
- until 0.00001 sec: **quark soup era** during which ordinary matter and dark matter arose
- 0.00001 to 300 sec: **neutrons and protons**, then nuclei of the lightest elements were created
- 100,000 years to 5 billion years: gravity of dark matter builds **cosmic structure** from quantum seeds
- 5 billion years on: Dark energy takes over and speeds up the expansion

*a jiffy = 10^{-40} ish sec

How we come there

- a jiffy* after the beginning: tremendous burst of expansion (inflation) that smoothed spacetime, **created hot quark soup**, and turned subatomic **quantum fluctuations** into seeds for galaxies
- until 0.00001 sec: **quark soup era** during which ordinary matter and dark matter arose
- 0.00001 to 300 sec: **neutrons and protons**, then nuclei of the lightest elements were created
- 100,000 years to 5 billion years: gravity of dark matter builds **cosmic structure** from quantum seeds
- 5 billion years on: **Dark energy takes over** and speeds up the expansion

*a jiffy = 10^{-40} ish sec

Quark Soup



Quark Soup

Inflation

Quark Soup

Inflation

Early burst of enormous expansion

Quark Soup

Inflation

Early burst of enormous expansion

Baryogenesis

Quark Soup

Dark Matter

Inflation

Early burst of enormous expansion

Baryogenesis

Quark Soup

Dark Matter

WIMPs

Inflation

Early burst of enormous expansion

Baryogenesis

Quark Soup

Dark Matter

Elementary particles as Dark Matter

WIMPs

Inflation

Early burst of enormous expansion

Baryogenesis

Quark Soup

Dark Matter

Elementary particles as Dark Matter

WIMPs

Inflation

Early burst of enormous expansion

Dark Energy

Baryogenesis

Quark Soup

Dark Matter

Elementary particles as Dark Matter

WIMPs

Inflation

Early burst of enormous expansion

Baryogenesis

Dark Energy

Repulsive vacuum energy!

Quark Soup

Dark Matter

Elementary particles as Dark Matter

WIMPs

Inflation

Early burst of enormous expansion


Baryogenesis

Dark Energy


Repulsive vacuum energy!

CDM

Quark soup revolution



*Basic questions
about
the very small and the very big*



*Basic questions
about
the very small and the very big*

- What is the dark matter particle?
or is that even the right question?
- What is the nature of dark energy
and what is our cosmic destiny?
- When did inflation take place
and what is the cause of it?
- How did ordinary matter originate?
- What happened before the big bang
and where did space-time come from?

*Basic questions
about
the very small and the very big*

- **What is the dark matter particle?**
or is that even the right question?
- What is the nature of dark energy
and what is our cosmic destiny?
- When did inflation take place
and what is the cause of it?
- How did ordinary matter originate?
- What happened before the big bang
and where did space-time come from?

*Basic questions
about
the very small and the very big*

- **What is the dark matter particle?**
or is that even the right question?
- **What is the nature of dark energy**
and what is our cosmic destiny?
- When did inflation take place
and what is the cause of it?
- How did ordinary matter originate?
- What happened before the big bang
and where did space-time come from?

*Basic questions
about
the very small and the very big*

- **What is the dark matter particle?**
or is that even the right question?
- **What is the nature of dark energy**
and what is our cosmic destiny?
- **When did inflation take place**
and what is the cause of it?
- How did ordinary matter originate?
- What happened before the big bang
and where did space-time come from?

*Basic questions
about
the very small and the very big*

- **What is the dark matter particle?**
or is that even the right question?
- **What is the nature of dark energy**
and what is our cosmic destiny?
- **When did inflation take place**
and what is the cause of it?
- **How did ordinary matter originate?**
- What happened before the big bang
and where did space-time come from?

*Basic questions
about
the very small and the very big*

- **What is the dark matter particle?**
or is that even the right question?
- **What is the nature of dark energy**
and what is our cosmic destiny?
- **When did inflation take place**
and what is the cause of it?
- **How did ordinary matter originate?**
- **What happened before the big bang**
and where did space-time come from?

*Basic questions
about
the very small and the very big*

- **What is the dark matter particle?**
or is that even the right question?
- **What is the nature of dark energy**
and what is our cosmic destiny?
- **When did inflation take place**
and what is the cause of it?
- **How did ordinary matter originate?**
- **What happened before the big bang**
and where did space-time come from?

What happened to the anti-matter

Our game

Find new particles/new symmetries/new forces?

Prove and confirm models

Our game

Find new particles/new symmetries/new forces?

Prove and confirm models

More questions

Our game

Find new particles/new symmetries/new forces?

Prove and confirm models

More questions

⇒ *Origin of Mass - Higgs boson(s)*

Our game

Find new particles/new symmetries/new forces?

Prove and confirm models

More questions

⇒ *Origin of Mass - Higgs boson(s)*



Our game

Find new particles/new symmetries/new forces?

Prove and confirm models

More questions

⇒ *Origin of Mass - Higgs boson(s)*



Still open questions

*Our game**Find new particles/new symmetries/new forces?**Prove and confirm models**More questions*

- ⇒ *Origin of Mass - Higgs boson(s)* ✓
- ⇒ *Supersymmetric particles - a new zoology of particles, dark matter particle? ...*

Still open questions

*Our game**Find new particles/new symmetries/new forces?**Prove and confirm models**More questions*

- ⇒ *Origin of Mass - Higgs boson(s)* ✓
- ⇒ *Supersymmetric particles - a new zoology of particles, dark matter particle? ...*
- ⇒ *Extra space-time dimensions: gravitons, Z' etc. ?*

Still open questions

*Our game**Find new particles/new symmetries/new forces?**Prove and confirm models**More questions*

- ⇒ *Origin of Mass - Higgs boson(s)* ✓
- ⇒ *Supersymmetric particles - a new zoology of particles, dark matter particle? ...*
- ⇒ *Extra space-time dimensions: gravitons, Z' etc. ?*
- ⇒ *The Unexpected !!*

Still open questions

*Our game**Find new particles/new symmetries/new forces?**Prove and confirm models**More questions*


- ⇒ *Origin of Mass - Higgs boson(s)* ✓
- ⇒ *Supersymmetric particles - a new zoology of particles, dark matter particle? ...*
- ⇒ *Extra space-time dimensions: gravitons, Z' etc. ?*
- ⇒ *The Unexpected !!*


Still open questions

*Our game**Find new particles/new symmetries/new forces?**Prove and confirm models**More questions*


- ⇒ *Origin of Mass - Higgs boson(s)* ✓
- ⇒ *Supersymmetric particles - a new zoology of particles, dark matter particle? ...*
- ⇒ *Extra space-time dimensions: gravitons, Z' etc. ?*
- ⇒ *The Unexpected !!*

*Studies of CP Violation and Quark Gluon Plasma**Still open questions*





- **Protons are composite** Partons (valence+sea quarks, gluons) carry **longitudinal momentum fraction of the proton** (x)
- **Longitudinal parton momenta are unknown**
- **Parton distribution functions** (PDFs): estimate the momentum fraction carried by a parton inside the proton



- **Protons are composite** Partons (valence+sea quarks, gluons) carry **longitudinal momentum fraction of the proton** (x)
- **Longitudinal parton momenta are unknown**
- **Parton distribution functions** (PDFs): estimate the momentum fraction carried by a parton inside the proton


What do we want to measure



- **Protons are composite** Partons (valence+sea quarks, gluons) carry **longitudinal momentum fraction of the proton** (x)
- **Longitudinal parton momenta are unknown**
- **Parton distribution functions** (PDFs): estimate the momentum fraction carried by a parton inside the proton

What do we want to measure

- **Count the Number of particles**
- **Event topologie**
- **momentum / Energie**
- **Particle identity**
- **Transverse Missing energy/momentun**



- **Protons are composite** Partons (valence+sea quarks, gluons) carry **longitudinal momentum fraction of the proton** (x)
- **Longitudinal parton momenta are unknown**
- **Parton distribution functions** (PDFs): estimate the momentum fraction carried by a parton inside the proton

What do we want to measure

- **Count the Number of particles**
- **Event topologie**
- **momentum / Energie**
- **Particle identity**
- **Transverse Missing energy/momentun**

Can't be achieved
with a single detector




- **Protons are composite** Partons (valence+sea quarks, gluons) carry **longitudinal momentum fraction of the proton** (x)
- **Longitudinal parton momenta are unknown**
- **Parton distribution functions** (PDFs): estimate the momentum fraction carried by a parton inside the proton

What do we want to measure



- **Count the Number of particles**
- **Event topologie**
- **momentum / Energie**
- **Particle identity**
- **Transverse Missing energy/momentun**

Can't be achieved with a single detector

Integrate detectors to a detector system






The LHC detectors: a modular concept









The LHC detectors: a modular concept




The LHC detectors: a modular concept



Pattern Recognition


New particles discovered in CMS will be typically unstable and rapidly transform into a cascade of lighter, more stable and better understood particles. Particles travelling through CMS leave behind characteristic patterns, or 'signatures', in the different layers, allowing them to be identified. The presence (or not) of any new particles can then be inferred.



Key:

- Muon
- Electron
- Charged Hadron (e.g. Pion)
- - - Neutral Hadron (e.g. Neutron)
- - - Photon

Courtesy of A. Abdelalim



3.8T Superconducting Solenoid

Lead Tungstate E/M Calorimeter (ECAL)

Hermetic ($|\eta| < 5.2$) Hadron Calorimeter (HCAL) [scintillators & brass]


All Silicon Tracker (Pixels and Microstrips)

Redundant Muon System (RPCs, Drift Tubes, Cathode Strip Chambers)

3 m

ICHEP, 3 July 2014

Courtesy of A. Abdelalim



3.8T Superconducting Solenoid

Lead Tungstate E/M Calorimeter (ECAL)

Hermetic ($|\eta| < 5.2$) Hadron Calorimeter (HCAL) [scintillators & brass]

All Silicon Tracker (Pixels and Microstrips)

Redundant Muon System (RPCs, Drift Tubes, Cathode Strip Chambers)

3 m

ICHEP, 3 July 2014

Courtesy of A. Abdelalim

ICHEP, 3 July 2014

3.8T Superconducting Solenoid

Lead Tungstate E/M Calorimeter (ECAL)


Hermetic ($|\eta| < 5.2$) Hadron Calorimeter (HCAL) [scintillators & brass]

3 m

All Silicon Tracker (Pixels and Microstrips)

Redundant Muon System (RPCs, Drift Tubes, Cathode Strip Chambers)

Courtesy of A. Abdelalim



3.8T Superconducting Solenoid

Lead Tungstate E/M Calorimeter (ECAL)

Hermetic ($|\eta| < 5.2$) Hadron Calorimeter (HCAL) [scintillators & brass]

All Silicon Tracker (Pixels and Microstrips)

Redundant Muon System (RPCs, Drift Tubes, Cathode Strip Chambers)

3 m

ICHEP, 3 July 2014

Courtesy of A. Abdelalim

ICHEP, 3 July 2014

3.8T Superconducting Solenoid

Lead Tungstate E/M Calorimeter (ECAL)


Hermetic ($|\eta| < 5.2$) Hadron Calorimeter (HCAL) [scintillators & brass]

3 m

All Silicon Tracker (Pixels and Microstrips)

Redundant Muon System (RPCs, Drift Tubes, Cathode Strip Chambers)

Courtesy of A. Abdelalim



3.8T Superconducting Solenoid

Lead Tungstate E/M Calorimeter (ECAL)

Hermetic ($|\eta| < 5.2$) Hadron Calorimeter (HCAL) [scintillators & brass]

All Silicon Tracker (Pixels and Microstrips)

Redundant Muon System (RPCs, Drift Tubes, Cathode Strip Chambers)

3 m

ICHEP, 3 July 2014



Detector Requirements : The challenges

- ◆ *High Interaction Rate*
 - ◆ *pp interaction rate 1 billion interactions/s*
 - ◆ *Data can be recorded for only ~1000 out of 40 million crossings/sec*





Detector Requirements : The challenges

- ◆ *High Interaction Rate*
 - ◆ *pp interaction rate 1 billion interactions/s*
 - ◆ *Data can be recorded for only ~1000 out of 40 million crossings/sec*
- ◆ *Large Particle Multiplicity*
 - ◆ *large number of superposed events in each crossing*
 - ◆ *several 1000 tracks stream into the detector every 25 ns*
 - ◆ *need highly granular detectors with good time resolution for low occupancy*
 - ◆ *large number of channels ($\sim 100 \text{ M ch}$)*





- ◆ *High Interaction Rate*
 - ◆ *pp interaction rate 1 billion interactions/s*
 - ◆ *Data can be recorded for only ~1000 out of 40 million crossings/sec*
- ◆ *Large Particle Multiplicity*
 - ◆ *large number of superposed events in each crossing*
 - ◆ *several 1000 tracks stream into the detector every 25 ns*
 - ◆ *need highly granular detectors with good time resolution for low occupancy*
 - ◆ *large number of channels (~ 100 M ch)*
- ◆ *Tracker sensors that can withstand an extremely high radiation environment*
 - ◆ *Good track resolution in a busy environment*
 - ◆ *Innovative triggering at level 1 to keep up with the flood of data.*
 - ◆ *Level-1 trigger decision takes ~2-3 μ s*
 - ◆ *electronics need to store data locally (pipelining)*

- ◆ **High Interaction Rate**
 - ◆ pp interaction rate 1 billion interactions/s
 - ◆ Data can be recorded for only ~ 1000 out of 40 million crossings/sec
- ◆ **Large Particle Multiplicity**
 - ◆ large number of superposed events in each crossing
 - ◆ several 1000 tracks stream into the detector every 25 ns
 - ◆ need highly granular detectors with good time resolution for low occupancy
 - ◆ large number of channels ($\sim 100 \text{ M ch}$)
- ◆ **Tracker sensors that can withstand an extremely high radiation environment**
 - ◆ *Good track resolution in a busy environment*
 - ◆ *Innovative triggering at level 1 to keep up with the flood of data.*
 - ◆ Level-1 trigger decision takes $\sim 2-3 \mu\text{s}$
 - ◆ electronics need to store data locally (pipelining)



What an event with 140 vertices looks like in the CMS tracker



- ◆ **High Interaction Rate**
 - ◆ *pp interaction rate 1 billion interactions/s*
 - ◆ *Data can be recorded for only ~1000 out of 40 million crossings/sec*
- ◆ **Large Particle Multiplicity**
 - ◆ *large number of superposed events in each crossing*
 - ◆ *several 1000 tracks stream into the detector every 25 ns*
 - ◆ *need highly granular detectors with good time resolution for low occupancy*
 - ◆ *large number of channels (~ 100 M ch)*
- ◆ **Tracker sensors that can withstand an extremely high radiation environment**
 - ◆ *Good track resolution in a busy environment*
 - ◆ *Innovative triggering at level 1 to keep up with the flood of data.*
 - ◆ *Level-1 trigger decision takes ~2-3 μs*
 - ◆ *electronics need to store data locally (pipelining)*



What an event with 140 vertices looks like in the CMS tracker

- ◆ **Challenge in photon-detection**
- ◆ **Challenge in silicon**
- ◆ **Challenge in data collection / trigger**
- ◆ **High Radiation Levels**
 - ◆ *Require radiation hard (tolerant) detectors and electronics*

- ◆ **High Interaction Rate**
 - ◆ pp interaction rate 1 billion interactions/s
 - ◆ Data can be recorded for only ~ 1000 out of 40 million crossings/sec
- ◆ **Large Particle Multiplicity**
 - ◆ large number of superposed events in each crossing
 - ◆ several 1000 tracks stream into the detector every 25 ns
 - ◆ need highly granular detectors with good time resolution for low occupancy
 - ◆ large number of channels ($\sim 100 \text{ M ch}$)
- ◆ **Tracker sensors that can withstand an extremely high radiation environment**
 - ◆ Good track resolution in a busy environment
 - ◆ Innovative triggering at level 1 to keep up with the flood of data.
 - ◆ Level-1 trigger decision takes $\sim 2\text{-}3 \mu\text{s}$
 - ◆ electronics need to store data locally (pipelining)



What an event with 140 vertices looks like in the CMS tracker


- ◆ **Challenge in photon-detection**
- ◆ **Challenge in silicon**
- ◆ **Challenge in data collection / trigger**
- ◆ **High Radiation Levels**
 - ◆ Require radiation hard (tolerant) detectors and electronics





ECAL

Scintillating PbWO₄ crystals

4 Teslar
Superconducting
COIL



Plastic scintillator/brass sandwich
HCAL




IRON YOKE

TRACKER
Silicon Microstrips
Si Pixels



MUON ENDCAPS




MUON BARREL




Drift Tube
Chambers (DT)
Resistive Plate
Chambers (RPC)

Cathode Strip Chambers (CSC)
Resistive Plate Chambers (RPC)


Length: 21.6 m
Diameter: 15 m
Weight: ~12500 tons




all 5 coil modules finished in 2004
assembly in CMS hall, Jan. 2005





Insertion of coil in vacuum
tank in September 05










s.c cable: all 21 lengths (53 km) finished in 2003



Insert with superconductor









TIB+ completed

Tracker of CMS:
10 million Si-microstrips
and 70 million Si-pixels







TIB+ inserted into TOB

TEC inserted into TST






The Silicon CMS tracker





TIB+ completed

Tracker of CMS:
 10 million Si-microstrips
 and 70 million Si-pixels






TIB+ inserted into TOB



TEC inserted into TST



- Homogeneous Lead tungstate PbWO₄ crystals
- Fast scintillation response, excellent time resolution
 - about 80% of the light emitted in 25 ns
- Compact & high granularity
 - Molière radius 2.2 cm
 - Radiation length X₀ 0.89 cm
- Barrel $|\eta| < 1.48$:
 - ~61K crystals in 36 SuperModules (SM)
 - 2x2x23 cm³ covering 26 X₀
 - Photodetector: Avalanche Photo Diodes (APD)
- Endcap $1.48 < |\eta| < 3.0$
 - ~15k crystals in 4 Dees
 - 3x3x22 cm³ covering 24 X₀
 - Photodetector: Vacuum Photo Triodes (VPT)
- Preshower $1.65 < |\eta| < 2.6$
 - ~137k silicon strips in 2 planes per endcap
 - 3X₀ of lead radiator
- No longitudinal segmentation
- Energy resolution for electrons impinging on the center of a 3x3 barrel crystal matrix from Test Beam (no upstream material, no magnetic field, etc...)




$$\frac{\sigma_E}{E} = \frac{2.8\%}{\sqrt{E \text{ (GeV)}}} \oplus \frac{0.128}{E \text{ (GeV)}} \oplus 0.3\%$$




- **HCAL Barrel (HB) $0 < |\eta| < 1.3$ and Endcap (HE) $1.3 < |\eta| < 3$**
 - Sampling calorimeter, alternating layers of brass absorber and plastic scintillator tiles.
 - Hybrid photo-detector (HPD) readout
- **Outer (HO): Outside solenoid**
 - Tail catcher with scintillator layers
 - HPD readout
- **Forward (HF) at $|z|=11$ m: $2.9 < |\eta| < 5$**
 - Cherenkov light from scintillating quartz fibers in steel absorber
 - read out with conventional PMTs
- **Stability of photo-detector gains monitored using LED system**
- **Pedestals, and signal synchronization (timing) monitored using Laser data**

HF side view




- **Drift Tubes (DT) $|\eta| < 1.2$**
 - 4 stations/wheel
 - cell $42 \times 13 \text{ mm}^2$
 - gas mixture 85% Ar, 15% CO₂
 - drift velocity $\sim 55 \mu\text{m/ns}$, maximum drift time $\sim 400 \text{ ns}$
 - Time resolution $< 3 \text{ ns}$, spatial $\sim 100 \mu\text{m}$






$$x_{\text{hit}} = t_{\text{drift}} \cdot v_{\text{drift}} \equiv (t_{\text{TDC}} - t_{\text{ped}}) \cdot v_{\text{drift}}$$

- **Cathode Strip Chambers (CSC) $0.9 < |\eta| < 1.2$ (MWPC)**
 - 1 CSC has 6 layers, strips measure $r-\phi$, wires radial
 - gas 50% CO₂, 40% Ar, 10% CF₄
 - 4 stations subdivided in rings
 - Time resolution $\sim 3 \text{ ns}$, spatial $50\text{-}150 \mu\text{m}$



- **Resistive Plate Chambers (RPC) $|\eta| < 1.6$**
 - Double-gap chambers in avalanche mode
 - gas 95.2% Freon, 4.5% isobutane
 - Triggering redundancy, time resolution $< 3 \text{ ns}$ (spatial $\sim 1 \text{ cm}$)








Most front-ends follow a similar architecture :

- Very small signals (fC) -> need **amplification** and **optimisation of S/N (filter)**
- Measurement of **amplitude** and/or **time** (ADCs, discris, TDCs)
- Several thousands to millions of channels needs time to decide to keep or not the event : **memory**



Pipelined-multilevel-triggers LM



Pipelined-multilevel-triggers LM



Pipelined-multilevel-triggers LM




40 MHz LVL1 100 kHz (LVL2+LVL3) 100Hz

synchronous

asynchronous

3 μ s







40 MHz
COLLISION RATE

100 kHz
LEVEL-1 TRIGGER

1 Terabit/s
(50000 DATA CHANNELS)

500 Gigabit/s

Gigabit/s SERVICE LAN



16 Million channels
3 Gigacell buffers

1 Megabyte EVENT DATA

200 Gigabyte BUFFERS
500 Readout memories


EVENT BUILDER. A large switching network (512+512 ports) with a total throughput of approximately 500 Gbit/s forms the interconnection between the sources (Readout Dual Port Memory) and the destinations (switch to Farm Interface). The Event Manager collects the status and request of event filters and distributes event building commands (read/clear) to RDPMs.


5 TeraIPS

EVENT FILTER. It consists of a set of high performance commercial processors organized into many farms convenient for on-line and off-line applications. The farm architecture is such that a single CPU processes one event.

Petabyte ARCHIVE

CMS data flow and on(off) line computing





1 PB par an

The Worldwide LHC Computing Grid



The Worldwide LHC Computing Grid




WLCG:

An International collaboration to distribute and analyse LHC data

Integrates computer centres worldwide that provide computing and storage resource into a single infrastructure accessible by all LHC physicists

The Worldwide LHC Computing Grid



WLCG:

An International collaboration to distribute and analyse LHC data

Integrates computer centres worldwide that provide computing and storage resource into a single infrastructure accessible by all LHC physicists


The Worldwide LHC Computing Grid



Tier-0
(CERN and Hungary):
data recording,
reconstruction and
distribution

Tier-1: permanent
storage, re-
processing,
analysis

Tier-2: Simulation,
end-user analysis



nearly 160 sites,
35 countries

~250'000 cores

173 PB of storage

> 2 million jobs/day


10 Gb links

WLCG:

An International collaboration to distribute and analyse LHC data

Integrates computer centres worldwide that provide computing and storage resource into a single infrastructure accessible by all LHC physicists

Present detector status and performance



Si Strip Tracker
Lower operating temperature

Electromagnetic Calorimeter
New DAQ links

Pixel Detector
upgraded in 2017
replaced some electronics in 2018

Hadron Calorimeter
Replaced HPDs → SiPMs in Endcaps

Forward Hadron Calorimeter
upgraded in 2017



Muon Detectors
Drift tubes (VME → μ TCA ROS)
Resistive Plate chambers
Cathode strip chambers
GEM slice test (GE1/1)

Major upgrade of L1 trigger done by 2016

Trigger in 2018
L1 hardware ~100kHz
HLT software ~1kHz


Features of New Design

- Robust design: 4 barrel layers and 3 endcap disks at each end
- Smaller inner radius (new beampipe), large outer
- New readout chip with expanded buffers, embedded digitization and high speed data link
- Reduced mass with 2-phase CO₂ cooling, electronics moved to high eta, DC-DC converters




Features of New Design

- Robust design: 4 barrel layers and 3 endcap disks at each end
- Smaller inner radius (new beampipe), large outer
- New readout chip with expanded buffers, embedded digitization and high speed data link
- Reduced mass with 2-phase CO₂ cooling, electronics moved to high eta, DC-DC converters




Installed (2017)




Features of New Design


- Robust design: 4 barrel layers and 3 endcap disks at each end
- Smaller inner radius (new beampipe), large outer
- New readout chip with expanded buffers, embedded digitization and high speed data link
- Reduced mass with 2-phase CO₂ cooling, electronics moved to high eta, DC-DC converters




Installed (2017)



Initial detector




upgrade detector




Features of New Design

- Robust design: 4 barrel layers and 3 endcap disks at each end
- Smaller inner radius (new beampipe), large outer
- New readout chip with expanded buffers, embedded digitization and high speed data link
- Reduced mass with 2-phase CO₂ cooling, electronics moved to high eta, DC-DC converters





Installed (2017)



Initial detector

upgrade detector



Using same Higgs selections as 2012




Significant gain in signal reconstruction efficiency:

H →	4μ	+41%
H →	2μ2e	+48%
H →	4e	+51%

Primary vertex resolution improved by ~1.5 - 2

Mass distribution from various di-muon triggers

- Very good tracking performance for physics
- ...and very good muon trigger performance tool



Commissioning 2018:

- New optical links to CMS DAQ for faster data transmission from ECAL FEDs
- automatic recovery of front end errors for trigger and data links



Stability of the relative energy scale measured from the invariant mass distribution of $\pi^0 \rightarrow \gamma\gamma$ decays in Barrel

- continuously monitored via automatic prompt calibration tools


Alignment

- ECAL and Pre-shower (ES) aligned using 2018 data, after opening/closing CMS
- Information is used to tighten the identification cuts for electrons at HLT


Δx of the ES energy deposits wrt the tracks before and **after** alignment.



Loose electron ID






Zee invariant mass - comparing 2018 w/ 2016/17




Good electron **identification** efficiencies in 2017 data, and well modelled in simulation

Phase-I upgrade of front end electronics of HE - replaced all HPDs with SiPMs



CMS Preliminary 2018



The upgraded HE is running stably



Several benefits with the upgrade:

- Eliminated progressive HPD damage
- Increased photo detection efficiency by x2.5
- **Extend longevity of HE till the end of Run 3**
- Increased longitudinal segmentation
- Add per-channel timing information
- better S/N (e.g. for MIP)

Muon operations proceeding smoothly with good fraction of active electronics channels



DT readout system upgraded from VME → μ TCA

Excellent performance !!



Detectors performance (local hit & segment efficiencies, resolutions) are in agreement with 2017

Good bunch crossing assignment in the trigger based on RPC hits



Larger FPGAs, finer granularity input, high speed optical links

New hardware!
Limited number of boards.


Ambitious plan assume parallel running of a (part of) new system in 2015. Full replacement 2015/16 YEST

Global Trigger:
- more algorithms,
- flexibility

Calorimetry:
- improved algorithms, granularity, tower-level precision, pileup subtr.

Muons:
- 3 partitions (Barrel, Endcap, Overlap)
- explore the available information at early step of triggering.

Currently independent candidates from DTTF, CSCTF, PACT merged at GMT



More needed for phase 2!

Larger FPGAs, finer granularity input, high speed optical links

New hardware!
Limited number of boards.


Ambitious plan assume parallel running of a (part of) new system in 2015. Full replacement 2015/16 YEST

Global Trigger:
- more algorithms,
- flexibility

Calorimetry:
- improved algorithms, granularity, tower-level precision, pileup subtr.

Muons:
- 3 partitions (Barrel, Endcap, Overlap)
- explore the available information at early step of triggering.

Currently independent candidates from DTTF, CSCTF, PACT merged at GMT



More needed for phase 2!

★ *muon triggers: improved μ p_T resolution: full information from 3 systems in track finding, more FPGAs processing*

★ *calorimeter triggers: finer granularity,*
 ★ *more FPGAs processing means better: $e/\gamma/\mu$ isolation, better E_T jet/ τ resolution with PU subtraction.*

★ *Increased system flexibility and algorithm sophistication*

Larger FPGAs, finer granularity input, high speed optical links

New hardware!
Limited number of boards.

Ambitious plan assume parallel running of a (part of) new system in 2015. Full replacement 2015/16 YEST

Global Trigger:

- more algorithms,
- flexibility


Calorimetry:

- improved algorithms, granularity, tower-level precision, pileup subtr.

Muons:

- 3 partitions (Barrel, Endcap, Overlap)
- explore the available information at early step of triggering.

Currently independent candidates from DTTF, CSCTF, PACT merged at GMT



More needed for phase 2!

Trigger efficiency @ $210^{34} \text{ cm}^{-2} \text{ s}^{-1}$

Channel	Current	Upgrade
W(eν),H(bb)	37.5%	71.5%
W(μν),H(bb)	69.6%	97.9%
VBF H(ττ(μτ))	19.4%	48.4%
VBF H(ττ(ετ))	14.0%	39.0%
VBF H(ττ(ττ))	14.9%	50.1%
H(WW(eeνν))	74.2%	95.3%
H(WW(μμνν))	89.3%	99.9%
H(WW(eμνν))	86.9%	99.3%
H(WW(μeνν))	90.7%	99.7%

★ *muon triggers: improved μ p_T resolution: full information from 3 systems in track finding, more FPGAs processing*

★ *calorimeter triggers: finer granularity,*
 ★ *more FPGAs processing means better: $e/\gamma/\mu$ isolation, better E_T jet/ τ resolution with PU subtraction.*

★ *Increased system flexibility and algorithm sophistication*




Run 2 pp achievements




CMS Integrated Luminosity, pp, $\sqrt{s} = 13$ TeV

Data included from 2015-06-03 08:41 to 2018-10-24 04:00 UTC



CMS Integrated Luminosity, pp, $\sqrt{s} = 13$ TeV

Data included from 2015-06-03 08:41 to 2018-10-24 04:00 UTC



- About 140 fb^{-1} certified good for physics in Run 2. Overall efficiencies:
 - 92.3% recording efficiency in Run2
 - 92.8% validation efficiency in Run2




Run 2 pp achievements




CMS Integrated Luminosity, pp, $\sqrt{s} = 13$ TeV

Data included from 2015-06-03 08:41 to 2018-10-24 04:00 UTC



CMS Integrated Luminosity, pp

Data included from 2010-03-30 11:22 to 2018-10-24 04:00 UTC



- About 140 fb⁻¹ certified good for physics in Run 2. Overall efficiencies:
 - 92.3% recording efficiency in Run2
 - 92.8% validation efficiency in Run2




Run 2 pp achievements




CMS Integrated Luminosity, pp, $\sqrt{s} = 13$ TeV

Data included from 2015-06-03 08:41 to 2018-10-24 04:00 UTC



CMS Integrated Luminosity, pp

Data included from 2010-03-30 11:22 to 2018-10-24 04:00 UTC



- About 140 fb⁻¹ certified good for physics in Run 2. Overall efficiencies:
 - 92.3% recording efficiency in Run2
 - 92.8% validation efficiency in Run2

- Final score is:
 - 68.2 fb⁻¹ (offline preliminary) delivered to CMS in 2018
 - 163 fb⁻¹ delivered overall in Run 2
 - 192.5 fb⁻¹ from 2010




Publications



- Show all
- Total
- Exotica
- Standard Model
- Supersymmetry
- Higgs
- Top Physics
- Heavy Ion
- B Physics
- Forward Physics
- Beyond 2 Generations
- Detector Performance



829 collider data papers submitted as of 2018-11-27



CMS has submitted, as of December 1st 2018, 829 publications on collisions data in a wide variety of physics (and detector) topics.

- 265 Run 2 publications*

Typically 3 papers per week





Detector challenges:

- x 10 more radiation ($\sim 10^{16} \text{ neq/cm}^2$; 10 MGy)
- x 10 more pile-up

- Run1: $\langle \mu \rangle = 20$; $\langle n_{\text{PU jets } pT > 30 \text{ GeV}} \rangle \sim 0.04$

- HL-LHC: $\times 10$ $\langle \mu \rangle = 200$; $\langle n_{\text{PU jets } pT > 30 \text{ GeV}} \rangle \sim 7.4$ $\times 185$



Detector challenges:

- x 10 more radiation ($\sim 10^{16} \text{ neq/cm}^2$; 10 MGy)
- x 10 more pile-up

- **Run1:** $\langle \mu \rangle = 20$; $\langle n_{\text{PU jets } pT > 30 \text{ GeV}} \rangle \sim 0.04$

- **HL-LHC:** $\times 10$ $\langle \mu \rangle = 200$; $\langle n_{\text{PU jets } pT > 30 \text{ GeV}} \rangle \sim 7.4$ $\times 185$

Upgrades needed to:

- keep performance (tracking, b -tag, jet/ E_{miss} ,...)
- Trigger rates acceptable with low $P_{\mathcal{T}}$ thresholds