

VH and VBF Higgs production ATLAS and CMS

Joe Taenzer (Tel Aviv University)

On behalf of the ATLAS and CMS collaborations

Higgs Couplings 2018

Tokyo, Japan

November 26th, 2018



New results since last year

ATLAS

- $H \rightarrow \gamma\gamma$ with 80 fb⁻¹, **Preliminary** ([LINK](#))
- $H \rightarrow ZZ$ with 80 fb⁻¹, **Preliminary** ([LINK](#))
- $H \rightarrow WW$ with 36 fb⁻¹, Submitted to **Phys. Lett. B** ([LINK](#))
- H (125 GeV) combination with up to 80 fb⁻¹, **Preliminary** ([LINK](#))

CMS

- $H \rightarrow \gamma\gamma$ with 36 fb⁻¹, Accepted for publication in **J. High Energy Phys.** ([LINK](#))
- $H \rightarrow ZZ$ with 80 fb⁻¹, **Preliminary** ([LINK](#))
- $H \rightarrow WW$ with 36 fb⁻¹, Submitted to **Phys. Lett. B** ([LINK](#))
- H (125 GeV) combination with 36 fb⁻¹, Submitted to **Eur. Phys. J. C** ([LINK](#))


Covered in other talks:

- ATLAS $H \rightarrow \tau\tau$ with 36 fb⁻¹, Submitted to **Phys. Rev. D** ([LINK](#))
- CMS $H \rightarrow \tau\tau$ with 36 fb⁻¹, Submitted to **J. High Energy Phys.** ([LINK](#))
- V($H \rightarrow bb$) observation with 80 fb⁻¹ by both ATLAS and CMS ([ATLAS PUBLICATION](#), [CMS PUBLICATION](#))

Outline




- Higgs production/decays - VH/VBF
- Simplified Template Cross-Sections (STXS)
- VH/VBF results in $H \rightarrow ZZ$ from ATLAS / CMS
- VH/VBF results in $H \rightarrow \gamma\gamma$ from ATLAS / CMS
- VH/VBF results in $H \rightarrow WW$ from ATLAS / CMS

Higgs production modes




VBF cross-section around 1 order of magnitude less than **ggF**, **VH** cross-section around $\frac{1}{2}$ that of **VBF**

Direct **HVV** coupling in LO **VBF/VH** production, while **ggF** has a fermion loop






Higgs decay modes



Highest BRs for 125 GeV Higgs are bb , WW , gg

Can't ignore detector signature! e.g.
 $H \rightarrow ZZ \rightarrow 4l$ is much cleaner than $H \rightarrow bb \rightarrow \text{jetjet!}$



Simplified Template X-Sections

STXS

Goals:

- More finely grained measurements than production mode measurements
- Reduce theoretical uncertainties folded into measurements
→ Shift dominant theory uncertainties to the interpretation level
- Isolate possible BSM effects into STXS bins designed for BSM sensitivity


Stages:

(increasing granularity)

- Stage 0 → Higgs production mode cross-section measurements in $|y_H| < 2.5$
ggF, VBF, VH, ttH
- Stage 1 → 31 particle level categories (bins)
Current data lacks sensitivity to resolve all Stage 1 categories
→ Reduced stage 1 measurements merge stage 1 bins where necessary

VH/VBF $H \rightarrow ZZ$ – ATLAS

Event categorization and reconstruction level signal composition




- Reconstructed event categories aimed at reduced stage 1 STXS measurement
- ggF production dominant even in VBF enriched VH hadronic categories (60-80% ggF)
- BDTs are used to improve the discrimination between production modes in each category

VH/VBF $H \rightarrow ZZ$ – ATLAS

BDT Discriminants


Reconstructed event category	BDT discriminant	Input variables
$0j$	BDT_{ggF}	$p_T^{4\ell}, \eta_{4\ell}, D_{ZZ^*}$
$1j\text{-}p_T^{4\ell}\text{-Low}$	$\text{BDT}_{\text{VBF}}^{1j\text{-}p_T^{4\ell}\text{-Low}}$	$p_T^j, \eta_j, \Delta R(j, 4\ell)$
$1j\text{-}p_T^{4\ell}\text{-Med}$	$\text{BDT}_{\text{VBF}}^{1j\text{-}p_T^{4\ell}\text{-Med}}$	$p_T^j, \eta_j, \Delta R(j, 4\ell)$
$1j\text{-}p_T^{4\ell}\text{-High}$	-	-
VBF-enriched- $p_T^j\text{-Low}$	BDT_{VBF}	$m_{jj}, \Delta\eta_{jj}, p_T^{j1}, p_T^{j2}, \eta_{4\ell}^*, \Delta R_{jZ}^{\text{min}}, (p_T^{4\ell jj})_{\text{constrained}}$
VBF-enriched- $p_T^j\text{-High}$	-	-
VH-Had-enriched	$\text{BDT}_{\text{VH-Had}}$	$m_{jj}, \Delta\eta_{jj}, p_T^{j1}, p_T^{j2}, \eta_{4\ell}^*, \Delta R_{jZ}^{\text{min}}, \eta_{j1}$
VH-Lep-enriched	-	-
$t\bar{t}H$ -enriched	-	-

Discriminants in the 1-jet VBF categories are designed to disentangle VBF and ggF production




VH/VBF $H \rightarrow ZZ$ – ATLAS

Cross-sections



STXS Stage 0 Cross-sections:

- VBF observed cross-section 3x larger than SM prediction
- VH observed cross-section consistent with SM prediction



Reduced STXS stage 1 cross-sections

- VBF observed cross-sections: Larger deviation from SM prediction in the low leading jet p_T category
- VH observed cross-sections are consistent with the SM predictions

VH/VBF H \rightarrow ZZ – CMS

Event categorization and kinematic discriminant definitions

Full kinematic information from each event is used

- Higgs decays and associated particles extracted using ME calculations
- Used to form kinematic discriminants, e.g.

$$\mathcal{D}_{\text{bkg}}^{\text{VBF+dec}} = \frac{\mathcal{P}_{\text{sig}}^{\text{VBF+VH+dec}}(\vec{\Omega})}{\mathcal{P}_{\text{sig}}^{\text{VBF+VH+dec}}(\vec{\Omega}) + c^{\text{VBF2jet}}(m_{4\ell}) \times (\mathcal{P}_{\text{bkg}}^{\text{VBS+VVV}}(\vec{\Omega}) + \mathcal{P}_{\text{bkg}}^{\text{QCD+dec}}(\vec{\Omega}))}$$

$$\mathcal{D}_{\text{bkg}}^{\text{VH+dec}} = \frac{\mathcal{P}_{\text{sig}}^{\text{VBF+VH+dec}}(\vec{\Omega})}{\mathcal{P}_{\text{sig}}^{\text{VBF+VH+dec}}(\vec{\Omega}) + c^{\text{had.VH}}(m_{4\ell}) \times (\mathcal{P}_{\text{bkg}}^{\text{VBS+VVV}}(\vec{\Omega}) + \mathcal{P}_{\text{bkg}}^{\text{QCD+dec}}(\vec{\Omega}))'}$$

Category	Leptons	Jets	Discriminant
VBF-2jet	4	2-3 (< 1 b-tag) 4 (0 b-tags)	$D_{2\text{jet}} > 0.5$
VH-hadronic	4	2-3 (< 1 b-tag) 4 (0 b-tags)	$\max(D_{\text{WH}}, D_{\text{ZH}}) > 0.5$
VH-leptonic	4 + 1 (WH) 4 + 2 (ZH)	< 3 jets (0 b-tags)	-
VBF-1jet	4	1	$D_{1\text{jet}} > 0.5$
Untagged			



Categorization order

Note: ttH not shown



VH/VBF H \rightarrow ZZ – CMS




Kinematic discriminant distributions (2017 dataset)



Good separation between VBF/VH and other Higgs production modes
→ Working as designed

VH/VBF H → ZZ – CMS

Cross-sections and signal strengths




Combination of 2016 and 2017 datasets

Cross-sections and signal strengths are generally consistent with the SM

VH/VBF H $\rightarrow \gamma\gamma$ – ATLAS



Event categorization and reconstruction level signal composition



- Reconstructed event categories aimed at stage 1 STXS measurement
→ 29 categories in total!
- VH leptonic categories signal composition ~70-80% VH
- Hadronic VH 25-40% of signal is VH
→ Large ggF contamination
- VBF categories signal composition varies from 25-90% VBF
→ Large ggF contamination

VH/VBF $H \rightarrow \gamma\gamma$ – ATLAS

Mass Spectra




- Signal modeled by a double-sided Crystal Ball function
- Continuum background modeled by a function that depends on the region
 → background fits are performed in $m_{\gamma\gamma}$ sidebands
- Other backgrounds (e.g. $V\gamma\gamma$ for VH) are obtained from simulation
 → Other Higgs production modes are included in the “Total background” line in the above plots


VH/VBF $H \rightarrow \gamma\gamma$ – ATLAS

Cross-sections

STXS Stage 0 Cross-sections



Reduced STXS stage 1 cross-sections




VBF and VH observed cross-sections are compatible with the SM prediction

VH/VBF $H \rightarrow \gamma\gamma$ – CMS

Event categorization and reconstruction level signal composition

CMS Simulation $H \rightarrow \gamma\gamma$


35.9 fb⁻¹ (13 TeV)



10-25% ggF contamination in VBF and VH hadronic categories

VH/VBF $H \rightarrow \gamma\gamma$ – CMS




Mass Spectra



- Signal model for each production process: sum of up to 5 Gaussians
 - Final fit function in each category: sum of normalized function for each production process
- Background model: Fit function included as a discrete nuisance parameter
 - Exponential, power law, polynomial functions (and more!) are all tried
 - Statistical fit penalized for N degrees of freedom in fit function

VH/VBF $H \rightarrow \gamma\gamma$ – CMS

cross-sections and signal strengths





VBF cross-section and signal strength measurements are consistent with the SM

VH cross-sections and signal strengths slightly larger than the SM prediction.

VBF H → WW – ATLAS

Post-fit BDT scores (VBF)



Signal strength

$$\mu_{\text{VBF}} = 0.62^{+0.30}_{-0.28}(\text{stat.}) \pm 0.13(\text{theo syst.}) \pm 0.16(\text{exp syst.}) = 0.62^{+0.37}_{-0.36}$$

$$\sigma_{\text{VBF}} * \text{BR}_{H \rightarrow WW}$$

Expected: $0.81 \pm 0.02 \text{ pb}$

$$\sigma_{\text{VBF}} \cdot \mathcal{B}_{H \rightarrow WW^*} = 0.50^{+0.24}_{-0.23}(\text{stat.}) \pm 0.11(\text{theo syst.}) \pm 0.13(\text{exp syst.}) \text{ pb} = 0.50^{+0.30}_{-0.29} \text{ pb}$$

Significance observed (expected):


1.9σ (2.7σ)

Discriminant variable || BDT
 BDT input variables || $m_{jj}, \Delta y_{jj}, m_{\ell\ell}, \Delta\phi_{\ell\ell}, m_T, \sum_{\ell} C_{\ell}, \sum_{\ell,j} m_{\ell j}, p_T^{\text{tot}}$

BDT used to enhance discriminating power between signal (VBF) and backgrounds, **including ggF!**

VH/VBF H \rightarrow WW – CMS


CMS Simulation 35.9 fb⁻¹ (13 TeV)



Significant ggF contamination in **VBF-tagged** and **VH-tagged** categories.

WH-tagged (3 lepton) and **ZH-tagged** (4 lepton) very pure in WH/ZH

CMS 35.9 fb⁻¹ (13 TeV)



VH leptonic and **VBF** cross-sections are consistent with SM, **VH hadronic** cross-section is larger than the SM prediction

Summary

ATLAS



- Stage 0 and reduced Stage 1 STXS measurements from $H \rightarrow ZZ$ and $H \rightarrow \gamma\gamma$
 - Both preliminary results used 80fb^{-1} of integrated luminosity
 - Generally consistent with SM predictions
- VBF $H \rightarrow WW$ $\sigma_{\text{VBF}} \cdot \text{BR}_{H \rightarrow WW}$ and signal strength measurement with 36fb^{-1}
 - Consistent with SM prediction

CMS

- Stage 0 STXS measurements from $H \rightarrow ZZ$, $H \rightarrow \gamma\gamma$, $H \rightarrow WW$
 - $H \rightarrow ZZ$ result used 80fb^{-1} of integrated luminosity, $H \rightarrow \gamma\gamma$ and $H \rightarrow WW$ used 36fb^{-1}
 - Generally consistent with SM predictions



BACKUP SLIDES

H(125GeV) Combination ATLAS



- Combined measurements of Higgs production cross-sections in the ZZ, yy, WW, bb, $\tau\tau$, and $\mu\mu$ decay modes
- Not all analyses were performed with the same integrated luminosity:
 - ZZ, yy, and $\mu\mu \Rightarrow 80 \text{ fb}^{-1}$
 - WW and $\tau\tau \Rightarrow 36 \text{ fb}^{-1}$
 - tt(H \rightarrow bb), ttH multi lep $\Rightarrow 36 \text{ fb}^{-1}$
- Generally consistent with the SM prediction(s)!


H(125GeV) Combination CMS



- Combined measurements of Higgs production cross-sections in the ZZ, $\gamma\gamma$, WW, bb, $\tau\tau$, and $\mu\mu$ decay modes
- Integrated luminosity of 35.9 fb⁻¹ for all analyses
- Generally consistent with SM predictions


VH/VBF H → ZZ – ATLAS

Event categorization for STXS stage 0 and reduced stage 1



VH/VBF H \rightarrow yy – ATLAS

Event categorization for STXS



*VBF-like: $m_{jj} > 400$ GeV, $|\Delta y_{jj}| > 2.8$

†VH-like: $60 < m_{jj} < 120$ GeV

VH → tautau – CMS

Event Selection

WH selection

$$p_T^{\tau_h} > 20 \text{ GeV}, |\eta^{\tau_h}| < 2.3, I^e < 0.1, I^\mu < 0.15, \text{b veto}$$

Channel	Trigger ($p_T/ \eta $)	Lepton selection: p_T (GeV)	τ_h selection: isolation
$e\mu\tau_h$	$\mu(22/2.1)$ or $e(25/2.1)$	$p_T^e > 15$ or $26, p_T^\mu > 23$ or 15	MVA τ_h (60% eff.)
$\mu\mu\tau_h$	$\mu(22/2.1)$	$p_T^\mu > 23, p_T^\mu > 15$	MVA τ_h (60% eff.)
$e\tau_h\tau_h$	$e(25/2.1)$	$p_T^e > 26$	MVA τ_h (55 or 65% eff.)
$\mu\tau_h\tau_h$	$\mu(22/2.1)$	$p_T^\mu > 23$	MVA τ_h (55 or 65% eff.)

ZH selection

Z boson reconstructed from opposite charge, same-flavor light leptons, $60 < m_{\ell\ell} < 120$ GeV, b veto

τ_h baseline requirements: $p_T^{\tau_h} > 20, |\eta^{\tau_h}| < 2.3, \text{MVA } \tau_h$ (65% efficiency)



e baseline requirements: $p_T^e > 10, |\eta^e| < 2.5, \text{MVA ID}$ (90% efficiency)

μ baseline requirements: $p_T^\mu > 10, |\eta^\mu| < 2.4, \mu \text{ ID}$ (> 99% efficiency), $I^\mu < 0.25$

Channel	Trigger ($p_T/ \eta $)	Lepton selection: p_T (GeV)	Lepton selection: isolation
$ee\mu\tau_h$			$I^\mu < 0.15$
$eee\tau_h$	$[e_1(23/2.5) \& e_2(12/2.5)]$	$[p_T^{e1} > 24 \& p_T^{e2} > 13]$	e ID (80% eff.), $I^e < 0.15$
$ee\tau_h\tau_h$	or $e_1(27/2.5)$	or $p_T^{e1} > 28$	baseline selection listed above
$eee\mu$			e ID (80% eff.), $I^e < 0.15, I^\mu < 0.15$
$\mu\mu\mu\tau_h$			$I^\mu < 0.15$
$\mu\mu e\tau_h$	$[\mu_1(17/2.4) \& \mu_2(8/2.4)]$	$[p_T^{\mu1} > 18 \& p_T^{\mu2} > 10]$	e ID (80% eff.), $I^e < 0.15$
$\mu\mu\tau_h\tau_h$	or $\mu_1(24/2.4)$	or $p_T^{\mu1} > 25$	baseline selection listed above
$\mu\mu e\mu$			e ID (80% eff.), $I^e < 0.15, I^\mu < 0.15$

VH \rightarrow $\tau\tau$ – CMS

$M_{\tau\tau}$



VH \rightarrow tautau – CMS

Cross-section measurement

