


Exotic Hadrons and Flavor Physics: May 28 – June 1, 2018

Interplay Between B-Anomalies and High- p_T Physics

Jernej F. Kamenik



Institut
"Jožef Stefan"
Ljubljana, Slovenija



Univerza v Ljubljani

Fakulteta za matematiko in fiziko

Stony Brook
31/5/2018

Prelude: Lepton Flavor Universality in SM

SM gauge sector respects accidental flavor symmetry

$$G_F^{\text{SM}} = U(3)_Q \times U(3)_U \times U(3)_D \times \boxed{U(3)_L \times U(3)_E} \quad \text{*neutrino masses}$$

LFU

Broken by Higgs Yukawas

$$G_{\text{acc.}}^{\text{SM}} = U(1)_B \times U(1)_e \times U(1)_\mu \times U(1)_\tau$$

⇒ Unique source of LFU breaking: $m_e \neq m_\mu \neq m_\tau$

Any LFU violation beyond lepton mass effects sign of NP!

Crash course on B-anomalies: charged currents

Charged-current semileptonic B decays:

$$\frac{d^2\Gamma(B \rightarrow D^{(*)}\ell\nu)}{dq^2 d\cos\theta} = |V_{cb}|^2 \mathcal{N}_0 \sum_{\lambda} h_{\lambda}(\cos\theta) \Phi_{\lambda}(q^2) \quad (\ell = e, \mu)$$

Hadronic form factor normalization
computed using Lattice QCD

Helicity basis functions
(determine angular distributions)

(Relative) form factor shapes
extracted (fitted) from data

*fit model dependence
Bernlochner et al., 1703.05330
Bigi et al., 1707.09509
see also talk by Michele

Comparison with measured rates yields


$$|V_{cb}|_{\text{excl.}} = (38.99 \pm 1.27) \times 10^{-3}$$

Crash course on B-anomalies: charged currents

Charged-current

d^2

Hadronic
computed



q^2)

$(\ell = e, \mu)$

shapes
data

*fit model dependence
Bernlochner et al., 1703.05330
Bigi et al., 1707.09509
see also talk by Michele

Comparison with measured rates yields

$$|V_{cb}|_{\text{excl.}} = (38.99 \pm 1.27) \times 10^{-3}$$

Crash course on B-anomalies: charged currents

Charged-current semitauonic B decays:

$$\frac{d^2\Gamma(B \rightarrow D^{(*)}\tau\nu)}{dq^2 d\cos\theta} = |V_{cb}|^2 \mathcal{N}_0 \sum_{\lambda} h_{\lambda}(\cos\theta) \Phi_{\lambda}(q^2, m_{\tau})$$

- Kinematical (phase space) effects
- One additional relative f.f. shape (tau helicity suppressed)

Can be systematically computed (Lattice QCD, HQET)

JFK & Mescia, 0802.3790
Fajfer et al., 1203.2654
Bernlochner et al., 1703.05330
Bigi et al., 1707.09509
see also talk by Michele


LFU ratios can be determined very precisely

$$R(D^{(*)}) \equiv \frac{\mathcal{B}(B \rightarrow D^{(*)}\tau\nu)}{\mathcal{B}(B \rightarrow D^{(*)}\ell\nu)}$$

*QED effects
de Boer et al., 1803.05881

Semileptonic B anomalies: LFU in charged currents

Intriguing experimental situation



* $R(B_c \rightarrow J/\psi)$
LHCb, 1711.05623

SM (subleading f.f.) uncertainties insignificant at this point

Crash course on B-Anomalies: neutral currents

Neutral-current mediated rare semileptonic B decays:

$$\frac{d^2\Gamma(B \rightarrow K^{(*)}\ell^+\ell^-)}{dq^2 d\Omega} = \frac{\alpha|V_{tb}V_{ts}|^2}{4\pi} \mathcal{N}_0 \sum_{\lambda} h_{\lambda}(\Omega) I_{\lambda}(q^2)$$



FCNCs - loop suppressed in SM

LD hadronic effects due to
hadronic substructure of the photon
not captured by local f.f.'s (c \bar{c} res.)

Can form th. clean LFU ratios: $R_{K^{(*)}} \equiv \frac{\Gamma(B \rightarrow K^{(*)}\mu\mu)}{\Gamma(B \rightarrow K^{(*)}ee)} \bigg|_{q_{\min}^2}^{q_{\max}^2}$

Non LFU effects tiny away from threshold $R_{K^{(*)}} = 1 + \mathcal{O}(m_{\mu}^2/m_B^2)$

Semileptonic B anomalies: neutral currents




Corroborating evidence
from angular analyses,
rate measurements:

$$\mathcal{H}_{\text{eff}}^{\text{NP}} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \frac{e^2}{16\pi^2} \sum_{i,\ell} (C_i^\ell O_i^\ell + C_i^{\prime\ell} O_i^{\prime\ell}) + \text{h.c.},$$

$$O_9^\ell = (\bar{s}\gamma_\mu P_L b)(\bar{\ell}\gamma^\mu \ell),$$

$$O_{10}^\ell = (\bar{s}\gamma_\mu P_L b)(\bar{\ell}\gamma^\mu \gamma_5 \ell),$$



Implications for high p_T : general considerations

B-anomalies in presence of (heavy) NP:

$$\mathcal{L}_{\text{BSM}} \rightarrow \mathcal{L}_{\nu\text{SM}} + \sum_{i, (d>4)} \frac{\mathcal{Q}_i^{(d)}}{\Lambda^{d-4}}$$

Deviations in flavor \Rightarrow indications of NP scale

$$[\text{scale}] = \frac{[\text{mass}]}{[\text{coupling}]}$$

Unitarity/Perturbativity

\Rightarrow upper bound on coupling

\Rightarrow upper bound on NP d.o.f. mass

Implications for high p_T : general considerations

LFUV in $R(D^{(*)})$: $\Lambda \simeq 2.5 \text{ TeV}$ e.g. $\mathcal{Q} = (\bar{c}\gamma_\mu P_L b)(\bar{\tau}\gamma_\mu P_L \nu)$

\Rightarrow tree-unitarity $M_{\text{NP}} \lesssim 6.5 \text{ TeV}$

up to the edge of LHC kinematical reach

see e.g.
Altmannshofer et al., 1704.06659

LFUV in $R_{K^{(*)}}$ (& other obs.) : $\Lambda \sim 40 \text{ TeV}$

e.g. $\mathcal{Q} = (\bar{s}\gamma_\mu P_L b)(\bar{\mu}\gamma^\mu P_L \mu)$

\Rightarrow NP d.o.f.s accessible at LHC only if their couplings to bs and/or $\mu\mu$ suppressed!

Implications of LFUV for NP flavor breaking

NP needs to respect SM gauge symmetry

$$\mathcal{Q}_i[Q, D, U, L, E]$$

At EW scale: in terms of four-fermion operators

$$R_{K^{(*)}} \left(\begin{array}{l} \epsilon_{ij}^L \epsilon_{kl}^Q (\bar{L}_i L_j) (\bar{Q}_k Q_l) \\ \epsilon_{ij}^E \epsilon_{kl}^Q (\bar{E}_i E_j) (\bar{Q}_k Q_l) \end{array} \right) \quad \begin{array}{l} \epsilon_{ij}^{EL} \epsilon_{kl}^{QD} (\bar{E}_i H^\dagger L_j) (\bar{Q}_k H D_l) \\ \epsilon_{ij}^{LE} \epsilon_{kl}^{QU} (\bar{L}_i H E_j) (\bar{Q}_k \tilde{H} U_l) \end{array} R(D^{(*)})$$

Buttazzo et al., 1706.07808

$$\epsilon_{\mu\mu}^{L,E}, \epsilon_{sb}^Q \neq 0$$

$$\epsilon_{\tau i}^{L,EL}, \epsilon_{cb}^{Q,QD} \neq 0$$

$$\epsilon_{i\tau}^{LE}, \epsilon_{bc}^{QU} \neq 0$$

*B_c lifetime, decays
Alonso et al., 1611.06676
Akeroyd & Chen, 1708.04072

*right-handed currents
Greljo et al., 1804.04642

Simplest UV:

Z'/W'

LQ's

H^\pm

Implications of LFUV for NP flavor breaking


Absence of BSM LFUV, FCNCs in Kaon, Charm, Tau decays requires approximate alignment with the 3rd generation

$$\epsilon_{sb}^Q \propto V_{tb} V_{ts} \quad \epsilon_{cb}^Q \propto V_{cb}$$

$U(2)_F$, MFV

$$\epsilon_{ij}^Q \simeq \cancel{1} + \mathcal{O}(\mathbf{Y}\mathbf{Y}^\dagger)$$

Not universal! High pT di-muon searches



Fajfer et al., 1206.1872
Bordone et al., 1702.07238

Immediate implications for LHC

Flavor alignment implies lower NP scale:

$$(\bar{Q}_3 Q_3)(\bar{L}_3 L_3) \rightarrow V_{cb}(\bar{c}b)(\bar{\tau}\nu)$$

$\Rightarrow R(D^{(*)})$ anomaly

$$\Lambda\sqrt{|V_{cb}|} \sim 500 \text{ GeV}$$

Well within LHC reach!

$$(\bar{Q}_3 Q_3)(\bar{L}_2 L_2) \rightarrow V_{tb}V_{ts}(\bar{s}b)(\bar{\mu}\mu)$$

$\Rightarrow R_{K^{(*)}}$ anomaly

$$\Lambda\sqrt{|V_{ts}|} \sim 8 \text{ TeV}$$

Still only marginally!

Immediate implications for LHC: $R(D^{(*)})$

Enhanced LFUV in top processes:



$$(\bar{Q}_3 Q_3)(\bar{L}_3 L_3) \rightarrow V_{cb}(\bar{c}b)(\bar{\tau}\nu) + \boxed{V_{tb}(\bar{t}b)(\bar{\tau}\nu)}$$

Currently tested to O(10%)

$$\mathcal{B}_e = 13.3(4)(4)\%, \mathcal{B}_\mu = 13.4(3)(5)\%, \mathcal{B}_{\tau_h} = 7.0(3)(5)\%,$$

ATLAS, 1506.05074

Example simplified model: charged spin-1 boson (W')



LHC measurements starting to constrain $m_{W'} < m_t$ region.


Immediate implications for LHC: $R(D^{(*)})$

Weak gauge invariance \Rightarrow neutral currents

$$(\bar{Q}_3 Q_3)(\bar{L}_3 L_3) \rightarrow V_{cb}(\bar{c}b)(\bar{\tau}\nu) + V_{tb}(\bar{t}b)(\bar{\tau}\nu) + (\bar{b}b)(\bar{\tau}\tau)$$

Constraints from existing $pp \rightarrow \tau^+ \tau^-$ searches at LHC

Faroughy, Greljo & JFK, 1609.07138



W'/Z' explanation only allowed if light ($M < 400$ GeV) or broad ($\Gamma/M > 20\%$)

Leptoquark, charged scalar explanations disfavored

Departures from strict $U(2)_F$ limit can ameliorate the bounds

see e.g.
Buttazzo et al., 1706.07808

Importance of EW radiative corrections: LFV

LFUV in absence of LFV:

MLFV


$$\epsilon_{\mu i}^L \propto \delta_{\mu i}$$

$$\epsilon_{\tau i}^L \propto \delta_{\tau i}$$

See e.g.
Alonso et al., 1505.05164

see also
Glashow et al., 1411.0565

Mixing of 4-fermion operators under EW RGE evolution



Feruglio et al., 1606.00524
1705.00929

Induces LFUV in decays of heavy leptons; in Z,W couplings

⇒ Severely constrains combined explanations of LFUV
in charged and neutral current B decays

⇒ Implications for UV model building (cancelations)

see also
Buttazzo et al., 1706.07808
Cornella et al., 1803.00945


di Luzio et al., 1708.08450
Bordone et al., 1712.01368
1805.09328

Importance of EW radiative corrections: B-anomalies without new quark flavor violation

Starting with flavor conserving non-universal operators:

$$(\bar{L}_2 L_2)(\bar{U}_3 U_3) \qquad (\bar{E}_2 E_2)(\bar{U}_3 U_3)$$

EW matching & RGE induce LFUV in rare FCNC B decays



Aebischer et al., 1512.02830
Faroughy et al., 1805.04917

see also
Blanger, Delaunay, & Westhoff, 1507.06660
Bauer et al., 1511.01900
Becirevic & Sumensari, 1704.05835

Effective NP scale now loop-suppressed: $\Lambda \frac{\sqrt{|V_{ts}|}}{4\pi} \sim 600 \text{ GeV}$

\Rightarrow automatically respects 3rd gen. alignment

\Rightarrow d.o.f.'s mediating R_K well within LHC kinematical reach


$R_{K^{(*)}}$ without new flavor violation: a UV completion

JFK, Soreq & Zupan, 1704.06005

VL quark partner of right-handed top (T),

- charged under gauged $U(1)'$ (Z' , h')
- T - t_R mix after $U(1)'$ breaking - induced $U(1)'$ charge of t_R (similar mechanism possible to induce muon $U(1)'$ charge)

R_K induced through W loops



Dominant signatures:


- neutrino trident

- $pp \rightarrow 4t$

Alvarez et al., 1611.05032

- $pp \rightarrow 2\mu 2t$

see also
Fox et al., 1801.03505



$R_{K^{(*)}}$ without new flavor violation: a UV completion

JFK, Soreq & Zupan, 1704.06005


VL quark partner of right-handed top (T),

- charged under gauged $U(1)'$ (Z' , h')
 - T - t_R mix after $U(1)'$ breaking - induced $U(1)'$ charge of t_R
- (similar mechanism possible to induce muon $U(1)'$ charge)

Complementary constraints:

- Flavor (LHCb, BelleII)
- LHC top & muon production
- Low energy colliders: neutrino trident

Existing experiments should confirm/exclude



Conclusions

Flavor is powerful guide to high- p_T searches at LHC:

- In case of significant signals of NP in flavor observables can identify prospective LHC experimental targets

Generally, NP d.o.f.'s accommodating tentative B-anomalies could be beyond LHC reach.

Phenomenological and model-building considerations point towards more optimistic scenarios

- Low energy constraints can point to lighter mediators!
- Example of fruitful interplay between NP searches at energy and intensity frontiers

BSM penguin at high p

