

Resonance suppression from color reconnection

Based on Phys. Rev. D 97, 036010 (2018)

R. Acconcia¹, **D. D. Chinellato**¹, R. Derradi de Souza¹,
C. Markert², J. Takahashi¹, G. Torrieri¹


¹ - Universidade Estadual de Campinas

² - The University of Texas at Austin




UNICAMP

Introduction to resonances: the heavy ion view




nuclear collisions:
chemically equilibrated particle production

Introduction to resonances: the heavy ion view



- Description of hadron yields in a **thermally equilibrated regime**: same conclusions from different implementations, $T_{\text{ch}} \sim 156$ MeV
- dN/dy of many species in Pb-Pb well described: $\chi^2/\text{ndf} \sim 2$


Introduction to resonances: the heavy ion view



- However, the description is not good for the K^*
- Extra component [after hadronization](#): final state effect?




Resonances, Rescattering and Regeneration



- Consider the short-lived K^* resonance ($\sim 4 \text{ fm}/c$)
 - Usual scenario: decay in the hadronic phase
 - Daughters may **re-scatter** and yield may not be visible
 - **Regeneration (pseudo-elastic scattering)** may **recover** part of the yield
- Resonances: **probe** (the duration of) **the hadronic stage**

K^* and ϕ production rates in Pb-Pb



- Clear suppression observed in K^*/K^- when going from pp to central Pb-Pb collisions
- Not observed for ϕ/K^-
- Likely due to **re-scattering** of K^* decay daughters **with final-state hadronic medium**


$$\tau_{K^*} (\sim 4 \text{ fm}/c) \ll \tau_{\phi}$$

→ What about the smaller systems?

ALI-PREL-132043



K^* and ϕ production rates in Pb-Pb vs pp, p-Pb



- Clear suppression observed in K^*/K^- when going from pp to central Pb-Pb collisions: present vs multiplicity for **all systems**!
- Could there still be rescattering even in pp and p-Pb collisions?
 - **UrQMD says no**: system too small, would be negligible!

- Can we look at **alternative mechanisms** for this K^* yield reduction?

ALI-PREL-132043




Simulating pp: The PYTHIA event generator



- Monte Carlo event generator developed originally by the Lund group;
- Generates events based on
 - scattering amplitude from pQCD
 - additional empirical corrections
- Simulates the entire event, including hadronization, allowing for any experimental observable to be predicted
- Multi-purpose (e^+e^- , pp and more)


A proton-proton collision for PYTHIA



Parton momenta: p.d.f.

Select interacting **parton momenta** from p.d.f

A proton-proton collision for PYTHIA



Parton momenta: p.d.f.

Hard Parton interaction

Single Hard Parton Interaction


+ Resonance **Decays**

(N.B.: 'resonance' in this context:

"all particles with mass above the b quark system")



A proton-proton collision for PYTHIA



Parton momenta: p.d.f.


Hard Parton interaction

Initial state radiation (ISR)

Final state radiation (FSR)

Initial and Final State Radiation

A proton-proton collision for PYTHIA



Parton momenta: p.d.f.


Multiple interactions (MPI)

Initial state radiation (ISR)

Final state radiation (FSR)

Additional parton interactions included
Multiple interactions allowed to have **ISR** and **FSR**

A proton-proton collision for PYTHIA



Parton momenta: p.d.f.

Multiple interactions (MPI)

Initial state radiation (ISR)

Final state radiation (FSR)


Color (re)connection (CR)

Hadronization, decays

Final Partons of the Event
connected via **color confinement** strings
and **hadrons are created + decayed**

Multiple string
hadronization: non-trivial,
many approaches


Multiple string hadronization: the problem




- In PYTHIA: fragmentation via 'string breaking': flux tubes of interacting gluons of approximately constant energy density
- If MPIs available: which string arrangement should be used?

G. Gustafson, Acta Phys. Polon. **B40**, 1981 (2009)


"MPI-based" CR scheme



First Interaction




Second Interaction



Color reconnection


Multiple string hadronization: the problem



- In PYTHIA: **fragmentation via 'string breaking'**: flux tubes of interacting gluons of approximately constant energy density
- If **MPIs** available: **which string arrangement** should be used?

G. Gustafson, Acta Phys. Polon. **B40**, 1981 (2009)

"More QCD-based" CR scheme




Are there **other ways** to connect? → Yes! Introducing 'junctions'

...originally introduced to improve the description of **baryon production**

The K^*/K ratio in PYTHIA

p-Pb: Eur. Phys. J. C (2016) 76:245




- Single parton interaction: baseline $K^*/K \sim 0.33$
- Multiple interactions: higher N_{ch} , K^*/K at baseline value
- MPI-based CR: no significant change
- More-QCD based CR: signs of suppression?
- Effect within uncertainties of p-Pb (borderline)?

→ What is the origin of this suppression?...

...could this be due to particle decays?




The K^*/K ratio and the role of decays



- If all particles are allowed to decay, resonances will feed into the K (but not the K^*)
- Tested by looking at the K^*/K directly after hadronization (all decays disabled)
- Suppression still there!

Let's study the strings that produced these mesons...

String mass distribution



- For strings: $\text{length} \propto \text{energy}$
- CR produces shorter strings , especially in the More QCD-based scheme
- How does this depend on $\text{number of parton-parton interactions}$?


String mass distribution

- String lengths:

No CR > MPI-based


MPI-based > More QCD-based

- Effect is most important for high N_{PI} -> high multiplicity!




→ What about K^*/K for a given string mass?

The K^*/K vs string mass



- Vertical lines: averages for each CR scheme at $N_{\text{pl}} = 20$
- Suppression from lower average string mass

Other resonances: the ρ_0/π and η'/π



- ρ_0/π : also affected by rescattering in classical picture
 - Smaller strings also reduce this ratio!
- η'/π : interesting, η' is spin zero
 - Spin has nothing to do with this suppression

Conclusions

- Traditional heavy ion picture:
 - Short-lived resonances suppressed because of rescattering
 - From pp pQCD-inspired models:
 - They may also be suppressed in high-density hadronization because of shorter strings
 - This conclusion was drawn in a specific CR scheme but the mechanism is general.
-



- ❑ Could we use resonance suppression to tune CR? How much suppression could CR account for?
- ❑ What about baryonic resonances? Description harder in pQCD-inspired models, more parameter-dependent: more study needed
- ❑ How to distinguish experimentally? Is hadron mass or lifetime the driving factor in suppression measurements?

Thank you!



Backup


Other resonances: the ρ_0/π and ϕ/π



- ρ_0/π : also affected by rescattering in classical picture
→ Smaller strings also reduce this ratio!
- ϕ/π : no suppression in classical picture but beware different quark content, more investigation required

The K^*/K for $q\bar{q}$ strings and junctions

In the More QCD-based scheme





- General reduction of average mass for both cases
- At lower masses, junctions prefer forming baryons, leading to lower K^*/K ratios
- But: $q\bar{q}$ strings still responsible for majority of K , K^* production -> majority of suppression



3. Understanding of the late hadronic stage of the collision

K^*/K and the lifetime of the hadronic phase


- Model of Torrieri, Rafelski et al predicts particle ratios as a function of chemical freeze-out temperature and lifetime of hadronic phase
- Model predictions:



*References:

- G. Torrieri and J. Rafelski, *J. Phys. G* **28**, 1911 (2002)
- J. Rafelski et al., *Phys. Rev. C* **64**, 054907 (2001)
- J. Rafelski et al., *Phys. Rev. C* **65**, 069902(E) (2002)
- C. Markert et al., arXiv:hep-ph/0206260v2 (2002)

Baryon production: p/π



- Single parton interaction:
baseline $p/\pi \sim 0.06$
- Multiple interactions:
higher N_{ch} , no significant
change in ratio
- More QCD-based: p/π
increases, contrary to
data

pp: Nature Physics 13, 535–539 (2017)

p-Pb: Phys. Lett. B 728 (2014) 25–38

Pb-Pb: Phys. Rev. C 88, 044910 (2013)