


Overview of recent open heavy-flavour results with ALICE at the LHC

Xiaoming Zhang

- **Introduction**
- **R_{AA} and v_2 of open heavy-flavour particles**
- **Collectivity in smaller systems**
- **Charmed baryon production**
- **Open heavy flavour jets**

Introduction

Heavy quarks (charm and beauty): powerful probes of the Quark-Gluon Plasma (QGP)



Total charm cross section in A–A collisions is expected to scale w. r. t. the number of binary collisions in pp-like collisions

S. Radhakrishnan at QM'18

	Charm Hadron	Cross Section $d\sigma/dy$ (μb)
Au+Au 200 GeV (10-40%)	D^0	$41 \pm 1 \pm 5$
	D^+	$18 \pm 1 \pm 3$
	D_s^+	$15 \pm 1 \pm 5$
	Λ_c^+	$78 \pm 13 \pm 28^*$
	Total	$152 \pm 13 \pm 29$
p+p 200 GeV	Total	$130 \pm 30 \pm 26$

* derived using Λ_c^+ / D^0 ratio in 10-80% STAR Preliminary

- Produced in initial hard scatterings (high Q^2) at the early stage of heavy-ion collisions: $\tau_{c/b} \sim 0.01 - 0.1 \text{ fm}/c < \tau_{\text{QGP}} (\sim 0.3 \text{ fm}/c)$
- Production cross section calculable with pQCD ($m_c, m_b \gg \Lambda_{\text{QCD}}$)
- Experience the entire evolution of the QCD medium – probe transport properties of the deconfined medium

Introduction

Heavy quarks (charm and beauty): powerful probes of the Quark-Gluon Plasma (QGP)

Nuclear modification factor (R_{AA}): heavy quark in-medium energy loss


- Elastic (radiative) vs. inelastic (collisional) processes
- Radiative energy loss: color charge (Casimir factor) and mass (dead cone effect) dependence

$$R_{AA}(p_T) = \frac{dN_{AA}/dp_T}{\langle T_{AA} \rangle d\sigma_{pp}/dp_T} \begin{matrix} \text{QCD medium} \\ \text{QCD vacuum} \end{matrix}$$

- $R_{AA} = 1$, if no medium modification

$$\Delta E_g > \Delta E_q > \Delta E_c > \Delta E_b$$

→ $R_{AA}(\text{light hadron}) < R_{AA}(D) < R_{AA}(B)$?




Medium modification of heavy-flavour hadron formation

- Hadronization via quark coalescence which may modify the $D_s^+/\text{non-strange D}$ and Λ_c / D ratio

Introduction

Heavy quarks (charm and beauty): powerful probes of the Quark-Gluon Plasma (QGP)

$$E \frac{d^3\sigma}{d^3\vec{p}} = \frac{d^2\sigma}{2\pi p_T dp_T dy} [1 + \sum_{n=1}^{\infty} 2v_n \cos n(\varphi - \Psi_R)]$$



Azimuthal anisotropy: Fourier decomposition of particle azimuthal distribution relative to the reaction plane (Ψ_{RP})

- **Elliptic flow (v_2):** coefficient of second order harmonic
- Low and intermediate p_T : collective motion and possible heavy-quark thermalization in the QCD medium
- High p_T : path-length dependence of heavy-quark in-medium energy loss

Introduction

Open heavy-quark correlations and jets

- Complementary to open heavy-flavour measurements
- Possible modification of heavy-quark fragmentation
- Flavour dependence of the jet quenching / redistribution of the lost energy



Smaller systems: pp and p–Pb collisions


- Control experiments
 - Important to test pQCD calculations
 - Provide a necessary baseline for heavy-ion studies
 - Understanding of Cold Nuclear Matter (CNM) effects
- New collectivity-like effects observed at high multiplicity in smaller systems
 - Insight into Multiple-Parton-Interaction (MPI) phenomena
 - Understand the interplay of soft and hard processes

ALICE, Phys. Lett. B753 (2016) 126


ALICE
p-Pb $\sqrt{s_{NN}} = 5.02$ TeV
V0S: (0-20%)-(60-100%)

$0.5 < p_T^t$ (GeV/c) < 1
Assoc. tracklets

Trigger particle: muon at forward rapidity $2.5 < |\eta| < 4$
Associate particle: mid-rapidity tracklet $|\eta| < 1$



ALICE apparatus



Mid-rapidity ($|\eta| < 0.9$)

- ITS, TPC, TOF: vertexing, tracking, PID
- EMCal: high- p_T electron trigger, PID

Hadronic decays:

- $D^0 \rightarrow K^- \pi^+$
- $D^+ \rightarrow K^- \pi^+ \pi^+$
- $D^{*+} \rightarrow D^0 \pi^+$
- $D_s^+ \rightarrow \phi \pi^+ \rightarrow K^- K^+ \pi^+$

Semi-leptonic decays:

- $D, B \rightarrow e^+ X$
- $D, B \rightarrow \mu^+ X$

Forward MUON ($-4 < \eta < -2.5$)




- Muon trigger, tracking, PID

Smaller detectors: V0, T0, ZDC

- Event trigger, characterization

Nuclear modification factor




ALICE arXiv:1804.09083





- Increasing suppression from peripheral to central collisions
- R_{AA} of HFe at mid-rapidity is consistent with HFm at forward rapidity
 - Heavy quarks undergone significant interactions in a wide rapidity window in the most central Pb–Pb collisions
 - Confirmed the RUN-I measurements

Nuclear modification factor

ALICE arXiv:1804.09083




- Increasing suppression from peripheral to central collisions
- R_{AA} of HFe at mid-rapidity is consistent with HFm at forward rapidity



- Similar suppression at 5.02 TeV and 2.76 TeV
- Counterbalance between an increased medium temperature / density and harder quark p_T spectra

D meson R_{AA} vs. light-hadron R_{AA}

ALICE arXiv:1804.09083 ALICE arXiv:1802.09145




- $R_{AA}(D) > R_{AA}(\pi^\pm)$ for $p_T < 8$ GeV/c in central and semi-central collisions
- $R_{AA}(D) \approx R_{AA}(\pi^\pm) \approx R_{AA}(h^\pm)$ in peripheral collisions and for $p_T > 8$ GeV/c in central and semi-central collisions

D meson R_{AA} vs. light-hadron R_{AA}

ALICE arXiv:1804.09083



ALICE arXiv:1802.09145



- Double ratio at low p_T
 - N_{coll} vs N_{part} scaling
 - Different fragmentation and initial spectrum shapes
 - Possible mass and Casimir factor effects
 - Different coalescence and radial flow...
 - Expected by models and semi-central collisions

- $R_{AA}(D) \approx R_{AA}(\pi^\pm) \approx R_{AA}(h^\pm)$ in peripheral collisions and for $p_T > 8$ GeV/c in central and semi-central collisions



$R_{AA}(e \leftarrow b)$ vs. $R_{AA}(e \leftarrow c, b)$



- Hint of a smaller suppression for beauty-decay electrons for $p_T < 6$ GeV/c
- Data is reproduced by models within uncertainties, implementing quark mass dependent energy loss

Xe-Xe vs. Pb-Pb collisions

ALICE arXiv:1805.04399




- Similar R_{AA} observed with similar $\langle dN_{ch}/d\eta \rangle$ and N_{par}
 - Possible interplay of geometry and path-length dependence
- [arXiv:1805.04030]

Elliptic flow of open heavy flavours¹³

ALICE Phys. Rev. Lett. 120 (2018) 102301

ALICE arXiv:1809.09371




- Positive v_2 of D mesons for p_T in 2 – 8 GeV/c in semi-central collisions
- v_2 of D_s compatible with non-strange D-mesons within uncertainties

Elliptic flow of open heavy flavours

ALICE Phys. Rev. Lett. 120 (2018) 102301



ALICE arXiv:1809.09371

ALICE arXiv:1805.04390



- Positive v_2 of D mesons for p_T in 2 – 8 GeV/c in semi-central collisions
- v_2 of D_s compatible with non-strange D-mesons within uncertainties
- $v_2(D) \approx v_2(\pi^\pm)$ for $p_T > 4$ GeV/c, hint of $v_2(D) < v_2(\pi^\pm)$ for $p_T < 4$ GeV/c

R_{AA} and v_2 of D mesons vs. models





- ALICE LHC RUN-II: improved data precision w. r. t. RUN-I and provided important constraints on models
- Models in which charm quarks pick up **collective flow via recombination or subsequent elastic collisions** in expanding medium better describe both R_{AA} and v_2 at low p_T (**LBT**, **MC@sHQ**, **PHSD**, **POWLANG**)
- Recombination and collisional energy loss: important for heavy quarks
- Charm quark diffusion coefficient at the LHC: $(1.5 - 7) / 2\pi T_c$

Event-shape engineering

- Event eccentricity quantified by q_2 :

$$\rightarrow \langle (q_2)^2 \rangle \approx 1 + \langle M-1 \rangle \langle (v_2)^2 \rangle$$

- Opportunity to study the charm-quark coupling to the light-hadron bulk by measuring v_2 at different q_2 values





- Significant separation of D-meson v_2 in events with **large** and **small** q_2
- Charm quarks sensitive to the light-hadron bulk collectivity and event-by-event initial condition fluctuations

Autocorrelation and non-flow effects between q_2 determination and D-meson reconstruction are present

ALICE arXiv:1809.09371

Event-shape engineering

- Event eccentricity quantified by q_2 :
- $\langle (q_2)^2 \rangle \approx 1 + \langle M-1 \rangle \langle (v_2)^2 \rangle$
- Opportunity to study the charm-quark coupling to the light-hadron bulk by measuring v_2 at different q_2 values






ALICE arXiv:1809.09371

- $dN/dp_T(\text{ESE}) / dN/dp_T(\text{unbiased})$ compatible with unity within errors
- Promising observable to study interplay between elliptic flow and radial flow (at low/intermediate p_T) and in-medium energy (at high p_T)

Directed flow of open charm




- Sensitive to the early time EM fields in the collisions
 - Provide constraint for CME related physics
- Charm dragged by tilted bulk: production points are shifted from the bulk at $y \neq 0$ – larger v_1 for D mesons than for light flavours
 - Probe the longitudinal profile of the initial matter



Hint of positive slope with a significance of 2.7σ at low p_T

Directed flow of open charm

- Sensitive to the early time EM fields in the collisions
 - Provide constraint for CME related physics
- Charm dragged by tilted bulk: production points are shifted from the bulk at $y \neq 0$ – larger v_1 for D mesons than for light flavours
 - Probe the longitudinal profile of the initial matter




Hint of positive slope with a significance of 2.7σ at low p_T

Similar trend observed for charged particles, but different magnitude

HF-decay lepton ν_2 in p-Pb collisions²⁰

ALICE arXiv:1805.04367





ALI-PUB-310817

- Positive HFe ν_2 in $1.5 < p_T < 4$ GeV/c ($>5\sigma$) in high multiplicity events
 - Possible lower than ν_2 of charged particles at intermediate- p_T
 - Similar to inclusive muons at large rapidity

HF-decay lepton ν_2 in p-Pb collisions²¹

ALICE arXiv:1805.04367





ALI-PUB-310817

ALI-PREL-308084

- Positive HFe ν_2 in $1.5 < p_T < 4$ GeV/c ($>5\sigma$) in high multiplicity events
 - Possible lower than ν_2 of charged particles at intermediate- p_T
 - Similar to inclusive muons at large rapidity
- New: inclusive muon ν_2 at 8.16 TeV, Q-cumulants with 2-particle correlations
 - Positive ν_2 in $2 < p_T < 6$ GeV/c ($>3\sigma$) – HFm components dominated


Ξ_c production in pp collisions

ALICE Phys. Lett. B781 (2018) 8




- $\Xi_c(\rightarrow e \Xi^- \nu_e) / D^0$ ratio higher than theoretical predictions
 - PYTHIA8 with enhanced color reconnection mechanisms closer to data
 - $\text{BR}(\Xi_c \rightarrow e \Xi^- \nu_e)$ unknown, high uncertainty bands in theoretical predictions

Λ_c / D^0 ratio in smaller systems



- Decreasing trend from $p_T = 4 \text{ GeV}/c$ observed in pp and p-Pb collisions
- Similar trend to baryon-to-meson ratio in the light-flavour sector

Λ_c / D^0 ratio in smaller systems




- Similar trend to baryon-to-meson ratio in the light-flavour sector


served in pp and $p\text{-Pb}$ collisions

Λ_c production in Pb–Pb collisions

ALICE arXiv:1809.10922



ALI-PUB-311318




ALI-PUB-311322


- First measurement in Pb–Pb at the LHC
- Λ_c / D^0 ratio in Pb–Pb: higher than ($>2\sigma$) pp and p–Pb collisions
- Described by model calculations including only coalescence

Λ_c production in Pb–Pb collisions

ALICE arXiv:1809.10922



ALI-PUB-311314




ALI-PUB-311310

- First measurement in Pb–Pb at the LHC
- Hint of $R_{AA}(\Lambda_c) > R_{AA}(D_s) > R_{AA}(\text{non-strange } D) > R_{AA}(h^\pm)$
- A significant fraction of charm quarks hadronize via coalescence


D⁰-tagged jets R_{AA} in Pb–Pb collisions

- Strong suppression of D⁰-tagged jets in the most 10% central Pb–Pb collisions
- Hint of more suppression of low p_T D⁰-tagged jets than inclusive jets at higher p_T
- D⁰-tagged jets: more quark-seeded jets compared to inclusive jets



D⁰-tagged jets R_{AA} in Pb–Pb collisions

- Strong suppression of D⁰-tagged jets in the most 10% central Pb–Pb collisions
- Hint of more suppression of low p_T D⁰-tagged jets than inclusive jets at higher p_T
- D⁰-tagged jets: more quark-seeded jets compared to inclusive jets
- Similar suppression of D⁰-jets and D mesons



- New constraint on understanding charm quark in-medium energy loss

Conclusion

R_{AA} and v_2 of open heavy-flavour particles

- ALICE LHC RUN-II: improved data precision w. r. t. RUN-I
- Recombination and collisional energy loss: important for heavy quarks

Collectivity in smaller systems: positive v_2 in high multiplicity events

Charmed baryon production

- PYTHIA8 with enhanced color reconnection closer to data in smaller syst.
- Λ_c production in Pb–Pb: first measurement at the LHC
 - Described by model calculations including only coalescence
 - Suggests a significant fraction of charm hadronize via coalescence

Open heavy flavor tagged jets: similar suppression of D^0 -jets and D mesons

Thanks!