



# A proton imagining via Double Parton Scattering

Matteo Rinaldi<sup>1</sup>

<sup>1</sup>Dipartimento di Fisica e Geologia. Università degli studi di Perugia and INFN section of Perugia.

In collaboration with:

Federico Alberto Ceccopieri, Sergio Scopetta, Marco Traini and Vicente Vento



**DIPARTIMENTO DI  
FISICA E GEOLOGIA**

Università degli Studi di  
Perugia, Italia



# CONTENTS

01

## Introduction:

- 3D structure of the proton
- Double Parton Distribution Functions (dPDFs)
- Double parton correlations in dPDFs

03

## Analysis of correlations in dPDFs

M. R., S. Scopetta, M. Traini and V.Vento, JHEP 10, 063(2016)  
M. R., F. A. Ceccopieri, PRD 95, no. 3, 034040 (2017)

02

## dPDFs in constituent quark models, a hadron “imaging” via DPS?

M.R., S. Scopetta and V. Vento, PRD 87, 114021 (2013)  
M.R., S. Scopetta, M. Traini and V.Vento, JHEP 12, 028 (2014)  
M.R., S. Scopetta, M. Traini and V.Vento, EPJC 781 (2018)  
M. R. and F. A. Ceccopieri, PRD 97, no. 7, 071501 (2018)  
M. R. and F. A. Ceccopieri. Accepted in JHEP, arXiv: 1812.04286.

02

## Calculation of experimental observables: effects of correlations

M. R., S. Scopetta, M. Traini and V.Vento, PLB 752, 40 (2016)  
M. Traini, S. Scopetta, M. R. and V. Vento, PLB 768, 270 (2017)  
F. A. Ceccopieri, M. R., S. Scopetta, PRD 95, 114030(2017)  
M. R. and F. A. Ceccopieri, Accepted in JHEP arXiv: 1812.04286

# CONTENTS

01

## Introduction:

- 3D structure of the proton
- Double Parton Distribution Functions (dPDFs)
- Double parton correlations in dPDFs

03

## Analysis of correlations in dPDFs

M. R., S. Scopetta, M. Traini and V.Vento, JHEP 10, 063(2016)  
M. R., F. A. Ceccopieri, PRD 95, no. 3, 034040 (2017)

02

## dPDFs in constituent quark models, a hadron “imaging” via DPS?

M.R., S. Scopetta and V. Vento, PRD 87, 114021 (2013)  
M.R., S. Scopetta, M. Traini and V.Vento, JHEP 12, 028 (2014)  
M.R., S. Scopetta, M. Traini and V.Vento, EPJC 781 (2018)  
M. R. and F. A. Ceccopieri, PRD 97, no. 7, 071501 (2018)  
M. R. and F. A. Ceccopieri. Accepted in JHEP, arXiv: 1812.04286.

02

## Calculation of experimental observables: effects of correlations

M. R., S. Scopetta, M. Traini and V.Vento, PLB 752, 40 (2016)  
M. Traini, S. Scopetta, M. R. and V. Vento, PLB 768, 270 (2017)  
F. A. Ceccopieri, M. R., S. Scopetta, PRD 95, 114030(2017)  
M. R. and F. A. Ceccopieri, Accepted in JHEP arXiv: 1812.04286

# CONTENTS

01

## Introduction:

- 3D structure of the proton
- Double Parton Distribution Functions (dPDFs)
- Double parton correlations in dPDFs

03

## Analysis of correlations in dPDFs

M. R., S. Scopetta, M. Traini and V.Vento, JHEP 10, 063(2016)  
M. R., F. A. Ceccopieri, PRD 95, no. 3, 034040 (2017)

02

## dPDFs in constituent quark models, a hadron “imaging” via DPS?

M.R., S. Scopetta and V. Vento, PRD 87, 114021 (2013)  
M.R., S. Scopetta, M. Traini and V.Vento, JHEP 12, 028 (2014)  
M.R., S. Scopetta, M. Traini and V.Vento, EPJC 781 (2018)  
M. R. and F. A. Ceccopieri, PRD 97, no. 7, 071501 (2018)  
M. R. and F. A. Ceccopieri. Accepted in JHEP, arXiv: 1812.04286.

02

## Calculation of experimental observables: effects of correlations

M. R., S. Scopetta, M. Traini and V.Vento, PLB 752, 40 (2016)  
M. Traini, S. Scopetta, M. R. and V. Vento, PLB 768, 270 (2017)  
F. A. Ceccopieri, M. R., S. Scopetta, PRD 95, 114030(2017)  
M. R. and F. A. Ceccopieri, Accepted in JHEP arXiv: 1812.04286



# CONTENTS

01

## Introduction:

- 3D structure of the proton
- Double Parton Distribution Functions (dPDFs)
- Double parton correlations in dPDFs

03

## Analysis of correlations in dPDFs

M. R., S. Scopetta, M. Traini and V.Vento, JHEP 10, 063(2016)  
M. R., F. A. Ceccopieri, PRD 95, no. 3, 034040 (2017)

02

## dPDFs in constituent quark models, a hadron “imaging” via DPS?


M.R., S. Scopetta and V. Vento, PRD 87, 114021 (2013)  
M.R., S. Scopetta, M. Traini and V.Vento, JHEP 12, 028 (2014)  
M.R., S. Scopetta, M. Traini and V.Vento, EPJC 781 (2018)  
M. R. and F. A. Ceccopieri, PRD 97, no. 7, 071501 (2018)  
M. R. and F. A. Ceccopieri. Accepted in JHEP, arXiv: 1812.04286.

02


## Calculation of experimental observables: effects of correlations

M. R., S. Scopetta, M. Traini and V.Vento, PLB 752, 40 (2016)  
M. Traini, S. Scopetta, M. R. and V. Vento, PLB 768, 270 (2017)  
F. A. Ceccopieri, M. R., S. Scopetta, PRD 95, 114030(2017)  
M. R. and F. A. Ceccopieri, Accepted in JHEP arXiv: 1812.04286


# THE 3D STRUCTURE OF THE PROTON



The 3D structure of a strongly interacting system (e.g. nucleon, nucleus...) could be accessed through different processes (e.g. SIDIS, DVCS ...), measuring different kind of parton distributions, providing different kind of information. The parton distribution puzzle is:



# THE 3D STRUCTURE OF THE PROTON



The 3D structure of a strongly interacting system (e.g. nucleon, nucleus...) could be accessed through different processes (e.g. SIDIS, DVCS ...), measuring different kind of parton distributions, providing different kind of information. The parton distribution puzzle is:


All these distributions are ONE-BODY functions!

How can we access new information as two particle correlations?




Theoretical tools

« Physical » objects




[Lorce, Pasquini, Vanderhaeghen (2011)]

# Answer: MULTIPARTON INTERACTIONS



Multiparton interaction (MPI) can contribute to the, pp and pA, cross section @ the LHC:



The cross section for a double parton scattering (DPS) event can be written in the following way:

N. Paver, D. Treleani, *Nuovo Cimento* 70A, 215 (1982)

$$d\sigma = \frac{1}{S} \sum_{i,j,k,l} \hat{\sigma}_{ij}(\mathbf{x}_1, \mathbf{x}_3, \mu_A) \hat{\sigma}_{kl}(\mathbf{x}_2, \mathbf{x}_4, \mu_B) \int d^2z_{\perp} \underbrace{F_{ik}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{z}_{\perp}, \mu_A, \mu_B)}_{\text{dPDF}} \underbrace{F_{jl}(\mathbf{x}_3, \mathbf{x}_4, \mathbf{z}_{\perp}, \mu_A, \mu_B)}_{\text{Momentum scales}}$$


Momentum fractions carried by the parton inside the proton

Transverse distance between partons

DPS processes are important for fundamental studies, e.g. the background for the research of new physics and to grasp information on the 3D PARTONIC STRUCTURE OF THE PROTON






# Answer: MULTIPARTON INTERACTIONS



1-body





2-body



LFWFs



Phase-space (Wigner) distribution



dPDFs  
 $x_1, x_2, \vec{z}_\perp$

TMDs  
 $x, \vec{k}_\perp$

GTMDs  
 $x, \vec{k}_\perp, \Delta$

PDFs  
 $x$

FFs  
 $\Delta$

Charges


GPDs  
 $x, \Delta$

$\Delta = 0$   
 $\int dx$   
 $\int d^2k_\perp$

$\int d^2z_\perp \int dx_2$



# Parton correlations and dPDFs



@ LHC kinematics it is often used a factorized form of the **dPDFs**:  $(\mathbf{x}_1, \mathbf{x}_2) - \mathbf{z}_\perp$  factorization:


$$F_{ij}(x_1, x_2, \vec{z}_\perp, \mu) = F_{ij}(x_1, x_2, \mu) T(\vec{z}_\perp, \mu) \quad \text{and } x_1, x_2 \text{ factorization:}$$

\* Here and in the following:  
 $\mu = \mu_A = \mu_B$

$$F_{ij}(x_1, x_2, \mu) = \underbrace{q_i(x_1, \mu)}_{\text{PDF (1-Body)}} \underbrace{q_j(x_2, \mu)}_{\text{PDF (1-Body)}} \theta(1 - x_1 - x_2) (1 - x_1 - x_2)^n$$

unknown

**NO CORRELATION ANSATZ**



In this scenario, parton correlations inside the proton are neglected



**NO NEW INFORMATION!**

BUT:

- Correlations are present
- dPDFs** are non perturbative in QCD and DPCs cannot be directly evaluated within QCD


**HOW CAN WE BE SURE OF THE ACCURACY OF SUCH APPROXIMATION**



**WHAT CAN WE LEARN ABOUT dPDFs AND THE PROTON STRUCTURE?**



# DPCs in Constituent quark models (CQMs)



- Main features:
  - Effective potential
  - Effective particles strongly bound and **correlated**

CQM are a proper framework to describe DPCs, but their predictions are reliable **ONLY** in the valence quark region at low energy scale

**WHILE**

LHC data are available at small  $\mathcal{Q}$  in this region, due to the large population of partons, the role of correlations could be less relevant **BUT** theoretical microscopic estimates are necessary!

i) dPDF evaluated at the initial scale of the model


**pQCD evolution of dPDFs**

ii) dPDF evaluated at high generic scale

CQM calculations are useful tools for the interpretation of data and for the planning of measurements of unknown quantities (e.g., TMDs in SiDIS, GPDs in DVCS...)

**Similar expectations motivate the present investigation of**  
**dPDFs**

# The Light-Front approach



Relativity can be implemented, for a CQM, by using a **Light-Front** (LF) approach. In the Relativistic Hamiltonian Dynamics (RHD) of an interacting system, introduced by Dirac (1949), one has:

{

• RHD
 


Instant Form:  $t_0=0$

Evolution Operator:  $P_0= E$

**Front Form (LF):**  $x_+ = t_0+z=0$

**Evolution Operator:**  $P_-$

$a^\pm = a_0 \pm a_3$


• Fixed number of off-shell particles
  




• Full Poincare' covariance

• 7 Kinematical generators (maximum number): i) three LF boosts (at variance with the dynamical nature of the Instant-form boosts), ii)  $P^+$ ,  $P_\perp$ , iii) Rotation around z.

• The proton state can be represented in the following way:

see e.g.: S. J. Brodsky, H. -C. Pauli, S. S. Pinsky, Phys.Rept. 301, 299 (1998)


$$|p, P^+ \vec{P}_\perp\rangle = \psi_{qqq} |qqq\rangle + \psi_{qqq g} |qqq g\rangle + \psi_{qqq q\bar{q}} |qqq q\bar{q}\rangle$$



$\psi_n =$  LF wave function

Invariant under LF boosts

# The Light-Front approach



Relativity can be implemented, for a CQM, by using a **Light-Front** (LF) approach. In the Relativistic Hamiltonian Dynamics (RHD) of an interacting system, introduced by Dirac (1949), one has:

{

RHD


Instant Form:  $t_0=0$

Evolution Operator:  $P_0= E$

**Front Form (LF):**  $x_+ = t_0+z=0$

**Evolution Operator:**  $P_-$

$a^\pm = a_0 \pm a_3$






- Fixed number of off-shell particles
- Full Poincare' covariance

• 7 Kinematical generators (maximum number): i) three LF boosts (at variance with the dynamical nature of the Instant-form boosts), ii)  $\mathbf{P}^+$ ,  $\mathbf{P}_\perp$ , iii) Rotation around z.

• The proton state can be represented in the following way:

see e.g.: S. J. Brodsky, H. -C. Pauli, S. S. Pinsky, *Phys.Rept.* 301, 299 (1998)


$$|p, P^+ \vec{P}_\perp\rangle = \psi_{qqq} |qqq\rangle + \psi_{qqq g} |qqq g\rangle + \psi_{qqq q\bar{q}} |qqq q\bar{q}\rangle$$



$\psi_n =$  LF wave function

Invariant under LF boosts

# The Light-Front approach



Relativity can be implemented, for a CQM, by using a **Light-Front** (LF) approach. In the Relativistic Hamiltonian Dynamics (RHD) of an interacting system, introduced by Dirac (1949), one has:

RHD {

- Instant Form:  $t_0=0$
- Evolution Operator:  $P_0= E$
- Front Form (LF):**  $x_+ = t_0+z=0$
- Evolution Operator:**  $P_-$

$a^\pm = a_0 \pm a_3$


- Fixed number of off-shell particles
- Full Poincare' covariance

7 Kinematical generators (maximum number): i) three LF boosts (at variance with the dynamical nature of the Instant-form boosts), ii)  $\mathbf{P}^+$ ,  $\mathbf{P}_\perp$ , iii) Rotation around z.


The proton state can be represented in the following way:

see e.g.: **S. J. Brodsky, H. -C. Pauli, S. S. Pinsky, Phys.Rept. 301, 299 (1998)**


$$|p, P^+ \vec{P}_\perp\rangle = \psi_{qqq} |qqq\rangle + \psi_{qqq g} |qqq g\rangle + \psi_{qqq q\bar{q}} |qqq q\bar{q}\rangle$$



LC2019



Matteo Rinaldi




$\psi_n =$  **LF wave function**

Invariant under LF boosts

# dPDFs in a Light-Front approach

M. R., S. Scopetta, M. Traini and V.Vento, JHEP 12, 028 (2014)



Extending the procedure developed in S. Boffi, B. Pasquini and M. Traini, Nucl. Phys. B 649, 243 (2003) for GPDs, we obtained the following expression of the **dPDF** in momentum space, often called  **$_2$ GPDs**:

$$F_{ij}(x_1, x_2, k_\perp) = 3(\sqrt{3})^3 \int \prod_{i=1}^3 d\vec{k}_i \delta\left(\sum_{i=1}^3 \vec{k}_i\right) \Phi^*({\vec{k}_i}, k_\perp) \Phi({\vec{k}_i}, -k_\perp) \\ \times \delta\left(x_1 - \frac{k_1^+}{P_+}\right) \delta\left(x_2 - \frac{k_2^+}{P_+}\right)$$

$\Phi({\vec{k}_i}, \pm k_\perp) = \Phi\left(\vec{k}_1 \pm \frac{\vec{k}_\perp}{2}, \vec{k}_2 \mp \frac{\vec{k}_\perp}{2}, \vec{k}_3\right)$

Conjugate to  $z_\perp$

**GOOD SUPPORT**

$$x_1 + x_2 > 1 \Rightarrow F_{ij}(x_1, x_2, k_\perp) = 0$$


$$\Phi(\vec{k}_1, \vec{k}_2, \vec{k}_3) = D^{\dagger 1/2}(R_{il}(\vec{k}_1)) D^{\dagger 1/2}(R_{il}(\vec{k}_2)) D^{\dagger 1/2}(R_{il}(\vec{k}_3)) \psi^{[i]}(\vec{k}_1, \vec{k}_2, \vec{k}_3)$$

**Melosh operator rotates canonical spin in LF one**

**Instant form** proton w.f.  
**We need a CQM!**




# What we would like to learn: partonic mean distance



M. R. and F. A. Ceccopieri, arXiv: 1812.04286, JHEP accepted

Since, in coordinates space, dPDFs get a number density interpretation, in principle one can calculate the mean distance between partons!

For example, for 2 gluons perturbatively generated:




1) **HP** model

2) **HO** model

$$x_1 = 10^{-4} \text{ and } x_2 = 10^{-2}$$

$$\vec{d}_\perp = \vec{b}_\perp = \vec{z}_\perp$$



M. Traini *et al*, Nucl. Phys. A 656, 400-420 (1999), non relativistic Hyper-Central CQM (potential by M. Ferraris *et al*, PLB 364 (1995)) (**HP**)

The harmonic oscillator (**HO**)


One can also define the mean transverse distance  $(x_1 - x_2)$  distribution as follows:

$$\langle d_\perp^2 \rangle_{x_1, x_2}^{ij} = \frac{\int d^2 b_\perp b_\perp^2 F_{ij}(x_1, x_2, b_\perp, Q^2 = M_W^2)}{\int d^2 b_\perp F_{ij}(x_1, x_2, b_\perp, Q^2 = M_W^2)}$$

For example, for 2 gluons and two different models, one gets:

$$\begin{aligned} \sqrt{\langle d_\perp^2 \rangle_{10^{-2}, 10^{-2}}} &\begin{cases} \rightarrow 0.404 \text{ fm HP} \\ \rightarrow 0.365 \text{ fm HO} \end{cases} \\ \sqrt{\langle d_\perp^2 \rangle_{10^{-4}, 10^{-4}}} &\begin{cases} \rightarrow 0.391 \text{ fm HP} \\ \rightarrow 0.393 \text{ fm HO} \end{cases} \end{aligned}$$



# What we would like to learn: partonic mean distance



M. R. and F. A. Ceccopieri, arXiv: 1812.04286, JHEP accepted

Since, in coordinates space, dPDFs get a number density interpretation, in principle one can calculate the mean distance between partons!

For example, for 2 gluons perturbatively generated:





1) **HP** model

2) **HO** model

$$x_1 = 10^{-4} \text{ and } x_2 = 10^{-2}$$

$$\vec{d}_\perp = \vec{b}_\perp = \vec{z}_\perp$$



M. Traini *et al*, Nucl. Phys. A 656, 400-420 (1999), non relativistic Hyper-Central CQM (potential by M. Ferraris *et al*, PLB 364 (1995)) (**HP**)

The harmonic oscillator (**HO**)

One can also define the mean transverse distance  $(x_1 - x_2)$  distribution as follows:


$$\langle d_\perp^2 \rangle_{x_1, x_2}^{ij} = \frac{\int d^2 b_\perp b_\perp^2 F_{ij}(x_1, x_2, b_\perp, Q^2 = M_W^2)}{\int d^2 b_\perp F_{ij}(x_1, x_2, b_\perp, Q^2 = M_W^2)}$$

**Are two slow partons closer (in  $\perp$  plane) than two fast partons?**





# What we learned:

M. R., S. Scopetta, M. Traini and V.Vento, JHEP 12, 028 (2014)




Ratios, sensitive to correlations, are shown in order to test the factorization ansatz! Use has been made of relativistic HP CQM.



The  $(x_1, x_2) - k_\perp$  and  $x_1 - x_2$  factorizations are **violated** in **all quark model analyses!**

M.R., S. Scopetta and V. Vento, PRD 87, 114021 (2013), H.-M. Chang, A.V. Manohar, and W.J. Waalewijn, PRD 87, 034009 (2013)

# What we learned: a link between dPDFs and GPDs?



The **dPDF** is formally defined through the Light-cone correlator:


$$F_{12}(x_1, x_2, \vec{z}_\perp) \propto \left( \sum_X \right) \int dz^- \left[ \prod_{i=1}^2 dl_i^- e^{ix_i l_i^- p^+} \right] \langle p | O(z, l_1) | X \rangle \langle X | O(0, l_2) | p \rangle \Big|_{\substack{\vec{l}_{1\perp} = \vec{l}_{2\perp} = 0 \\ l_1^+ = l_2^+ = z^+ = 0}}$$

Approximated by the proton state!

$$\int \frac{dp'^+ d\vec{p}'_\perp}{p'^+} |p'\rangle \langle p'|$$

**GPDS**

$$F_{12}(x_1, x_2, \vec{k}_\perp) \sim \underbrace{f(x_1, 0, \vec{k}_\perp)}_{\text{dPDF}} \underbrace{f(x_2, 0, \vec{k}_\perp)}_{\text{GPD}}$$



..... dPDF = GPD x GPD


— dPDF

In GPDs, the variables  $k_\perp$  and  $x$  are correlated!



Correlations between  $\vec{z}_\perp$  and  $x_1, x_2$  could be present in **dPDFs** !

# The Effective X-section



A fundamental tool for the comprehension of the role of DPS in hadron-hadron collisions is the so called “effective X-section”.

This object can be defined through a “pocket formula”:

Sensitive to correlations

$$\sigma_{eff} = \frac{m}{2} \frac{\sigma_A^{pp'} \sigma_B^{pp'}}{\sigma_{double}^{pp}}$$

Differential cross section for the process:  $pp' \rightarrow A(B) + X$


Combinatorial factor

Differential cross section for a DPS event:  $pp' \rightarrow A + B + X$

## ....EXPERIMENTAL STATUS:

- Difficult extraction, approved analysis for the same
- sign  $W$ 's production @LHC (RUN 2)
- the model dependent extraction of  $\sigma_{eff}$  from data is almost consistent with a “constant” (within errors) (**uncorrelated ansatz usually assumed!**)
- different ranges in  $X_i$  accessed in different



*Within our CQM framework, we can calculate  $\sigma_{eff}$  without any approximations!*



Yamanaka's talk  
mpi@LHC (2018)

# Same sign $W$ 's in pp collisions at $\sqrt{s} = 13$ TeV at the LHC

F. A. Ceccopieri, M. R., S. Scopetta, Phys.Rev. D95 (2017) no.11, 114030





In this channel, the single parton scattering (usually dominant w.r.t to the double one) starts to contribute to higher order in strong coupling constant.



*“Same-sign  $W$  boson pairs production is a promising channel to look for signature of double Parton interactions at the LHC.”*

# Same sign $W$ 's in pp collisions at $\sqrt{s} = 13$ TeV at the LHC

F. A. Ceccopieri, M. R., S. Scopetta, Phys.Rev. D95 (2017) no.11, 114030




In this channel, the single parton scattering (usually dominant w.r.t to the double one) starts to contribute to higher order in strong coupling constant.




**Can double parton correlations be observed for the first time in the next LHC run ?**

# Same sign W's in pp collisions at $\sqrt{s} = 13$ TeV at the LHC



F. A. Ceccopieri, M. R., S. Scopetta, Phys.Rev. D95 (2017) no.11, 114030



## Kinematical cuts

$$\begin{aligned}
 & pp, \sqrt{s} = 13 \text{ TeV} \\
 & p_{T,\mu}^{\text{leading}} > 20 \text{ GeV}, \quad p_{T,\mu}^{\text{subleading}} > 10 \text{ GeV} \\
 & |p_{T,\mu}^{\text{leading}}| + |p_{T,\mu}^{\text{subleading}}| > 45 \text{ GeV} \\
 & |\eta_\mu| < 2.4 \\
 & 20 \text{ GeV} < M_{\text{inv}} < 75 \text{ GeV} \text{ or } M_{\text{inv}} > 105 \text{ GeV}
 \end{aligned}$$

## DPS cross section:

$$\frac{d^4\sigma_{pp \rightarrow \mu^\pm \mu^\pm X}}{d\eta_1 dp_{T,1} d\eta_2 dp_{T,2}} = \sum_{i,k,j,l} \frac{1}{2} \int d^2\vec{b}_\perp F_{ij}(x_1, x_2, \vec{b}_\perp, M_W) F_{kl}(x_3, x_4, \vec{b}_\perp, M_W) \frac{d^2\sigma_{ik}^{pp \rightarrow \mu^\pm X}}{d\eta_1 dp_{T,1}} \frac{d^2\sigma_{jl}^{pp \rightarrow \mu^\pm X}}{d\eta_2 dp_{T,2}} \mathcal{I}(\eta_i, p_{T,i})$$


$M_W \longrightarrow$  Momentum scale

In order to estimate the role of double parton correlations we have used as input of dPDFs:


1) Longitudinal and transverse correlations arise from the relativistic CQM model describing three valence quarks

2) These correlations propagate to sea quarks and gluons through pQCD evolution

# Same sign W's in pp collisions at $\sqrt{s} = 13$ TeV at the LHC



F. A. Ceccopieri, M. R., S. Scopetta, Phys.Rev. D95 (2017) no.11, 114030



## Kinematical cuts

$$\begin{aligned}
 & pp, \sqrt{s} = 13 \text{ TeV} \\
 & p_{T,\mu}^{\text{leading}} > 20 \text{ GeV}, \quad p_{T,\mu}^{\text{subleading}} > 10 \text{ GeV} \\
 & |p_{T,\mu}^{\text{leading}}| + |p_{T,\mu}^{\text{subleading}}| > 45 \text{ GeV} \\
 & |\eta_\mu| < 2.4 \\
 & 20 \text{ GeV} < M_{\text{inv}} < 75 \text{ GeV} \text{ or } M_{\text{inv}} > 105 \text{ GeV}
 \end{aligned}$$

## DPS cross section:

$$\frac{d^4 \sigma_{pp \rightarrow \mu^\pm \mu^\pm X}}{d\eta_1 dp_{T,1} d\eta_2 dp_{T,2}} = \sum_{i,k,j,l} \frac{1}{2} \int d^2 \vec{b}_\perp F_{ij}(x_1, x_2, \vec{b}_\perp, M_W) F_{kl}(x_3, x_4, \vec{b}_\perp, M_W) \frac{d^2 \sigma_{ik}^{pp \rightarrow \mu^\pm X}}{d\eta_1 dp_{T,1}} \frac{d^2 \sigma_{jl}^{pp \rightarrow \mu^\pm X}}{d\eta_2 dp_{T,2}} \mathcal{I}(\eta_i, p_{T,i})$$

$M_W \longrightarrow$  Momentum scale


In order to estimate the role of double parton correlations we have used as input of dPDFs:

Relativistic model: **QM** M. R., S. Scopetta, M. Traini and V.Vento, JHEP 12, 028 (2014)

Final Results:  $\sigma^{++} + \sigma^{--} [\text{fb}] \sim 0.69 \pm 0.18 (\delta\mu_F)_{-0.16}^{+0.12} (\delta Q_0)^*$




# Same sign W's in pp collisions at $\sqrt{s} = 13$ TeV at the LHC




F. A. Ceccopieri, M. R., S. Scopetta, Phys.Rev. D95 (2017) no.11, 114030

In order to understand whether correlations can be accessed in experimental observations, using dPDF evaluated within the QM model, the effective cross section has been calculated for this process and compared with its mean value:



$$\eta_1 \cdot \eta_2 \simeq \frac{1}{4} \ln \frac{x_1}{x_3} \ln \frac{x_2}{x_4}$$

$$\langle \tilde{\sigma}_{eff} \rangle = 21.04^{+0.07}_{-0.07} (\delta Q_0) {}^{+0.06}_{-0.07} (\delta \mu_F) \text{ mb} .$$



$$\tilde{\sigma}_{eff} = \frac{m}{2} \frac{\sigma_A^{pp'} \sigma_B^{pp'}}{\sigma_{double}^{pp}}$$


Difference between green and red line is due to correlations effects

x- dependence of effective x-section consistent with analyses:  
 M.Rinaldi et al PLB 752,40 (2016)  
 M. Traini, M. R. et al, PLB 768, 270 (2017)

“Assuming that the results of the first and the last bins can be distinguished if they differ by 1 sigma, we estimated that


$\mathcal{L} = 1000 \text{ fb}^{-1}$  is necessary to observe correlations”

# Same sign W's in pp collisions at $\sqrt{s} = 13$ TeV at the LHC




F. A. Ceccopieri, M. R., S. Scopetta, Phys.Rev. D95 (2017) no.11, 114030

In order to understand whether correlations can be accessed in experimental observations, using dPDF evaluated within the QM model, the effective cross section has been calculated for this process and compared with its mean value:




$$\eta_1 \cdot \eta_2 \simeq \frac{1}{4} \ln \frac{x_1}{x_3} \ln \frac{x_2}{x_4}$$

$$\langle \tilde{\sigma}_{eff} \rangle = 21.04^{+0.07}_{-0.07} (\delta Q_0) {}^{+0.06}_{-0.07} (\delta \mu_F) \text{ mb} .$$



$$\tilde{\sigma}_{eff} = \frac{m}{2} \frac{\sigma_A^{pp'} \sigma_B^{pp'}}{\sigma_{double}^{pp}}$$


Difference  between green and red line is due to correlations effects

To observe correlations,  $\mathcal{L} = 1000 \text{ fb}^{-1}$  is needed!



REACHABLE IN THE PLANNED LHC RUN

# Same sign W's in pp collisions at $\sqrt{s} = 13$ TeV at the LHC




F. A. Ceccopieri, M. R., S. Scopetta, Phys.Rev. D95 (2017) no.11, 114030


In order to understand whether correlations can be accessed in experimental observations, using dPDF evaluated within the QM model, the effective cross section has been calculated for this process and compared with its mean value:


$$\langle \tilde{\sigma}_{eff} \rangle = 21.04^{+0.07}_{-0.07} (\delta Q_0) {}^{+0.06}_{-0.07} (\delta \mu_F) \text{ mb} .$$

$$\tilde{\sigma}_{eff} = \frac{m}{2} \frac{\sigma_A^{pp'} \sigma_B^{pp'}}{\sigma_{double}^{pp}}$$




$$\eta_1 \cdot \eta_2 \simeq \frac{1}{4} \ln \frac{x_1}{x_3} \ln \frac{x_2}{x_4}$$



Difference  between green and red line is due to correlations effects

**IN THIS CHANNEL, WE ESTABLISHED THE POSSIBILITY TO OBSERVE, FOR THE FIRST TIME, TWO-PARTON CORRELATIONS IN THE NEXT LHC RUN!**

# A clue from data?




M. R. and F. A. Ceccopieri, PRD 97, no. 7, 071501 (2018) rapid communication

Considering the factorization ansatz, for which some estimates of  $\sigma_{\text{eff}}$  are available, one has:

$$\sigma_{eff} = \left[ \int \frac{d\vec{k}_{\perp}}{(2\pi)^2} \tilde{T}(\vec{k}_{\perp}) \tilde{T}(-\vec{k}_{\perp}) \right]^{-1}$$

Effective form factor (Eff)

# A clue from data?



M. R. and F. A. Ceccopieri, PRD 97, no. 7, 071501 (2018) rapid communication

Considering the factorization ansatz, for which some estimates of  $\sigma_{\text{eff}}$  are available, one has:  $\sigma_{\text{eff}} = \left[ \int \frac{d\vec{k}_{\perp}}{(2\pi)^2} \tilde{T}(\vec{k}_{\perp}) \tilde{T}(-\vec{k}_{\perp}) \right]^{-1}$  Effective form factor (Eff)


Eff can be formally defined as **FIRST MOMENT** of dPDF (like for GPDs) through the proton wave function:

$$\tilde{T}(k_{\perp}) = \frac{1}{2} \int dx_1 dx_2 F(x_1, x_2, k_{\perp}) = \int d\vec{k}_1 d\vec{k}_2 \Psi(\vec{k}_1 + \vec{k}_{\perp}, \vec{k}_2) \Psi^{\dagger}(\vec{k}_1, \vec{k}_2 + \vec{k}_{\perp})$$

From the above quantity the mean distance in the transverse plane between two partons can be defined:

$$\langle b^2 \rangle \sim -2 \frac{d}{k_{\perp} dk_{\perp}} \tilde{T}(k_{\perp}) \Big|_{k_{\perp}=0}$$

# A clue from data?



M. R. and F. A. Ceccopieri, PRD 97, no. 7, 071501 (2018) rapid communication

Considering the factorization ansatz, for which some estimates of  $\sigma_{eff}$  are available, one has:  $\sigma_{eff} = \left[ \int \frac{d\vec{k}_\perp}{(2\pi)^2} \tilde{T}(\vec{k}_\perp) \tilde{T}(-\vec{k}_\perp) \right]^{-1}$  Effective form factor (Eff)

Eff can be formally defined as **FIRST MOMENT** of dPDF (like for GPDs) through the proton wave function:

$$\tilde{T}(k_\perp) = \frac{1}{2} \int dx_1 dx_2 F(x_1, x_2, k_\perp) = \int d\vec{k}_1 d\vec{k}_2 \Psi(\vec{k}_1 + \vec{k}_\perp, \vec{k}_2) \Psi^\dagger(\vec{k}_1, \vec{k}_2 + \vec{k}_\perp)$$

From the above quantity the mean distance in the transverse plane between two partons can be defined:


$$\langle b^2 \rangle \sim -2 \frac{d}{k_\perp dk_\perp} \tilde{T}(k_\perp) \Big|_{k_\perp=0}$$

Eff is unknown but using general model independent properties and comparing Eff with standard proton ff, we found:

$$\frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi}$$

DPS processes:


The vertical line stands for the transverse proton radius



We also:

M. R. and F. A. Ceccopieri, arXiv: 1812.04286. JHEP accepted


- Extended the approach including splitting term



- Extended the approach to the most general unfactorized case




# Some extensions of the relation : $\frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi}$ |



M. R. and F. A. Ceccopieri, arXiv: 1812.04286. JHEP accepted.

Relevant for processes involving heavy particles is the splitting term (1v2 mechanism) i.e.:

$$* D_{j_1, j_2}(x_1, x_2) = \int d^2 b_{\perp} \tilde{F}_{j_1, j_2}(x_{1,2}, b_{\perp})$$



In pQCD evolution:

$$\frac{dD_{j_1 j_2}(x, x_2; t)}{dt} =$$

Gaunt J.R. and Stirling W. J., JHEP 03 (2010)

$$= \left\{ \begin{array}{l} \text{Homogeneous term (double DGLAP)} \\ + \\ \sum_{j'} F_{j'}(x_1 + x_2; t) \underbrace{\frac{1}{x_1 + x_2} P_{j' \rightarrow j_1 j_2}}_{\text{SPLITTING TERM}} \left( \frac{x_1}{x_1 + x_2} \right) \end{array} \right.$$

1v2

2v2

**SPLITTING TERM**



$$\frac{\sigma_{eff}}{3\pi} \left( 1 + \frac{3}{2} r_v \right) \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi} \left( 1 + 2 r_v \right)$$

$$r_v \sim \frac{F_{j_1 j_2}^{splitting}(x_1, x_2, k_{\perp} = 0; t)}{F_{j_1 j_2}(x_1, x_2, k_{\perp} = 0; t)}$$

Due to the difficulty in the estimate of the 2 contributions:

with:  
 $0 \leq r_v \leq 1$

Absolute minimum  
 $r_v = 0$


$$\frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq$$

$$\frac{3 \sigma_{eff}}{\pi}$$

Absolute maximum  
 $r_v = 1$




# Some extensions of the relation : $\frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi}$ |



M. R. and F. A. Ceccopieri, arXiv: 1812.04286. JHEP accepted.

Relevant for processes involving heavy particles is the splitting term (1v2 mechanism) i.e.:

$$* D_{j_1, j_2}(x_1, x_2) = \int d^2 b_{\perp} \tilde{F}_{j_1, j_2}(x_{1,2}, b_{\perp})$$




In pQCD evolution:  $\frac{dD_{j_1 j_2}(x_1, x_2; t)}{dt} = \left\{ \begin{array}{l} \text{Homogeneous term (double DGLAP)} \\ + \\ \sum_{j'} F_{j'}(x_1 + x_2; t) \underbrace{\frac{1}{x_1 + x_2} P_{j' \rightarrow j_1 j_2}}_{\text{SPLITTING TERM}} \left( \frac{x_1}{x_1 + x_2} \right) \end{array} \right.$

Gaunt J.R. and Stirling W. J., JHEP 03 (2010)


1v2

2v2

**SPLITTING TERM**



+



$$\frac{\sigma_{eff}}{3\pi} \left( 1 + \frac{3}{2} r_v \right) \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi} \left( 1 + 2 r_v \right)$$

$$r_v \sim \frac{F_{j_1 j_2}^{splitting}(x_1, x_2, k_{\perp} = 0; t)}{F_{j_1 j_2}(x_1, x_2, k_{\perp} = 0; t)}$$

Due to the difficulty in the estimate of the 2 contributions:


with:  
 $0 \leq r_v \leq 1$

Absolute minimum  
 $r_v = 0$

$$\frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq \frac{3 \sigma_{eff}}{\pi}$$

Absolute maximum  
 $r_v = 1$


# Some extensions of the relation : $\frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi}$ |



M. R. and F. A. Ceccopieri, arXiv: 1812.04286. JHEP accepted.

Relevant for processes involving heavy particles is the splitting term (1v2 mechanism) i.e.:

$$* D_{j_1, j_2}(x_1, x_2) = \int d^2 b_{\perp} \tilde{F}_{j_1, j_2}(x_{1,2}, b_{\perp})$$




In pQCD evolution:  $\frac{dD_{j_1 j_2}(x_1, x_2; t)}{dt} = \left\{ \begin{array}{l} \text{Homogeneous term (double DGLAP)} \\ + \\ \sum_{j'} F_{j'}(x_1 + x_2; t) \underbrace{\frac{1}{x_1 + x_2} P_{j' \rightarrow j_1 j_2}}_{\text{SPLITTING TERM}} \left( \frac{x_1}{x_1 + x_2} \right) \end{array} \right.$

Gaunt J.R. and Stirling W. J., JHEP 03 (2010)


1v2

2v2

**SPLITTING TERM**



+



$$\frac{\sigma_{eff}}{3\pi} \left( 1 + \frac{3}{2} r_v \right) \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi} \left( 1 + 2 r_v \right)$$

$$r_v \sim \frac{F_{j_1 j_2}^{splitting}(x_1, x_2, k_{\perp} = 0; t)}{F_{j_1 j_2}(x_1, x_2, k_{\perp} = 0; t)}$$

Due to the difficulty in the estimate of the 2 contributions:

with:  
 $0 \leq r_v \leq 1$

Absolute minimum  
 $r_v = 0$


$$\frac{\sigma_{eff}}{3\pi}$$

$$\leq \langle b^2 \rangle \leq$$

$$\frac{3 \sigma_{eff}}{\pi}$$

Absolute maximum  
 $r_v = 1$


# Some extensions of the relation : $\frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq \frac{\sigma_{eff}}{\pi}$ |



M. R. and F. A. Ceccopieri, arXiv: 1812.04286. JHEP accepted.

Relevant for processes involving heavy particles is the splitting term (1v2 mechanism) i.e.:

$$* D_{j_1, j_2}(x_1, x_2) = \int d^2 b_{\perp} \tilde{F}_{j_1, j_2}(x_{1,2}, b_{\perp})$$



In pQCD evolution:

$$\frac{dD_{j_1 j_2}(x_1, x_2; t)}{dt} =$$

Gaunt J.R. and Stirling V

Homogeneous term (double DGLAP)

$$+ \frac{1}{1+x_2} P_{j' \rightarrow j_1 j_2} \left( \frac{x_1}{x_1+x_2} \right)$$

IG TERM



$$r_v \sim \frac{F_{j_1 j_2}^{splitting}(x_1, x_2, k_{\perp} = 0; t)}{F_{j_1 j_2}(x_1, x_2, k_{\perp} = 0; t)}$$

with:

$$0 \leq r_v \leq 1$$

1v2

2v2




1) Minimum as function of  $r_v$   
 $m(r_v)$

2) Maximum as function of  $r_v$   
 $M(r_v)$

$$\text{Absolute minimum } r_v = 0 \quad \frac{\sigma_{eff}}{3\pi} \leq \langle b^2 \rangle \leq \frac{3 \sigma_{eff}}{\pi} \quad \text{Absolute maximum } r_v = 1$$

# What next: pion double PDF



M. R., S. Scopetta, M. Traini and V.Vento, EPJC 78, no. 9,782 (2018)

The dPDF expression, at the hadronic scale, evaluated in the intrinsic frame, in term of meson wave function:

$$f_2(x, k_\perp) = \frac{1}{2} \sum_{h, h'} \int \frac{d^2 k_{1\perp}}{2(2\pi)^3} \psi_{h, h'}(x, \vec{k}_{1\perp}) \underbrace{\psi_{h, h'}^*(x, \vec{k}_{1\perp} + \vec{k}_\perp)}_{\text{Meson wave function}}$$


Parton helicities

Intrinsic parton momentum

Meson wave function

\*See talk by W. Broniowski about the integrated dPDF in the NJL model

# What next: pion double PDF





M. R., S. Scopetta, M. Traini and V.Vento, EPJC 78, no. 9,782 (2018)

The dPDF expression, at the hadronic scale, evaluated in the intrinsic frame, in term of meson wave function:


$$f_2(x, k_\perp) = \frac{1}{2} \sum_{h,h'} \int \frac{d^2 k_{1\perp}}{2(2\pi)^3} \psi_{h,h'}(x, \vec{k}_{1\perp}) \psi_{h,h'}^*(x, \vec{k}_{1\perp} + \vec{k}_\perp)$$

Parton helicities
Intrinsic parton momentum
Meson wave function




1) Also for pion, model calculations indicate that factorization on  $x - k_\perp$  does not work!

2) Also for pion, model calculations indicate that dPDF can not be described in terms of GPDs (Dotted line=dPDF approximated).



# What next: pion double PDF




M. R., S. Scopetta, M. Traini and V.Vento, EPJC 78, no. 9,782 (2018)


The dPDF expression, at the hadronic scale, evaluated in the intrinsic frame, in term of meson wave function:

$$f_2(x, k_\perp) = \frac{1}{2} \sum_{h,h'} \int \frac{d^2 k_{1\perp}}{2(2\pi)^3} \psi_{h,h'}(x, \vec{k}_{1\perp}) \psi_{h,h'}^*(x, \vec{k}_{1\perp} + \vec{k}_\perp)$$

Parton helicities
Intrinsic parton momentum
Meson wave function




$$M_{ud}^{22}(y, Q^2) = \int_0^1 dx_1 \int_0^{1-x_1} dx_2 x_1 x_2 \bar{F}_{ud}(x_1, x_2, y, Q^2)$$



The latter is a quantity close to those evaluated in “new” lattice studies of DPS. Future comparison are in principle possible to obtain new information on dPDF from lattice QCD.



# What next: pion double PDF



M. R., S. Scopetta, M. Traini and V.Vento, EPJC 78, no. 9,782 (2018)

The dPDF expression, at the hadronic scale, evaluated in the intrinsic frame, in term of meson wave function:


$$f_2(x, k_\perp) = \frac{1}{2} \sum_{h, h'} \int \frac{d^2 k_{1\perp}}{2(2\pi)^3} \psi_{h, h'}(x, \vec{k}_{1\perp}) \underbrace{\psi_{h, h'}^*(x, \vec{k}_{1\perp} + \vec{k}_\perp)}_{\text{Meson wave function}}$$

Parton helicities  $\leftarrow$  Intrinsic parton momentum  $\leftarrow$  Meson wave function


We also computed the  $\pi\pi$  mean  $\sigma_{eff}$ :

$$\bar{\sigma}_{eff} = 41 \text{ mb}$$

This result has been used in experimental analysis for DPS at COMPASS: **arXiv: 1909.06195**






# Conclusions





## A CQM calculation of the dPDFs with a Poincare' invariant approach

- ✓ longitudinal and transverse correlations are found;
- ✓ deep study on relativistic effects: **transverse and longitudinal model independent correlations have been found**;
- ✓ pQCD evolution of dPDFs, including non perturbative degrees of freedom into the scheme: **correlations are present at high energy scales and in the low  $x$  region**;
- ✓ calculation of the effective X-section within different models in the valence region:
- ✓ **x-dependent quantity obtained!**
- ✓ Calculation of mean partonic distance from present experimental analyses
- ✓ calculation of pion dPDF





## Study of DPS in same sign WW production at the LHC

- ✓ Calculations of the DPS cross section of same sign WW production
- ✓ **dynamical correlations are found to be measurable in the next run at the LHC**



A proton imaging (complementary to that investigated by means of electromagnetic probes) can/will be obtained in the next LHC runs!


# Further Information on:





Thanks

# Some extensions of the relation : $\frac{\sigma_{\text{eff}}}{3\pi} \leq \langle b^2 \rangle \leq \frac{\sigma_{\text{eff}}}{\pi} \quad ||$



M. R. and F. A. Ceccopieri, arXiv: 1812.04286. JHEP accepted.


IF WE DO NOT CONSIDER ANY FACTORIZATION ANSATZ IN DOUBLE PDFs:

$$\frac{\sigma_{\text{eff}}(x_1, x_2)}{3\pi} \left[ r^{2\nu_2}(x_1, x_2)^2 + \frac{3}{2} r^{2\nu_1}(x_1, x_2)^2 r_\nu \right] \leq \langle b^2 \rangle_{x_1, x_2} \leq \frac{\sigma_{\text{eff}}(x_1, x_2)}{\pi} \left[ r^{2\nu_2}(x_1, x_2)^2 + 2r^{2\nu_1}(x_1, x_2)^2 r_\nu \right]$$

$$r^{2\nu_2}(x_1, x_2) = \frac{F(x_1, x_2, k_\perp = 0; t)}{F(x_1; t)F(x_2; t)}$$

$$r^{2\nu_1}(x_1, x_2) = \frac{F^{\text{splitting}}(x_1, x_2, k_\perp = 0; t)}{F(x_1; t)F(x_2; t)}$$


# The Effective X-section calculation




M. R., S. Scopetta, M. Traini and V.Vento, PLB 752, 40 (2016)

Our predictions of  $\sigma_{eff}$ , **without any approximation**, in the valence region at different energy scales:


$$\sigma_{eff}(x_1, x_2, \mu_0^2) \xrightarrow[\text{pQCD evolution of dPDFs}]{\text{pQCD evolution of PDFs}} \sigma_{eff}(x_1, x_2, Q^2)$$



**Gluons ⊗ Gluons**



**Valence ⊗ Valence quarks**



**Valence quark ⊗ Sea quark**  
Partons involved in, e.g., same sign WW production.

The old data lie in the obtained range of  $\sigma_{eff}$


$$\bar{\sigma}_{eff} \sim 21 \text{ mb}$$

Similar results obtained with dPDFs calculated within AdS/QCD soft-wall model  
M. Traini, M. R., S. Scopetta and V.Vento, PLB 768, 270 (2017)

- $x_i$  dependence of  $\sigma_{eff}$  may be model independent feature
- Absolute value of  $\sigma_{eff}$  is a model dependent result




# The Effective X-section calculation



M. R., S. Scopetta, M. Traini and V.Vento, PLB 752, 40 (2016)

Our predictions of  $\sigma_{eff}$ , without any approximation, in the valence region at different energy scales:

$$\sigma_{eff}(x_1, x_2, \mu_0^2) \xrightarrow[\text{pQCD evolution of dPDFs}]{\text{pQCD evolution of PDFs}} \sigma_{eff}(x_1, x_2, Q^2 = 250 \text{ GeV}^2)$$



**ACCESS THE DOUBLE PARTON CORRELATIONS**

**ACCESS THE PARTONIC STRUCTURE OF THE PROTON**


The old data lie in the obtained range of  $\sigma_{eff}$

Valence quark  $\otimes$  Sea quark  
Partons involved in, e.g., same sign WW production.

Similar results obtained with dPDFs calculated within AdS/QCD soft-wall model  
M. Traini, M. R., S. Scopetta and V.Vento, PLB 768, 270 (2017)

- $x_i$  dependence of  $\sigma_{eff}$  may be model independent feature
- Absolute value of  $\sigma_{eff}$  is a model dependent result

# The Effective X-section calculation



M. R., S. Scopetta, M. Traini and V.Vento, PLB 752, 40 (2016)

$$\sigma_{eff} = \frac{m}{2} \frac{\sigma_A^{pp'} \sigma_B^{pp'}}{\sigma_{double}^{pp}}$$

This quantity can be written in terms of PDFs and dPDFs ( $_2$ GPDs)

Here the scale is omitted

$$\sigma_{eff}(x_1, x'_1, x_2, x'_2) = \frac{\sum_{i,k,j,l} \mathbf{F}_i(x_1) \mathbf{F}_k(x'_1) \mathbf{F}_j(x_2) \mathbf{F}_l(x'_2) C_{ik} C_{jl}}{\sum_{i,j,k,l} C_{ik} C_{jl} \int \mathbf{F}_{ij}(x_1, x_2; \mathbf{k}_\perp) \mathbf{F}_{kl}(x'_1, x'_2; -\mathbf{k}_\perp) \frac{d\mathbf{k}_\perp}{(2\pi)^2}}$$

Colour coefficient

Non trivial x-dependence

If factorization between dPDF and PDFs held:

$$F_{ab}(x_1, x_2, \vec{k}_\perp) = F_a(x_1) F_b(x_2) \tilde{T}(\vec{k}_\perp)$$

“EFFECTIVE FORM FACTOR”

Conjugated variable to  $\vec{k}_\perp$

$$\sigma_{eff}(x_1, x'_1, x_2, x'_2) \rightarrow \sigma_{eff} = \left[ \int \frac{d\vec{k}_\perp}{(2\pi)^2} \tilde{T}(\vec{k}_\perp) T(-\vec{k}_\perp) \right]^{-1} = \left[ \int d\vec{b}_\perp \Gamma(\vec{b}_\perp)^2 \right]^{-1}$$

Constant value w.r.t.  $x_i$



**NO CORRELATIONS!**