

# Double parton distributions of the pion in the NJL model

Wojciech Broniowski<sup>1,2</sup> and Enrique Ruiz Arriola<sup>3</sup>

<sup>1</sup>Institute of Nuclear Physics PAN, Cracow

<sup>2</sup>Jan Kochanowski U., Kielce

<sup>3</sup>U. of Granada

Light Cone 2019



École Polytechnique, Palaiseau, 16-20 September 2019

# Motivation for multi-parton distributions

- Old story ([Fermilab](#)), renewed interest (e.g., ATLAS measurement for  $pp \rightarrow W+2\text{ jets}$  2013) [Kuti, Weiskopf 1971, Konishi, Ukwa, Veneziano 1979, Gaunt, Stirling 2010, Diehl, Ostermeier, Schäfer 2012, ..., reviews: Bartalani et al. 2011, Snigirev 2011, Rinaldi, Ceccopieri 2018] [\(see Matteo Rinaldi's talk this afternoon\)](#)
- Model exploration [MIT bag: Chang, Manohar, Waalewijn 2013], valon [WB+ERA 2013], constituent quarks: Rinaldi, Scopetta, Vento 2013, Rinaldi, Scopetta, Traini, Vento 2018]
- Questions: factorization, interference of single- and double-parton scattering [Manohar, Waalewijn 2012, Gaunt 2013], longitudinal-transverse decoupling, positivity bounds [Diehl, Casemets 2013], transverse momentum dependence [Casemets, Gaunt 2019]
- [Gaunt-Stirling sum rules](#) [Gaunt, Stirling 2010, WB+ERA 2013, Diehl, Plöß, Schäfer 2019]

# Double parton scattering

[example from Łuszczak, Maciuła, Szczurek 2011]



DPS can be comparable to SPS at the LHC  
Assumption:  $D_{gg}(x_1, x_2, \mathbf{b}) = g(x_1)g(x_2)F(\mathbf{b})$   
– no correlations, transverse-longitudinal factorization

# Definition

Intuitive probabilistic definition:

Multi-parton distribution = probability distribution that struck partons have LC momentum fractions  $x_i$

Field-theoretic definition of (spin-averaged) sPDF and dPDF [Diehl, Ostermeier, Schaeffer 2012] of a hadron with momentum  $p$ :

$$D_j(x) = \int \frac{dz^-}{2\pi} e^{ixz^- p^+} \langle p | \mathcal{O}_j(0, z) | p \rangle \Big|_{z^+=0, z=0}$$

$$\begin{aligned} F_{j_1 j_2}(x_1, x_2, \mathbf{y}) &= 2p^+ \int dy^- \frac{dz_1^-}{2\pi} \frac{dz_2^-}{2\pi} e^{i(x_1 z_1^- + x_2 z_2^-) p^+} \\ &\quad \times \langle p | \mathcal{O}_{j_1}(y, z_1) \mathcal{O}_{j_2}(0, z_2) | p \rangle \Big|_{z_1^+ = z_2^+ = y^+ = 0, z_1 = z_2 = 0} \end{aligned}$$

$$\mathcal{O}_q(y, z) = \frac{1}{2} \bar{q}(y - \frac{z}{2}) \gamma^+ q(y + \frac{z}{2}), \dots$$


$$v^\pm = (v^0 \pm v^3)/\sqrt{2}$$

$\mathbf{y}$  plays the role of the transverse distance between the two quarks

# dPDF in momentum space

Fourier transform in  $y$

$$F_{j_1 j_2}(x_1, x_2, \mathbf{y}) \rightarrow \tilde{F}_{j_1 j_2}(x_1, x_2, \mathbf{q})$$



Special case of  $\mathbf{q} = \mathbf{0}$ :

$$D_{j_1 j_2}(x_1, x_2) = \tilde{F}_{j_1 j_2}(x_1, x_2, \mathbf{q} = \mathbf{0})$$

# Gaunt-Stirling sum rules

[Gaunt, Stirling, JHEP (2010) 1003:005, Gaunt, PhD Thesis]

Fock-space decomposition on LC + conservation laws →

$$|P\rangle = \sum_N \int d[x, \mathbf{k}]_N \Phi(\{x_i, \mathbf{k}_i\}) |\{x_i, \mathbf{k}_i\}\rangle_N$$

$$d[x, \mathbf{k}]_N = \prod_{i=1}^N \left[ \frac{dx_i d^2 k_i}{\sqrt{2(2\pi)^3 x_i}} \right] \delta \left( 1 - \sum_{i=1}^N x_i \right) \delta^{(2)} \left( 1 - \sum_{i=1}^N \mathbf{k}_i \right)$$

# Gaunt-Stirling sum rules

[Gaunt, Stirling, JHEP (2010) 1003:005, Gaunt, PhD Thesis]

Fock-space decomposition on LC + conservation laws →

$$\sum_i \int_0^{1-x_2} dx_1 x_1 D_{ij}(x_1, x_2) = (1 - x_2) D_j(x_2) \quad (\text{momentum})$$

$$\int_0^{1-x_2} dx_1 D_{i_{\text{val}} j}(x_1, x_2) = (N_{i_{\text{val}}} - \delta_{ij} + \delta_{\bar{i}\bar{j}}) D_j(x_2) \quad (\text{quark number})$$

$$(A_{i_{\text{val}}} \equiv A_i - A_{\bar{i}})$$

$$N_{i_{\text{val}}} = \int_0^1 dx D_{i_{\text{val}}}(x)$$

- Preserved by DGLAP evolution
- Non-trivial to satisfy with the (guessed) function
- Checked in light-front perturbation theory and in lowest-order covariant calculations in [Diehl, Plößl, Schäfer 2019]

Important and fundamental constraints!

## Simple example (valon model)

$|\Lambda\rangle = |uds\rangle$  (to avoid the complications of indistinguishable partons)

$$D_{uds}(x_1, x_2, x_3) = f(x_1, x_2, x_3)\delta(1 - x_1 - x_2 - x_3)$$

$$D_{ud}(x_1, x_2) = \int dx_3 D_{uds}(x_1, x_2, x_3) = f(x_1, x_2, 1 - x_1 - x_2)$$

$$D_{us}(x_1, x_3) = \dots$$

$$D_u(x_1) = \int_0^{1-x_1} dx_2 D_{ud}(x_1, x_2) = \int_0^{1-x_1} dx_3 D_{us}(x_1, x_3)$$

$$\int_0^{1-x_1} dx_2 \textcolor{blue}{x}_2 D_{ud}(x_1, x_2) + \int_0^{1-x_1} dx_3 \textcolor{blue}{x}_3 D_{us}(x_1, x_3)$$

$$= \int dx_2 dx_3 (\textcolor{blue}{x}_2 + \textcolor{blue}{x}_3) D_{uds}(x_1, x_2, x_3) = \int dx_2 dx_3 (1 - \textcolor{blue}{x}_1) D_{uds}(x_1, x_2, x_3)$$
$$= (1 - \textcolor{blue}{x}_1) D_u(x_1)$$

# Attempts of bottom-up construction

- Gaunt, Stirling (2011)

$$D_{ij}(x_1, x_2) = D_i(x_1) D_j(x_2) \frac{(1 - x_1 - x_2)^2}{(1 - x_1)^{2+n_1} (1 - x_2)^{2+n_2}}$$

(do not satisfy the GS sum rules)

- Lewandowska, Golec-Biernat 2014

$$D_{ij}(x_1, x_2) = \frac{1}{1 - x_2} D_i\left(\frac{x_1}{1 - x_2}\right) D_j(x_2)$$

...

(no parton exchange symmetry, negative  $D_{qq}$  at large  $x$ )

- Can never be unique: marginal projections do not determine the two-particle distribution


Problems!

# Message

- Construct the multiparticle distribution (model, data?) and go down with marginal projections

[cf. a similar in spirit “top-down” study by M. Rinaldi et al. 2018 with the Brodsky - de Teramond AdS/CFT soft wall pion wave function]

# Chiral quark models




- $\chi$ SB breaking  $\rightarrow$  massive quarks
- Point-like interactions
- Soft matrix elements with pions (and photons,  $W, Z$ )
- Large- $N_c$   $\rightarrow$  one-quark loop
- Regularization

pion – Goldstone boson of  $\chi$ SB, fully relativistic  $q\bar{q}$  bound state of the Bethe-Salpeter equation

Quantities evaluated at the quark model scale  
(where **constituent quarks are the only degrees of freedom**)

# Chiral quark models



- $\chi$ SB breaking  $\rightarrow$  massive quarks
- Point-like interactions
- Soft matrix elements with pions (and photons,  $W, Z$ )
- Large- $N_c$   $\rightarrow$  one-quark loop
- Regularization


pion – Goldstone boson of  $\chi$ SB, fully relativistic  $q\bar{q}$  bound state of the Bethe-Salpeter equation

Need for evolution

Gluon dressing, renorm-group improved

# sPDF in NJL

[Davidson, Arriola, 1995]



$$q_{\text{val}}(x; Q_0) = 1 \times \theta[x(1-x)]$$

(proper treatment of symmetries with regularization)

Quarks are the only degrees of freedom, hence saturate the sPDF sum rules:


$$\int_0^1 dx q_{\text{val}}(x; Q_0) = 1 \text{ (valence)}, \quad 2 \int_0^1 dx x q_{\text{val}}(x; Q_0) = 1 \text{ (momentum)}$$

# Scale and evolution

QM provide non-perturbative result at a low scale  $Q_0$

$$F(x, Q_0)|_{\text{model}} = F(x, Q_0)|_{\text{QCD}}, \quad Q_0 - \text{the matching scale}$$

Quarks carry 100% of momentum at  $Q_0$ , adjusted such that when evolved to  $Q = 2 \text{ GeV}$ , they carry the experimental value of 47% (radiative generation of gluons and sea quarks)




LO DGLAP evolution

$$Q_0 = 313^{+20}_{-10} \text{ MeV}$$

NLO close to LO

$$\sim (1-x)^{p+\frac{4C_F}{\beta_0} \log \frac{\alpha(Q_0)}{\alpha(Q)}}$$

# Pion valence quark PDF, NJL vs E615




points: Fermilab E615  
Drell-Yan,  $\pi^\pm W \rightarrow \mu^+ \mu^- X$

band: QM + LO DGLAP  
from  $Q_0 = 313^{+20}_{-10}$  MeV to  
 $Q = 4$  GeV

Many predictions for related quantities: DA, GPD, TDA, TMD,  
quasi/pseudo DA/PDF...

# dPDF of the pion in NJL model



$$D_{u\bar{d}}(x_1, x_2) = 1 \times \delta(1 - x_1 - x_2) \theta(x_1) \theta(x_2)$$

- Special case of the valon model
- GS sum rules satisfied
- ... at the quark-model scale → need for evolution

# dDGLAP evolution in the Mellin space

[Kirschner 1979, Shelest, Snigirev, Zinovev 1982]: method of solving dDGLAP based on the Mellin moments, similarly to sPDF

$$M_j^n = \int_0^1 dx x^n D_j(x), \quad M_{j_1 j_2}^{n_1 n_2} = \int_0^1 dx_1 \int_0^1 dx_2 \theta(1-x_1-x_2) x_1^{n_1} x_2^{n_2} D_{j_1 j_2}(x_1, x_2)$$

$$\frac{d}{dt} M_{j_1 j_2}^{n_1 n_2} = \sum_i P_{i \rightarrow j_1}^{n_1} M_{ij_2}^{n_1 n_2} + \sum_i P_{i \rightarrow j_2}^{n_2} M_{j_1 i}^{n_1 n_2} + \sum_i \left( P_{i \rightarrow j_1 j_2}^{n_1 n_2} + \tilde{P}_{i \rightarrow j_1 j_2}^{n_1 n_2} \right) M_i^{n_1+n_2}$$

$t = \frac{1}{2\pi\beta} \log [1 + \alpha_s(\mu)\beta \log(\Lambda_{\text{QCD}}/\mu)]$  (**single scale** for simplicity),  $\beta = \frac{11N_c - 2N_f}{12\pi}$   
(inhomogeneous term from coupling to sPDF's)

For valence-valence distributions there are no partons  $i$  decaying into a pair of valence quarks ( $P_{i \rightarrow j_1 j_2} = 0$ )  $\rightarrow$  **inhomogeneous term vanishes**

$$\text{dPDF : } \frac{d}{dt} M_{j_1 j_2}^{n_1 n_2}(t) = (P_{j_1 \rightarrow j_1}^{n_1} + P_{j_2 \rightarrow j_2}^{n_2}) M_{j_1, j_2}^{n_1 n_2}(t)$$

$$\text{sPDF : } \frac{d}{dt} M_j^n(t) = P_{j \rightarrow j}^n M_j^n(t)$$

# Solution

[10 lines in Mathematica (!)]

$$M_j^n(t) = e^{P_{j \rightarrow j}^n(t-t_0)} M_j^n(t_0)$$

$$M_{j_1 j_2}^{n_1 n_2}(t) = e^{[P_{j_1 \rightarrow j_1}^{n_1} + P_{j_2 \rightarrow j_2}^{n_2}](t-t_0)} M_{j_1 j_2}^{n_1 n_2}(t_0)$$

inverse Mellin transform:


$$D_j(x; t) = \int_C \frac{dn}{2\pi i} x^{-n-1} M_j^n(t)$$

$$D_{j_1 j_2}(x_1, x_2; t) = \int_C \frac{dn_1}{2\pi i} x_1^{-n_1-1} \int_{C'} \frac{dn_2}{2\pi i} x_2^{-n_2-1} M_{j_1, j_2}^{n_1, n_2}(t)$$


$n$  and  $n'$  are complex variables and the contours  $C$  and  $C'$  lie right to singularities of  $M$

- correlations  $\rightarrow M_{j_1 j_2}^{n_1 n_2}(t) \neq M_{j_1}^{n_1}(t) M_{j_2}^{n_2}(t)$  – no separability
- valence-valence:  $M_{j_1 j_2}^{n_1 n_2}(t)/[M_{j_1}^{n_1}(t) M_{j_2}^{n_2}(t)]$  independent of  $t$





$$x_1 x_2 D_{u\bar{d}}^{\pi^+}(x_1, x_2)$$



# $D_{ud}^{\pi^+}(x_1, x_2) - \log \text{ scale}$




# Correlation



# Gluon correlation





[Golec-Biernat, Lewandowska, Serino, Snyder, Staśto 2015]



# Gluon correlation

[Golec-Biernat, Lewandowska, Serino, Snyder, Staśto 2015]

$x_2=0.5$




# Valence moments in NJL

$$\frac{\langle x_1^n x_2^m \rangle}{\langle x_1^n \rangle \langle x_2^m \rangle} = \frac{(1+n)!(1+m)!}{(1+n+m)!} \quad (\text{NJL, any scale})$$

(independent of the evolution scale)

1	1	1	1	1	1
1	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{2}{5}$	$\frac{1}{3}$	$\frac{2}{7}$
1	$\frac{1}{2}$	$\frac{1}{10}$	$\frac{1}{5}$	$\frac{1}{7}$	$\frac{3}{28}$
1	$\frac{5}{1}$	$\frac{1}{5}$	$\frac{35}{1}$	$\frac{14}{5}$	$\frac{1}{21}$
1	$\frac{1}{3}$	$\frac{1}{7}$	$\frac{14}{1}$	$\frac{1}{126}$	$\frac{1}{42}$
1	$\frac{2}{7}$	$\frac{1}{28}$	$\frac{1}{21}$	$\frac{1}{42}$	$\frac{1}{77}$



Double moments reduced compared to product of single moments  
[lattice results coming shortly, Zimmermann et al.]

# Summary

- Top-down strategy of constructing multi-parton distributions → formal features guaranteed, in particular GS sum rules
- Phenomenological sPDF's as constraints
- NJL → valon initial condition,  $\text{const} \times \delta(1 - x_1 - x_2)$ , dDGLAP
- Correlations decrease with increasing evolution scale and are probably not very important ( $\pm 25\%$ ) in the range probed by experiments, justifying the product ansatz in that limit
- Moments measure the  $x_1 - x_2$  factorization breaking; can be verified in forthcoming lattice calculations