



Contribution ID: 1173

Type: Parallel Session Talk

Measurements of $|V_{us}|$ and Second Class Currents and Searches for Violation of Lepton Universality and CPT in Tau Decays at BABAR

Friday 23 July 2010 10:15 (13 minutes)

We report on a variety of results involving decays of the tau lepton using the very large sample of tau+tau- pairs produced in e+e- annihilation data collected with the BaBar detector at the PEP-II asymmetric-energy B Factory near a center-of-mass energy of 10.58 GeV. From measurements of the ratios of branching fractions: $B(\tau \rightarrow \mu \nu \text{ nubar}) / B(\tau \rightarrow e \nu \text{ nubar})$, $B(\tau \rightarrow \pi \nu) / B(\tau \rightarrow e \nu \text{ nubar})$, and $B(\tau \rightarrow K \nu) / B(\tau \rightarrow e \nu \text{ nubar})$ we test with high precision the Standard Model assumption of mu-e and tau-mu charged current lepton universality and provide a determination of the Cabibbo-Kobayashi-Maskawa matrix element $|V_{us}|$. Furthermore, we report on preliminary measurements of $\tau^- \rightarrow K^- n \pi^0 \nu_{\tau}$ with $n = 0,1,2,3$ and $\tau^- \rightarrow \pi^- n \pi^0 \nu_{\tau}$ with $n = 3,4$ as well as on the measurements of the branching fractions and hadronic mass distributions of $\tau^- \rightarrow K_S^0 \pi^- \nu_{\tau}$, $\tau^- \rightarrow K_S^0 \pi^- \pi^0 \nu_{\tau}$, $\tau^- \rightarrow K_S^0 \pi^- K^0_L \nu_{\tau}$. Data from the inclusive strange tau decay results are used in a different determination of $|V_{us}|$. We also report on our search for second class currents in $\tau^- \rightarrow \pi^- \eta \nu_{\tau}$, where the eta decays into $\pi^+ \pi^- \pi^0$ and our measurement of the tau mass. We obtain a test of CPT by measuring the difference between the masses of the tau+ and tau-.

Author: BABAR, Collaboration (SLAC)**Presenter:** LUSIANI, Alberto (Dipartimento di Fisica)**Session Classification:** 06 - CP violation, CKM and Rare Decays**Track Classification:** 06 - CP violation, CKM and Rare Decays