



Contribution ID: 1096

Type: Parallel Session Talk

Hadronic $b \rightarrow c$ decays at Belle

Friday, 23 July 2010 16:30 (13 minutes)

We present a measurement of the unitarity triangle angle ϕ_3 using Dalitz plot analysis of three-body neutral D decays from the $B^+ \rightarrow D^0 K^+$ process. The results are based on a large sample of B anti- B pairs recorded at the Upsilon(4S) resonance with the Belle detector at the KEKB e^+e^- collider. The decay $B \rightarrow D^0 K^+$ ($D = D^0$ or anti- D^0) includes the $b \rightarrow u$ transition and plays a crucial role in the measurement of the CP-violating angle ϕ_3 . We present the result of a study of the decay $B \rightarrow D^0 K^+$ where the D meson is reconstructed from $K^+ \pi^-$. We also report improved measurements of the branching fractions for the decays $B^0 \rightarrow D_s^+ \pi^-$ and $anti-B^0 \rightarrow D_s^+ K^-$. Based on these results, we determine the ratio between the amplitudes of the doubly Cabibbo suppressed decay $B^0 \rightarrow D^+ \pi^-$ and the Cabibbo favored decay $B^0 \rightarrow D^0 \pi^+$. We studied the three-body baryonic B^+ decays, $B^+ \rightarrow p \text{ anti-Lambda } D^0$. The branching fractions as well as the differential branching fractions as a function of the mass of the p anti-Lambda system are presented. These results are compared with theoretical predictions based on the generalized factorization approach. We present a study of the exclusive decays $B^0 \rightarrow D_s^- K^0_S \pi^+$ and $B^- \rightarrow D_s^+ K^- K^-$. We use the $D_s^- \rightarrow \phi \pi^-$, $anti-K^0_S \rightarrow (892)0 K^-$ and $K^0_S K^-$ decay modes for D_s reconstruction.

Author: BELLE, Collaboration (KEK)

Presenter: Mr JOSHI, Nikhil Jayant (Tata Institute of Fundamental Research, Mumbai)

Session Classification: 06 - CP violation, CKM and Rare Decays

Track Classification: 06 - CP violation, CKM and Rare Decays