



Contribution ID: 911

Type: Poster

## Search for novel origins of cosmic-ray antiprotons and antimatter with BESS-Polar flight over Antarctica.

The primary aims of the BESS-Polar program are precise measurements of the low-energy antiproton spectrum and search for cosmologically significant antimatter, which would provide new clues to understand the early Universe. The second flight (BESS-Polar II) over Antarctica was successfully carried out in December 2007 - January 2008. We performed 24.5 days scientific observation just at the solar minimum. The payload worked well during the flight and 4.7 billion cosmic-ray events were collected, which corresponds to 10-20 times statistics of the BESS data taken in the previous solar minimum period (1995 and 1997). Based on the BESS-Polar II data, we will present recent preliminary results of cosmic-ray antiproton measurements and sensitive search for antimatter.

**Primary author:** Dr YOSHIMURA, Koji (High Energy Accelerator Research Organization (KEK))

**Co-authors:** Mr ITAZAKI, A. (Kobe University); Prof. YAMAMOTO, Akira (High Energy Accelerator Research Organization (KEK)); Dr MOISEEV, Alex (National Aeronautics and Space Administration, Goddard Space Flight Center (NASA/GSFC)); Dr HORIKOSHI, Atsushi (High Energy Accelerator Research Organization (KEK)); Prof. SEO, Eun-Suk (University of Maryland); Dr FUKE, Hideyuki (Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency (ISAS/JAXA)); Dr MITCHELL, John (National Aeronautics and Space Administration, Goddard Space Flight Center (NASA/GSFC)); Prof. ORMES, Jonathan (University of Denver); Prof. NISHIMURA, Jun (The University of Tokyo); Mr SUZUKI, Junichi (High Energy Accelerator Research Organization (KEK)); Mr MATSUMOTO, K. (High Energy Accelerator Research Organization (KEK)); TAKEUCHI, K. (High Energy Accelerator Research Organization (KEK)); Mr SAKAI, Kenichi (The University of Tokyo); Mr TANAKA, Kenichi (High Energy Accelerator Research Organization (KEK)); Dr KIM, Ki-Chun (University of Maryland); Dr ABE, Ko (Kobe University); Dr SASAKI, Makoto (National Aeronautics and Space Administration, Goddard Space Flight Center (NASA/GSFC)); Dr HASEGAWA, Masaya (High Energy Accelerator Research Organization (KEK)); Prof. NOZAKI, Mitsuaki (High Energy Accelerator Research Organization (KEK)); Dr LEE, MooHyun (University of Maryland); THAKUR, Neeharika (University of Denver); Dr ORITO, Reiko (High Energy Accelerator Research Organization (KEK)); Dr STREITMATTER, Robert (National Aeronautics and Space Administration, Goddard Space Flight Center (NASA/GSFC)); Ms SHINODA, Ryohko (High Energy Accelerator Research Organization (KEK)); Dr HAINO, Sadakazu (High Energy Accelerator Research Organization (KEK)); Dr MATSUDA, Shinya (High Energy Accelerator Research Organization (KEK)); Mr KUMAZAWA, T. (High Energy Accelerator Research Organization (KEK)); Prof. YAMAGAMI, Takamasa (Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency (ISAS/JAXA)); Prof. YOSHIDA, Tetsuya (Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency (ISAS/JAXA)); Dr HAMS, Thomas (National Aeronautics and Space Administration, Goddard Space Flight Center (NASA/GSFC)); Mr TAKASUGI, Y. (Kobe University); Dr MAKIDA, Yasuhiro (High Energy Accelerator Research Organization (KEK)); Dr SHIKAZE, Yoshiaki (Kobe University); Mr MATSUKAWA, Yosuke (Kobe University); Mr MYERS, Z. (National Aeronautics and Space Administration, Goddard Space Flight Center (NASA/GSFC))

**Presenter:** Dr YOSHIMURA, Koji (High Energy Accelerator Research Organization (KEK))

