



Contribution ID: 726

Type: Poster

Magical properties of 2540 km baseline Superbeam Experiment

The determination of the neutrino mixing matrix and mass-squared differences is one of the aims of neutrino physics today. This is a complicated affair, owing to the various parameter degeneracies. While the proposed 7500 km long magic baseline experiment simplifies the task considerably, the intense beam required for such an experiment seems futuristic by current standards. As an alternative, we highlight the magical properties of the 2540 km baseline. We propose a superbeam experiment at this much shorter baseline with a narrow band NuMI-like beam, and demonstrate the ability of this single setup to distinguish between the two mass hierarchies. This, we show, is possible with a moderate exposure and by running the experiment in the neutrino mode only.

Our results hold up to fairly small values of the mixing angle θ_{13} and irrespective of the value of the CP violating parameter. Unlike the magic baseline, it may also be possible to use this setup to measure CP violation in neutrino oscillation experiments.

Author: Prof. UMASANKAR, Sankagiri (I.I.T. Bombay, Mumbai, India)

Co-authors: Mr SINGH, Ravi Shanker (Department of Physics, Drown University); Mr RAUT, Sushant (I.I.T. Bombay, Mumbai, India)

Presenter: Prof. UMASANKAR, Sankagiri (I.I.T. Bombay, Mumbai, India)

Track Classification: 07 - Neutrinos