

Charm and Beauty Production from Secondary Vertexing at HERA

Paul Thompson (Birmingham)




for the H1 and ZEUS Collaborations




- Motivation and analysis method
- Heavy Flavour jets in photoproduction
- Heavy Flavour jets in Deep Inelastic Scattering
- Contribution of Heavy Flavours to proton structure

Heavy Flavour Analyses



- In total $\sim 500\text{pb}^{-1}$ of high energy data collected per experiment
- luminosity upgrade in 2001
- detectors adjusted
- ZEUS: silicon micro vertex detector




*Many heavy flavour final analyses on full HERA I+II data.
Working on publication of remaining preliminaries and
combination of results*

Production of Heavy Quarks


Contribution of quasi-real photons at low Q^2

Direct γ



Resolved γ

(flavour excitation)



$Q^2 < 1 \text{ GeV}^2$ Photoproduction, $Q^2 > 1 \text{ GeV}^2$ DIS

Predominantly via boson gluon fusion

Test of perturbative QCD:

multi-scale problem (M , Q , p_T)

Directly sensitive to gluon density in the proton (PDFs)

Heavy Quark Production

Number of theoretical approaches:

Massless (Zero Mass), massive (Fixed Flavour) and general mass (GM) flavour number schemes (combination of massless/massive should provide best theoretical model).

QCD Calculations:

Fixed order - massive FFNS NLO(α_s^2) (FMNR, HVQDIS)

GM-VFNS PDFs - used in latest PDF fits

MSTW08 to NLO (α_s^2) and NNLO (α_s^3)

CTEQ 6.6 to NLO (α_s)


Monte-Carlo: LO (α_s) + Parton shower:

Collinear factorisation, DGLAP (PYTHIA, RAPGAP)

Contribution to Cross Section (DIS)

HERA I+II result:


- fraction of total DIS cross section from **charm** and **beauty**
- large charm fraction (~30%). Has influence on PDFs!
- small beauty fraction ~% (lower at low Q^2)
- mass thresholds visible
- good description by NNLO QCD



Tagging Heavy Quarks

Heavy quarks rarely produced, use properties of beauty hadrons:

- lifetime and mass
 - reconstruction of a secondary vertex
 - decay length and mass of tracks from secondary vertex
 - impact parameter



Vertex method allows measurement of all tracks to low p_T – increase statistics and reduce extrapolations to full phase space. Can compare with other methods semi-leptonic (1163 Juengst), reconstruction of charmed meson decays (1160 Jung, 1162 Roloff)

H1 and ZEUS vertex measurements

H1

- Inclusive charm and beauty in DIS
[Eur.Phys.J. C65 \(2010\) 89](#)
[arXiv:0907.2643](#)
- Charm and beauty jets in DIS
[DESY 10-083](#)

ZEUS


- Beauty dijets in Photoproduction
[ZEUS-prel-09-005](#)
- Beauty jets in DIS and F_2^{bb}
[ZEUS-prel-10-004](#)

Methods to discriminate heavy flavours from light quarks and to disentangle c from b are very similar for H1 and ZEUS

Highlight the important features here...



Flavour Tagging - secondary vertex

- Use all tracks ("inclusive") with hits in silicon detectors $p_T > 0.3(0.5) \text{ GeV}$ H1(ZEUS)
- 2D(3D) hits H1(ZEUS). Calculate 2D secondary vertex decay length and decay length significance $S_L = L/\sigma(L)$
- Sign of vertex given w.r.t jet axis
- Use also signed impact parameter δ of individual tracks



$$\alpha < 90^\circ \rightarrow \delta = +|\delta|$$

$$\alpha > 90^\circ \rightarrow \delta = -|\delta|$$






Flavour Separation

Significance $S = \delta/\sigma(\delta)$

DESY 10-083

For >2 tracks use NN



S_1 highest $|S|$
 S_2 2nd highest $|S|$

resolution

lifetime

c/b separation


Charm and beauty asymm. due to lifetime

Light flavours mostly symmetric

Photoproduction background small

Neural Network
 inputs include
 S_1, S_2, S_3, S_L and
 number of
 silicon tracks

Fitting Flavour Fractions



ZEUS-prel-09-005

Lights suppressed



b dominates

Reduce contribution of lights by using “mirror image” i.e. subtract negative bins from positive.

ZEUS fit S_L in bins of M_{VTX} , H1 fit S_1 , S_2 and NN output

Perform l, c, b fits in bins of e.g. p_T^{jet} or x, Q^2 to extract F_2^{bb} 10

Photoproduction b Dijets (ZEUS)



$Q^2 < 1 \text{ GeV}^2, P_T^{\text{Jet}} > 7(6) \text{ GeV}, -1.6 < \eta^{\text{Jet}} < 1.3$



Beauty jet cross section vs $p_T^{\text{Jet}}, \eta^{\text{Jet}}$

ZEUS-prel-09-005

Well described by (massive) NLO QCD

Agreement found with measurements from muon tagging (864 Geiser)

Beauty Jets In DIS (H1)



$Q^2 > 6 \text{ GeV}^2, P_T^{\text{Jet}} > 6 \text{ GeV}, -1 < \eta^{\text{Jet}} < 1.5$



Beauty jet cross sections vs E_T^{jet} and η^{jet}

DESY 10-083



Well described by (massive) NLO QCD

Good description (as for H1 γp analysis hep-ex/0605016)

Beauty Jets In DIS (ZEUS)





ZEUS-prel-10-004



Beauty jet cross sections vs Q^2 and x . Agreement with NLO QCD, although QCD lower at low Q^2 and low x

Charm Jets In DIS (H1)



$Q^2 > 6 \text{ GeV}^2, P_T^{\text{Jet}} > 6 \text{ GeV}, -1 < \eta^{\text{Jet}} < 1.5$

Charm jet cross sections vs E_T^{jet} and η^{jet}

DESY 10-083

Sensitivity to scale choice. Reasonable description with scale choice.


Measurement of F_2^{cc} and F_2^{bb}

$$F_{2,meas}^b(x_i, Q_i^2) = \frac{\sigma_{meas,i}}{\sigma_{theo,i}} \times F_{2,theo}^b(x_i, Q_i^2)$$

- Extraction of inclusive structure functions (F_L is small)
- Double differential cross section
- Use HVQDIS to calculate theoretical predictions
- Extrapolation to full phase space small for beauty
- Larger for charm, but reduced compared to exclusive methods because of low p_T track acceptance


Measurement of F_2^{bb}

HERA



- Beauty structure function versus Q^2 for fixed x
- Vertex methods between H1 and ZEUS agree
- Agreement also found with semi-leptonic analyses
- NNLO predictions available
- Some differences between theories
- Data well described

Measurement of F_2^{cc}



- Charm structure function vs Q^2 for fixed x
- Higher precision tests theory
- Differences between MSTW NNLO and NLO predictions for charm. NNLO somewhat better description than NLO
- CTEQ NLO describes data
- Data being used to complement D meson and semi-leptonic measurements in combination of HERA data (1159 Corradi)


Summary

- Heavy Flavour production at HERA is a vital testing ground for **perturbative QCD**
- Vertex detectors are a **powerful tool** to extract heavy flavour cross sections
- In general a **good description** is provided by pQCD
- The vertexing method allows to make measurements of the contribution of heavy flavours to the proton structure function. Charm data precision **provides constraint** for theory. Beauty well described.
- Better discrimination to come from **combination** of results.


Extra Slides

Flavour Tagging - Vertex Detectors

ZEUS tracking (MicroVertexDetector)




H1
CentralSiliconTracker




H1 and ZEUS vertex detectors:

- Multi-layered single and double sided silicon microstrip detectors
- Combine precise spatial information from vertex detectors with tracks from central drift chambers
- Resolution of impact parameter in transverse plane $< 100 \mu\text{m}$




Half Wheel



Barrel module

Fitting Flavour Fractions


ZEUS



Example of ZEUS 2D M_{VTX} and S_L fitting for DIS

Measurement of F_2^{bb}


HERA




Comparison of vertexing results with semi-leptonic

Comparison with Muon Tagged Data

H1 Beauty Jet Cross section



H1 Beauty Breit Frame Jet Cross section



- Extrapolate muon data to full phase space (small uncertainty)
- H1 and ZEUS data from muon tagging lie systematically above vertex data at either high or low Q^2