

## Dark pion DM : WIMP vs. SIMP

*Wednesday, 17 April 2019 15:40 (25 minutes)*

Scale invariant extension of the SM with QCD-like strongly interacting hidden (dark) sector is interesting, since the dimensional transmutation and chiral symmetry breaking in the hidden sector could be the origin of electroweak symmetry breaking (EWSB), and all the masses of the SM particles as well as dark pions and dark baryons that could be good cold dark matter candidates. In this talk I discuss dark pion DM as WIMP vs. SIMP. Ignoring the West-Zumino-Witten (WZW) interaction, I first discuss dark pion as a WIMP using two different approaches, the chiral perturbation theory (ChPT) and AdS/QCD. Then I include the WZW interaction and discuss dark pion within SIMP scenario. However, the analysis based on ChPT indicates that the viable parameter space for SIMP seems to be outside the validity region of ChPT. I show that this problem can be resolved if we include dark vector mesons, and the SIMP idea can be realized in the dark pion sector.

Based on :

arXiv:1103.2571 WIMP scenario within Chiral perturbation (PRL)

arXiv:1606.02969 WIMP scenario within AdS/QCD (JHEP)

arXiv:1801.07726 SIMP scenario including dark vector mesons  
and Wess-Zumino-Witten term (PRD)

**Primary author:** Prof. KO, Pyungwon (Korea Institute for Advanced Study (KIAS) )

**Presenter:** Prof. KO, Pyungwon (Korea Institute for Advanced Study (KIAS) )

**Session Classification:** Astroparticle and cosmology

**Track Classification:** Astroparticle and cosmology