

Reconstructing (s)tops using fat jets

Michihisa Takeuchi
(Heidelberg University)

with Tilman Plehn, Michael Spannowsky, Dirk Zerwas

arxiv: 1006.2833[hep-ph]

Top tagger for low p_T range

Top quark: strongly related to EWSB sector

top tagging at LHC: important for new physics search

Hadronic top : full momentum reconstruction possible

Inspired by very heavy $X \rightarrow t\bar{t}$, several top taggers available

[D.E.Kaplan, K.Rehermann, M.D.Schwartz and B.Tweedie]

[D. Krohn, J. Thaler, L.T. Wang]

[L.G. Almeida, S.J. Lee, G. Perez, G. Sterman, I. Sung]

Look into jet substructure.

Designed for $p_T > 500 \text{ GeV}$,
not expected in the SM.


For example Standard Model



Establish top tagger

in low p_T range (200 ~ 500 GeV)

(Same for $pp \rightarrow \tilde{t}_1 \tilde{t}_1^* \rightarrow (t \tilde{\chi}_1^0) (\bar{t} \tilde{\chi}_1^0)$)




Fat jets


Top Tagger: C/A algorithm with large R

top: heavy mass \rightarrow with modest p_T , decay products well separated.

C/A distance at parton level



Fraction of tops within various R



R=1.5 to have top ~ 200 GeV inside fat jet.

Algorithm



1. Find fat jets using C / A algorithm with $R=1.5$, $p_T > 200$ GeV

Large R collects QCD (ISR, FSR, UE and other jets).

2. Find hard objects using mass drop criterion

hard objects inside fat jet by mass drop criterion:

Undoing clustering, $m_{j_1} < 0.8 m_j$ to keep j_1 and j_2 . stop when $m_j < 30$ GeV



(mass drop in $t \rightarrow Wb$ and $W \rightarrow jj$)

3. Filter and choose pairing

Take 3 sub-jets, re-cluster with R_{filt} and compute mass of first 5

keep pairing with best top mass.

top candidate

$$|m_{jjj} - 172.3 \text{ GeV}| < 25 \text{ GeV}$$

$$p_T^{\text{rec}} > 200 \text{ GeV}$$

No b-tag, no W-mass cut yet.


tt vs. QCD, W+jets

4. check mass ratios

Cluster top candidate constituents into 3 subjets p_1, p_2, p_3

$$m_{12}, m_{13}, m_{23} \quad m_t^2 = m_{123}^2 = m_{12}^2 + m_{13}^2 + m_{23}^2$$


After imposing top mass cut, 2 independent mass ratios.



$$R_{\min} < \frac{m_{23}}{m_{123}} < R_{\max} \quad \text{and} \quad 0.2 < \arctan \frac{m_{13}}{m_{12}} < 1.3$$

$$R_{\min}^2 \left(1 + \left(\frac{m_{13}}{m_{12}} \right)^2 \right) < 1 - \left(\frac{m_{23}}{m_{123}} \right)^2 < R_{\max}^2 \left(1 + \left(\frac{m_{13}}{m_{12}} \right)^2 \right) \quad \text{and} \quad \frac{m_{23}}{m_{123}} > 0.35$$

$$R_{\min}^2 \left(1 + \left(\frac{m_{12}}{m_{13}} \right)^2 \right) < 1 - \left(\frac{m_{23}}{m_{123}} \right)^2 < R_{\max}^2 \left(1 + \left(\frac{m_{12}}{m_{13}} \right)^2 \right) \quad \text{and} \quad \frac{m_{23}}{m_{123}} > 0.35$$




tt vs. QCD, W+jets

4. check mass ratios

Cluster top candidate constituents into 3 subjets p_1, p_2, p_3

$$m_{12}, m_{13}, m_{23} \quad m_t^2 = m_{123}^2 = m_{12}^2 + m_{13}^2 + m_{23}^2$$

After imposing top mass cut, 2 independent mass ratios.



$$R_{\min} < \frac{m_{23}}{m_{123}} < R_{\max} \quad \text{and} \quad 0.2 < \arctan \frac{m_{13}}{m_{12}} < 1.3$$


$$R_{\min}^2 \left(1 + \left(\frac{m_{13}}{m_{12}} \right)^2 \right) < 1 - \left(\frac{m_{23}}{m_{123}} \right)^2 < R_{\max}^2 \left(1 + \left(\frac{m_{13}}{m_{12}} \right)^2 \right) \quad \text{and} \quad \frac{m_{23}}{m_{123}} > 0.35$$

$$R_{\min}^2 \left(1 + \left(\frac{m_{12}}{m_{13}} \right)^2 \right) < 1 - \left(\frac{m_{23}}{m_{123}} \right)^2 < R_{\max}^2 \left(1 + \left(\frac{m_{12}}{m_{13}} \right)^2 \right) \quad \text{and} \quad \frac{m_{23}}{m_{123}} > 0.35$$

W mass

$$R_{\min} = 85\% \frac{m_W}{m_t}$$

$$R_{\max} = 115\% \frac{m_W}{m_t}$$




tt vs. QCD, W+jets

4. check mass ratios

Cluster top candidate constituents into 3 subjets p_1, p_2, p_3

$$m_{12}, m_{13}, m_{23} \quad m_t^2 = m_{123}^2 = m_{12}^2 + m_{13}^2 + m_{23}^2$$

After imposing top mass cut, 2 independent mass ratios.



$$R_{\min} < \frac{m_{23}}{m_{123}} < R_{\max} \quad \text{and} \quad 0.2 < \arctan \frac{m_{13}}{m_{12}} < 1.3$$

$$R_{\min}^2 \left(1 + \left(\frac{m_{13}}{m_{12}} \right)^2 \right) < 1 - \left(\frac{m_{23}}{m_{123}} \right)^2 < R_{\max}^2 \left(1 + \left(\frac{m_{13}}{m_{12}} \right)^2 \right) \quad \text{and} \quad \frac{m_{23}}{m_{123}} > 0.35$$


$$R_{\min}^2 \left(1 + \left(\frac{m_{12}}{m_{13}} \right)^2 \right) < 1 - \left(\frac{m_{23}}{m_{123}} \right)^2 < R_{\max}^2 \left(1 + \left(\frac{m_{12}}{m_{13}} \right)^2 \right) \quad \text{and} \quad \frac{m_{23}}{m_{123}} > 0.35$$

W mass



$$R_{\min} = 85\% \frac{m_W}{m_t}$$

$$R_{\max} = 115\% \frac{m_W}{m_t}$$

soft/collinear




Efficiencies




~40% hadronic tops tagged, 2~4% mis-tag.

	$t\bar{t}$ QCD W+jets					$t\bar{t}$ QCD W+jets					
$p_{T,t}^{\min}$ [GeV]	0	200	300			0	200	300			
one fat jet	92200	36100	8250	$4.10 \cdot 10^7$	$3.19 \cdot 10^5$	100%	100%	100%	100%	100%	
two fat jets	40700	20300	5810	$2.16 \cdot 10^7$	$1.60 \cdot 10^5$	44%	57%	70%	53%	50%	relative to one fat jet
one top tag	20900	13400	4160	$8.18 \cdot 10^5$	$1.27 \cdot 10^4$	23%	37%	51%	2.0%	3.9%	relative to one fat jet
two top tags	1880	1630	700	11000	233	2.0%	4.5%	8.5%	0.027%	0.07%	relative to one fat jet
						4.5%	8.0%	12%	0.05%	0.15%	relative to two fat jets

Parton-jet matching



95% in $\Delta R < 0.5$



80% in $\Delta p/p^{\text{rec}} < 20\%$

momentum reconstructed well

stop pairs

$$pp \rightarrow \tilde{t}_1 \tilde{t}_1^* \rightarrow (t \tilde{\chi}_1^0)(\bar{t} \tilde{\chi}_1^0) \rightarrow (bjj \tilde{\chi}_1^0)(\bar{b}jj \tilde{\chi}_1^0)$$

cuts:

2 fat jets: $p_{T,j} > 200/200$ GeV

lepton veto

$\cancel{p}_T > 150$ GeV

2 tagged tops: $p_T^{\text{rec}} > 200/200$ GeV

b tag for 1st tagged top

$m_{T2} > 250$ GeV

	$\tilde{t}_1 \tilde{t}_1^*$						$t\bar{t}$	QCD	W +jets	Z +jets	S/B	$S/\sqrt{B}_{10 \text{ fb}^{-1}}$
$m_{\tilde{t}}[\text{GeV}]$	340	390	440	490	540	640						340
$p_{T,j} > 200$ GeV, ℓ veto	728	447	292	187	124	46	87850	$2.4 \cdot 10^7$	$1.6 \cdot 10^5$	n/a	$3.0 \cdot 10^{-5}$	
$\cancel{p}_T > 150$ GeV	283	234	184	133	93	35	2245	$2.4 \cdot 10^5$	1710	2240	$1.2 \cdot 10^{-3}$	
first top tag	100	91	75	57	42	15	743	7590	90	114	$1.2 \cdot 10^{-2}$	
second top tag	15	12.4	11	8.4	6.3	2.3	32	129	5.7	1.4	$8.3 \cdot 10^{-2}$	
b tag	8.7	7.4	6.3	5.0	3.8	1.4	19	2.6	$\lesssim 0.2$	$\lesssim 0.05$	0.40	5.9
$m_{T2} > 250$ GeV	4.3	5.0	4.9	4.2	3.2	1.2	4.2	$\lesssim 0.6$	$\lesssim 0.1$	$\lesssim 0.03$	0.88	6.1

top tag : W + jets, Z + jets negligible

b tag : QCD negligible

m_{T2} : reduce $t\bar{t}$

Summary

Focus on low pt tops (200 to 500 GeV)


Fat jets killing combinatorics

Efficiency: top: ~40%, W+jets: 4%, QCD: 2%

Top momentum reconstructed



stop pairs: $S/B \sim 1$ and $S/\sqrt{B} \sim 5$ with 10 fb^{-1} .

Back up (R_bjj)



$$R_{bjj} = \max\{R_1, R_2\}$$

Back up



Back up

