

Statistics for Particle Physics

Lecture 1: Fundamentals

<https://indico.cern.ch/event/747653/>



INSIGHTS Workshop on
Statistics and Machine Learning

CERN, 17-21 September, 2018



Glen Cowan
Physics Department
Royal Holloway, University of London
g.cowan@rhul.ac.uk
www.pp.rhul.ac.uk/~cowan

Outline

→ Lecture 1: Fundamentals

Probability

Random variables, pdfs

Lecture 2: Statistical tests

Formalism of frequentist tests

Comments on multivariate methods (brief)

p-values

Discovery and limits

Lecture 3: Parameter estimation

Properties of estimators

Maximum likelihood


Some statistics books, papers, etc.

- G. Cowan, *Statistical Data Analysis*, Clarendon, Oxford, 1998
- R.J. Barlow, *Statistics: A Guide to the Use of Statistical Methods in the Physical Sciences*, Wiley, 1989
- Ilya Narsky and Frank C. Porter, *Statistical Analysis Techniques in Particle Physics*, Wiley, 2014.
- Luca Lista, *Statistical Methods for Data Analysis in Particle Physics*, Springer, 2017.
- L. Lyons, *Statistics for Nuclear and Particle Physics*, CUP, 1986
- F. James., *Statistical and Computational Methods in Experimental Physics*, 2nd ed., World Scientific, 2006
- S. Brandt, *Statistical and Computational Methods in Data Analysis*, Springer, New York, 1998 (with program library on CD)
- M. Tanabashi et al. (PDG), Phys. Rev. D 98, 030001 (2018); see also **pdg.lbl.gov** sections on probability, statistics, Monte Carlo

Theory \leftrightarrow Statistics \leftrightarrow Experiment


Theory (model, hypothesis):

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i \bar{\psi} \not{D} \psi + \dots$$




$$\sigma = \frac{G_F \alpha_s^2 m_H^2}{288 \sqrt{2\pi}} \times \dots$$


+ simulation
of detector
and cuts



Experiment:



+ data
selection



Data analysis in particle physics

Observe events (e.g., pp collisions) and for each, measure a set of characteristics:

particle momenta, number of muons, energy of jets,...

Compare observed distributions of these characteristics to predictions of theory. From this, we want to:

Estimate the free parameters of the theory: $m_H = 125.4$

Quantify the uncertainty in the estimates: $\pm 0.4 \text{ GeV}$

Assess how well a given theory stands in agreement with the observed data:

0^+ good, 2^+ bad

To do this we need a clear definition of PROBABILITY

A definition of probability

Consider a set S with subsets A, B, \dots

For all $A \subset S, P(A) \geq 0$

$$P(S) = 1$$

If $A \cap B = \emptyset, P(A \cup B) = P(A) + P(B)$



Kolmogorov
axioms (1933)

Also define conditional probability of A given B :

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Subsets A, B independent if: $P(A \cap B) = P(A)P(B)$

If A, B independent, $P(A|B) = \frac{P(A)P(B)}{P(B)} = P(A)$

Interpretation of probability

I. Relative frequency

A, B, \dots are outcomes of a repeatable experiment

$$P(A) = \lim_{n \rightarrow \infty} \frac{\text{times outcome is } A}{n}$$

cf. quantum mechanics, particle scattering, radioactive decay...

II. Subjective probability

A, B, \dots are hypotheses (statements that are true or false)

$$P(A) = \text{degree of belief that } A \text{ is true}$$

- Both interpretations consistent with Kolmogorov axioms.
- In particle physics frequency interpretation often most useful, but subjective probability can provide more natural treatment of non-repeatable phenomena:

systematic uncertainties, probability that Higgs boson exists,...

Bayes' theorem


From the definition of conditional probability we have,

$$P(A|B) = \frac{P(A \cap B)}{P(B)} \quad \text{and} \quad P(B|A) = \frac{P(B \cap A)}{P(A)}$$

but $P(A \cap B) = P(B \cap A)$, so

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Bayes' theorem




First published (posthumously) by the
Reverend Thomas Bayes (1702–1761)

*An essay towards solving a problem in the
doctrine of chances*, Philos. Trans. R. Soc. 53
(1763) 370; reprinted in Biometrika, 45 (1958) 293.

The law of total probability

Consider a subset B of the sample space S ,

divided into disjoint subsets A_i such that $\cup_i A_i = S$,



$$\rightarrow B = B \cap S = B \cap (\cup_i A_i) = \cup_i (B \cap A_i), \quad B \cap A_i$$

$$\rightarrow P(B) = P(\cup_i (B \cap A_i)) = \sum_i P(B \cap A_i)$$

$$\rightarrow P(B) = \sum_i P(B|A_i)P(A_i) \quad \text{law of total probability}$$

Bayes' theorem becomes

$$P(A|B) = \frac{P(B|A)P(A)}{\sum_i P(B|A_i)P(A_i)}$$

An example using Bayes' theorem

Suppose the probability (for anyone) to have a disease D is:

$$\begin{aligned} P(D) &= 0.001 && \leftarrow \text{prior probabilities, i.e.,} \\ P(\text{no D}) &= 0.999 && \text{before any test carried out} \end{aligned}$$

Consider a test for the disease: result is + or -

$$\begin{aligned} P(+|D) &= 0.98 && \leftarrow \text{probabilities to (in)correctly} \\ P(-|D) &= 0.02 && \text{identify a person with the disease} \end{aligned}$$

$$\begin{aligned} P(+|\text{no D}) &= 0.03 && \leftarrow \text{probabilities to (in)correctly} \\ P(-|\text{no D}) &= 0.97 && \text{identify a healthy person} \end{aligned}$$

Suppose your result is +. How worried should you be?

Bayes' theorem example (cont.)

The probability to have the disease given a + result is

$$\begin{aligned} p(D|+) &= \frac{P(+|D)P(D)}{P(+|D)P(D) + P(+|\text{no } D)P(\text{no } D)} \\ &= \frac{0.98 \times 0.001}{0.98 \times 0.001 + 0.03 \times 0.999} \\ &= 0.032 \quad \leftarrow \text{posterior probability} \end{aligned}$$

i.e. you're probably OK!

Your viewpoint: my degree of belief that I have the disease is 3.2%.

Your doctor's viewpoint: 3.2% of people like this have the disease.

Frequentist Statistics – general philosophy

In frequentist statistics, probabilities are associated only with the data, i.e., outcomes of repeatable observations (shorthand: \vec{x}).

Probability = limiting frequency

Probabilities such as

P (Higgs boson exists),

P ($0.117 < \alpha_s < 0.121$),

etc. are either 0 or 1, but we don't know which.

The tools of frequentist statistics tell us what to expect, under the assumption of certain probabilities, about hypothetical repeated observations.

A hypothesis is preferred if the data are found in a region of high predicted probability (i.e., where an alternative hypothesis predicts lower probability).

Bayesian Statistics – general philosophy

In Bayesian statistics, use subjective probability for hypotheses:

probability of the data assuming hypothesis H (the likelihood)

prior probability, i.e., before seeing the data

$$P(H|\vec{x}) = \frac{P(\vec{x}|H)\pi(H)}{\int P(\vec{x}|H)\pi(H) dH}$$

posterior probability, i.e., after seeing the data

normalization involves sum over all possible hypotheses

Bayes' theorem has an “if-then” character: If your prior probabilities were $\pi(H)$, then it says how these probabilities should change in the light of the data.

No general prescription for priors (subjective!)

Random variables and probability density functions

A random variable is a numerical characteristic assigned to an element of the sample space; can be discrete or continuous.

Suppose outcome of experiment is continuous value x

$$P(x \text{ found in } [x, x + dx]) = f(x) dx$$

→ $f(x)$ = probability density function (pdf)

$$\int_{-\infty}^{\infty} f(x) dx = 1 \quad x \text{ must be somewhere}$$

Or for discrete outcome x_i with e.g. $i = 1, 2, \dots$ we have



$$P(x_i) = p_i \quad \text{probability mass function}$$

$$\sum_i P(x_i) = 1 \quad x \text{ must take on one of its possible values}$$

Cumulative distribution function

Probability to have outcome less than or equal to x is

$$\int_{-\infty}^x f(x') dx' \equiv F(x) \quad \text{cumulative distribution function}$$



Alternatively define pdf with $f(x) = \frac{\partial F(x)}{\partial x}$

Other types of probability densities

Outcome of experiment characterized by several values,
e.g. an n -component vector, (x_1, \dots, x_n)

→ joint pdf $f(x_1, \dots, x_n)$

Sometimes we want only pdf of some (or one) of the components

→ marginal pdf $f_1(x_1) = \int \dots \int f(x_1, \dots, x_n) dx_2 \dots dx_n$

x_1, x_2 independent if $f(x_1, x_2) = f_1(x_1)f_2(x_2)$

Sometimes we want to consider some components as constant


→ conditional pdf $g(x_1|x_2) = \frac{f(x_1, x_2)}{f_2(x_2)}$

Functions of a random variable

A function of a random variable is itself a random variable.

Suppose x follows a pdf $f(x)$, consider a function $a(x)$.

What is the pdf $g(a)$?



$$g(a) da = \int_{dS} f(x) dx$$

dS = region of x space for which a is in $[a, a+da]$.


For one-variable case with unique inverse this is simply

$$g(a) da = f(x) dx$$

$$\rightarrow g(a) = f(x(a)) \left| \frac{dx}{da} \right|$$

Functions without unique inverse

If inverse of $a(x)$ not unique,
include all dx intervals in dS
which correspond to da :



Example: $a = x^2, x = \pm\sqrt{a}, dx = \pm\frac{da}{2\sqrt{a}}$.

$$dS = \left[\sqrt{a}, \sqrt{a} + \frac{da}{2\sqrt{a}} \right] \cup \left[-\sqrt{a} - \frac{da}{2\sqrt{a}}, -\sqrt{a} \right]$$

$$g(a) = \frac{f(\sqrt{a})}{2\sqrt{a}} + \frac{f(-\sqrt{a})}{2\sqrt{a}}$$

Functions of more than one r.v.

Consider r.v.s $\vec{x} = (x_1, \dots, x_n)$ and a function $a(\vec{x})$.


$$g(a')da' = \int \dots \int_{dS} f(x_1, \dots, x_n) dx_1 \dots dx_n$$

dS = region of x -space between (hyper)surfaces defined by

$$a(\vec{x}) = a', \quad a(\vec{x}) = a' + da'$$

Functions of more than one r.v. (2)

Example: r.v.s $x, y > 0$ follow joint pdf $f(x,y)$,
consider the function $z = xy$. What is $g(z)$?



$$\begin{aligned} g(z) dz &= \int \dots \int_{dS} f(x, y) dx dy \\ &= \int_0^\infty dx \int_{z/x}^{(z+dz)/x} f(x, y) dy \\ \rightarrow g(z) &= \int_0^\infty f\left(x, \frac{z}{x}\right) \frac{dx}{x} \\ &= \int_0^\infty f\left(\frac{z}{y}, y\right) \frac{dy}{y} \end{aligned}$$

(Mellin convolution)

More on transformation of variables

Consider a random vector $\vec{x} = (x_1, \dots, x_n)$ with joint pdf $f(\vec{x})$.

Form n linearly independent functions $\vec{y}(\vec{x}) = (y_1(\vec{x}), \dots, y_n(\vec{x}))$ for which the inverse functions $x_1(\vec{y}), \dots, x_n(\vec{y})$ exist.

Then the joint pdf of the vector of functions is $g(\vec{y}) = |J|f(\vec{x})$

where J is the

Jacobian determinant:

$$J = \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \frac{\partial x_1}{\partial y_2} & \cdots & \frac{\partial x_1}{\partial y_n} \\ \frac{\partial x_2}{\partial y_1} & \frac{\partial x_2}{\partial y_2} & \cdots & \frac{\partial x_2}{\partial y_n} \\ \vdots & & & \vdots \\ & & \cdots & \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

For e.g. $g_1(y_1)$ integrate $g(\vec{y})$ over the unwanted components.

Expectation values

Consider continuous r.v. x with pdf $f(x)$.

Define expectation (mean) value as $E[x] = \int x f(x) dx$

Notation (often): $E[x] = \mu$ ~ “centre of gravity” of pdf.

For a function $y(x)$ with pdf $g(y)$,


$$E[y] = \int y g(y) dy = \int y(x) f(x) dx \quad (\text{equivalent})$$

Variance: $V[x] = E[x^2] - \mu^2 = E[(x - \mu)^2]$

Notation: $V[x] = \sigma^2$

Standard deviation: $\sigma = \sqrt{\sigma^2}$

$\sigma \sim$ width of pdf, same units as x .



Covariance and correlation

Define covariance $\text{cov}[x,y]$ (also use matrix notation V_{xy}) as

$$\text{cov}[x, y] = E[xy] - \mu_x \mu_y = E[(x - \mu_x)(y - \mu_y)]$$

Correlation coefficient (dimensionless) defined as

$$\rho_{xy} = \frac{\text{cov}[x, y]}{\sigma_x \sigma_y}$$

If x, y , independent, i.e., $f(x, y) = f_x(x)f_y(y)$, then


$$E[xy] = \int \int xy f(x, y) dx dy = \mu_x \mu_y$$

→ $\text{cov}[x, y] = 0$ x and y , ‘uncorrelated’


N.B. converse not always true.

Correlation (cont.)


$$\rho = 0.75$$




$$\rho = 0.95$$



$$\rho = -0.75$$



$$\rho = 0.25$$



Error propagation

Suppose we measure a set of values $\vec{x} = (x_1, \dots, x_n)$

and we have the covariances $V_{ij} = \text{cov}[x_i, x_j]$

which quantify the measurement errors in the x_i .

Now consider a function $y(\vec{x})$.

What is the variance of $y(\vec{x})$?

The hard way: use joint pdf $f(\vec{x})$ to find the pdf $g(y)$,

then from $g(y)$ find $V[y] = E[y^2] - (E[y])^2$.

Often not practical, $f(\vec{x})$ may not even be fully known.

Error propagation (2)

Suppose we had $\vec{\mu} = E[\vec{x}]$

in practice only estimates given by the measured \vec{x}

Expand $y(\vec{x})$ to 1st order in a Taylor series about $\vec{\mu}$

$$y(\vec{x}) \approx y(\vec{\mu}) + \sum_{i=1}^n \left[\frac{\partial y}{\partial x_i} \right]_{\vec{x}=\vec{\mu}} (x_i - \mu_i)$$

To find $V[y]$ we need $E[y^2]$ and $E[y]$.

$$E[y(\vec{x})] \approx y(\vec{\mu}) \text{ since } E[x_i - \mu_i] = 0$$

Error propagation (3)

$$\begin{aligned} E[y^2(\vec{x})] &\approx y^2(\vec{\mu}) + 2y(\vec{\mu}) \sum_{i=1}^n \left[\frac{\partial y}{\partial x_i} \right]_{\vec{x}=\vec{\mu}} E[x_i - \mu_i] \\ &+ E \left[\left(\sum_{i=1}^n \left[\frac{\partial y}{\partial x_i} \right]_{\vec{x}=\vec{\mu}} (x_i - \mu_i) \right) \left(\sum_{j=1}^n \left[\frac{\partial y}{\partial x_j} \right]_{\vec{x}=\vec{\mu}} (x_j - \mu_j) \right) \right] \\ &= y^2(\vec{\mu}) + \sum_{i,j=1}^n \left[\frac{\partial y}{\partial x_i} \frac{\partial y}{\partial x_j} \right]_{\vec{x}=\vec{\mu}} V_{ij} \end{aligned}$$

Putting the ingredients together gives the variance of $y(\vec{x})$

$$\sigma_y^2 \approx \sum_{i,j=1}^n \left[\frac{\partial y}{\partial x_i} \frac{\partial y}{\partial x_j} \right]_{\vec{x}=\vec{\mu}} V_{ij}$$

Error propagation (4)

If the x_i are uncorrelated, i.e., $V_{ij} = \sigma_i^2 \delta_{ij}$, then this becomes

$$\sigma_y^2 \approx \sum_{i=1}^n \left[\frac{\partial y}{\partial x_i} \right]_{\vec{x}=\vec{\mu}}^2 \sigma_i^2$$

Similar for a set of m functions $\vec{y}(\vec{x}) = (y_1(\vec{x}), \dots, y_m(\vec{x}))$

$$U_{kl} = \text{cov}[y_k, y_l] \approx \sum_{i,j=1}^n \left[\frac{\partial y_k}{\partial x_i} \frac{\partial y_l}{\partial x_j} \right]_{\vec{x}=\vec{\mu}} V_{ij}$$


or in matrix notation $U = A V A^T$, where

$$A_{ij} = \left[\frac{\partial y_i}{\partial x_j} \right]_{\vec{x}=\vec{\mu}}$$

Error propagation (5)


The ‘error propagation’ formulae tell us the covariances of a set of functions

$\vec{y}(\vec{x}) = (y_1(\vec{x}), \dots, y_m(\vec{x}))$ in terms of the covariances of the original variables.



Limitations: exact only if $\vec{y}(\vec{x})$ linear.

Approximation breaks down if function nonlinear over a region comparable in size to the σ_i .



N.B. We have said nothing about the exact pdf of the x_i , e.g., it doesn’t have to be Gaussian.

Error propagation – special cases

$$y = x_1 + x_2 \rightarrow \sigma_y^2 = \sigma_1^2 + \sigma_2^2 + 2\text{cov}[x_1, x_2]$$

$$y = x_1 x_2 \rightarrow \frac{\sigma_y^2}{y^2} = \frac{\sigma_1^2}{x_1^2} + \frac{\sigma_2^2}{x_2^2} + 2\frac{\text{cov}[x_1, x_2]}{x_1 x_2}$$

That is, if the x_i are uncorrelated:

add errors quadratically for the sum (or difference),
add relative errors quadratically for product (or ratio).



But correlations can change this completely...

Error propagation – special cases (2)

Consider $y = x_1 - x_2$ with

$$\mu_1 = \mu_2 = 10, \quad \sigma_1 = \sigma_2 = 1, \quad \rho = \frac{\text{cov}[x_1, x_2]}{\sigma_1 \sigma_2} = 0.$$

$$V[y] = 1^2 + 1^2 = 2, \rightarrow \sigma_y = 1.4$$

Now suppose $\rho = 1$. Then

$$V[y] = 1^2 + 1^2 - 2 = 0, \rightarrow \sigma_y = 0$$

i.e. for 100% correlation, error in difference $\rightarrow 0$.

Some distributions

<u>Distribution/pdf</u>	<u>Example use in HEP</u>
Binomial	Branching ratio
Multinomial	Histogram with fixed N
Poisson	Number of events found
Uniform	Monte Carlo method
Exponential	Decay time
Gaussian	Measurement error
Chi-square	Goodness-of-fit
Cauchy	Mass of resonance
Landau	Ionization energy loss
Beta	Prior pdf for efficiency
Gamma	Sum of exponential variables
Student's t	Resolution function with adjustable tails

Binomial distribution

Consider N independent experiments (Bernoulli trials):

outcome of each is ‘success’ or ‘failure’,
probability of success on any given trial is p .

Define discrete r.v. n = number of successes ($0 \leq n \leq N$).

Probability of a specific outcome (in order), e.g. ‘ssfsf’ is

$$pp(1-p)p(1-p) = p^n(1-p)^{N-n}$$

But order not important; there are

$$\frac{N!}{n!(N-n)!}$$

ways (permutations) to get n successes in N trials, total probability for n is sum of probabilities for each permutation.

Binomial distribution (2)

The binomial distribution is therefore

$$f(n; N, p) = \frac{N!}{n!(N-n)!} p^n (1-p)^{N-n}$$

random variable parameters







For the expectation value and variance we find:

$$E[n] = \sum_{n=0}^N n f(n; N, p) = Np$$

$$V[n] = E[n^2] - (E[n])^2 = Np(1-p)$$

Binomial distribution (3)

Binomial distribution for several values of the parameters:



Example: observe N decays of W^\pm , the number n of which are $W \rightarrow \mu\nu$ is a binomial r.v., p = branching ratio.

Multinomial distribution

Like binomial but now m outcomes instead of two, probabilities are

$$\vec{p} = (p_1, \dots, p_m), \quad \text{with } \sum_{i=1}^m p_i = 1.$$

For N trials we want the probability to obtain:

n_1 of outcome 1,

n_2 of outcome 2,

⋮

n_m of outcome m .

This is the multinomial distribution for $\vec{n} = (n_1, \dots, n_m)$

$$f(\vec{n}; N, \vec{p}) = \frac{N!}{n_1! n_2! \cdots n_m!} p_1^{n_1} p_2^{n_2} \cdots p_m^{n_m}$$

Multinomial distribution (2)

Now consider outcome i as ‘success’, all others as ‘failure’.

→ all n_i individually binomial with parameters N, p_i

$$E[n_i] = Np_i, \quad V[n_i] = Np_i(1 - p_i) \quad \text{for all } i$$

One can also find the covariance to be

$$V_{ij} = Np_i(\delta_{ij} - p_j)$$

Example: $\vec{n} = (n_1, \dots, n_m)$ represents a histogram
with m bins, N total entries, all entries independent.

Poisson distribution

Consider binomial n in the limit




$$N \rightarrow \infty, \quad p \rightarrow 0, \quad E[n] = Np \rightarrow \nu .$$

→ n follows the Poisson distribution:

$$f(n; \nu) = \frac{\nu^n}{n!} e^{-\nu} \quad (n \geq 0)$$

$$E[n] = \nu, \quad V[n] = \nu .$$

Example: number of scattering events n with cross section σ found for a fixed integrated luminosity, with $\nu = \sigma \int L dt$.




Uniform distribution

Consider a continuous r.v. x with $-\infty < x < \infty$. Uniform pdf is:

$$f(x; \alpha, \beta) = \begin{cases} \frac{1}{\beta - \alpha} & \alpha \leq x \leq \beta \\ 0 & \text{otherwise} \end{cases}$$

$$E[x] = \frac{1}{2}(\alpha + \beta)$$

$$V[x] = \frac{1}{12}(\beta - \alpha)^2$$



N.B. For any r.v. x with cumulative distribution $F(x)$, $y = F(x)$ is uniform in $[0,1]$.

Example: for $\pi^0 \rightarrow \gamma\gamma$, E_γ is uniform in $[E_{\min}, E_{\max}]$, with

$$E_{\min} = \frac{1}{2}E_\pi(1 - \beta), \quad E_{\max} = \frac{1}{2}E_\pi(1 + \beta)$$


Exponential distribution

The exponential pdf for the continuous r.v. x is defined by:

$$f(x; \xi) = \begin{cases} \frac{1}{\xi} e^{-x/\xi} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

$$E[x] = \xi$$

$$V[x] = \xi^2$$



Example: proper decay time t of an unstable particle

$$f(t; \tau) = \frac{1}{\tau} e^{-t/\tau} \quad (\tau = \text{mean lifetime})$$


Lack of memory (unique to exponential): $f(t - t_0 | t \geq t_0) = f(t)$

Gaussian distribution

The Gaussian (normal) pdf for a continuous r.v. x is defined by:

$$f(x; \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$$

$$\begin{aligned} E[x] &= \mu && \text{(N.B. often } \mu, \sigma^2 \text{ denote mean, variance of any r.v., not only Gaussian.)} \\ V[x] &= \sigma^2 \end{aligned}$$



Special case: $\mu = 0, \sigma^2 = 1$ ('standard Gaussian'):

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}, \quad \Phi(x) = \int_{-\infty}^x \varphi(x') dx'$$

If $y \sim \text{Gaussian with } \mu, \sigma^2$, then $x = (y - \mu) / \sigma$ follows $\varphi(x)$.

Gaussian pdf and the Central Limit Theorem

The Gaussian pdf is so useful because almost any random variable that is a sum of a large number of small contributions follows it. This follows from the Central Limit Theorem:

For n independent r.v.s x_i with finite variances σ_i^2 , otherwise arbitrary pdfs, consider the sum

$$y = \sum_{i=1}^n x_i$$

In the limit $n \rightarrow \infty$, y is a Gaussian r.v. with

$$E[y] = \sum_{i=1}^n \mu_i \quad V[y] = \sum_{i=1}^n \sigma_i^2$$

Measurement errors are often the sum of many contributions, so frequently measured values can be treated as Gaussian r.v.s.

Central Limit Theorem (2)

The CLT can be proved using characteristic functions (Fourier transforms), see, e.g., SDA Chapter 10.

For finite n , the theorem is approximately valid to the extent that the fluctuation of the sum is not dominated by one (or few) terms.



Beware of measurement errors with non-Gaussian tails.

Good example: velocity component v_x of air molecules.

OK example: total deflection due to multiple Coulomb scattering.
(Rare large angle deflections give non-Gaussian tail.)

Bad example: energy loss of charged particle traversing thin gas layer. (Rare collisions make up large fraction of energy loss, cf. Landau pdf.)

Multivariate Gaussian distribution

Multivariate Gaussian pdf for the vector $\vec{x} = (x_1, \dots, x_n)$:

$$f(\vec{x}; \vec{\mu}, V) = \frac{1}{(2\pi)^{n/2}|V|^{1/2}} \exp \left[-\frac{1}{2}(\vec{x} - \vec{\mu})^T V^{-1} (\vec{x} - \vec{\mu}) \right]$$

\vec{x} , $\vec{\mu}$ are column vectors, \vec{x}^T , $\vec{\mu}^T$ are transpose (row) vectors,

$$E[x_i] = \mu_i, , \quad \text{cov}[x_i, x_j] = V_{ij} .$$

For $n = 2$ this is

$$\begin{aligned} f(x_1, x_2; \mu_1, \mu_2, \sigma_1, \sigma_2, \rho) &= \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \\ &\times \exp \left\{ -\frac{1}{2(1-\rho^2)} \left[\left(\frac{x_1 - \mu_1}{\sigma_1} \right)^2 + \left(\frac{x_2 - \mu_2}{\sigma_2} \right)^2 - 2\rho \left(\frac{x_1 - \mu_1}{\sigma_1} \right) \left(\frac{x_2 - \mu_2}{\sigma_2} \right) \right] \right\} \end{aligned}$$

where $\rho = \text{cov}[x_1, x_2]/(\sigma_1\sigma_2)$ is the correlation coefficient.


Chi-square (χ^2) distribution

The chi-square pdf for the continuous r.v. z ($z \geq 0$) is defined by

$$f(z; n) = \frac{1}{2^{n/2}\Gamma(n/2)} z^{n/2-1} e^{-z/2}$$

$n = 1, 2, \dots$ = number of ‘degrees of freedom’ (dof)

$$E[z] = n, \quad V[z] = 2n.$$



For independent Gaussian x_i , $i = 1, \dots, n$, means μ_i , variances σ_i^2 ,

$$z = \sum_{i=1}^n \frac{(x_i - \mu_i)^2}{\sigma_i^2} \quad \text{follows } \chi^2 \text{ pdf with } n \text{ dof.}$$


Example: goodness-of-fit test variable especially in conjunction with method of least squares.

Cauchy (Breit-Wigner) distribution

The Breit-Wigner pdf for the continuous r.v. x is defined by

$$f(x; \Gamma, x_0) = \frac{1}{\pi} \frac{\Gamma/2}{\Gamma^2/4 + (x - x_0)^2}$$

($\Gamma = 2$, $x_0 = 0$ is the Cauchy pdf.)



$E[x]$ not well defined, $V[x] \rightarrow \infty$.

x_0 = mode (most probable value)

Γ = full width at half maximum

Example: mass of resonance particle, e.g. ρ , K^* , ϕ^0 , ...

Γ = decay rate (inverse of mean lifetime)

Landau distribution


For a charged particle with $\beta = v/c$ traversing a layer of matter of thickness d , the energy loss Δ follows the Landau pdf:

$$f(\Delta; \beta) = \frac{1}{\xi} \phi(\lambda) ,$$

$$\phi(\lambda) = \frac{1}{\pi} \int_0^\infty \exp(-u \ln u - \lambda u) \sin \pi u \, du ,$$

$$\lambda = \frac{1}{\xi} \left[\Delta - \xi \left(\ln \frac{\xi}{\epsilon'} + 1 - \gamma_E \right) \right] ,$$

$$\xi = \frac{2\pi N_A e^4 z^2 \rho \sum Z}{m_e c^2 \sum A} \frac{d}{\beta^2} , \quad \epsilon' = \frac{I^2 \exp \beta^2}{2m_e c^2 \beta^2 \gamma^2} .$$




L. Landau, J. Phys. USSR **8** (1944) 201; see also

W. Allison and J. Cobb, Ann. Rev. Nucl. Part. Sci. **30** (1980) 253.

Landau distribution (2)

Long ‘Landau tail’
→ all moments ∞



Mode (most probable value) sensitive to β ,
→ particle i.d.


Beta distribution

$$f(x; \alpha, \beta) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1}$$

$$E[x] = \frac{\alpha}{\alpha + \beta}$$

$$V[x] = \frac{\alpha\beta}{(\alpha + \beta)^2(\alpha + \beta + 1)}$$

Often used to represent pdf of continuous r.v. nonzero only between finite limits.



Gamma distribution


$$f(x; \alpha, \beta) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-x/\beta}$$

$$E[x] = \alpha\beta$$

$$V[x] = \alpha\beta^2$$

Often used to represent pdf of continuous r.v. nonzero only in $[0, \infty]$.

Also e.g. sum of n exponential r.v.s or time until n th event in Poisson process \sim Gamma



Student's t distribution

$$f(x; \nu) = \frac{\Gamma\left(\frac{\nu+1}{2}\right)}{\sqrt{\nu\pi}\Gamma(\nu/2)} \left(1 + \frac{x^2}{\nu}\right)^{-\left(\frac{\nu+1}{2}\right)}$$


$$E[x] = 0 \quad (\nu > 1)$$

$$V[x] = \frac{\nu}{\nu - 2} \quad (\nu > 2)$$

ν = number of degrees of freedom
(not necessarily integer)

$\nu = 1$ gives Cauchy,

$\nu \rightarrow \infty$ gives Gaussian.



Extra slides

The Monte Carlo method


What it is: a numerical technique for calculating probabilities and related quantities using sequences of random numbers.

The usual steps:

- (1) Generate sequence r_1, r_2, \dots, r_m uniform in $[0, 1]$.
- (2) Use this to produce another sequence x_1, x_2, \dots, x_n distributed according to some pdf $f(x)$ in which we're interested (x can be a vector).
- (3) Use the x values to estimate some property of $f(x)$, e.g., fraction of x values with $a < x < b$ gives $\int_a^b f(x) dx$.
→ MC calculation = integration (at least formally)

MC generated values = ‘simulated data’

- use for testing statistical procedures



Random number generators

Goal: generate uniformly distributed values in $[0, 1]$.

Toss coin for e.g. 32 bit number... (too tiring).

→ ‘random number generator’

= computer algorithm to generate r_1, r_2, \dots, r_n .

Example: multiplicative linear congruential generator (MLCG)

$$n_{i+1} = (a n_i) \bmod m , \quad \text{where}$$

n_i = integer

a = multiplier

m = modulus

n_0 = seed (initial value)

N.B. mod = modulus (remainder), e.g. $27 \bmod 5 = 2$.

This rule produces a sequence of numbers n_0, n_1, \dots

Random number generators (2)

The sequence is (unfortunately) periodic!

Example (see Brandt Ch 4): $a = 3, m = 7, n_0 = 1$

$$n_1 = (3 \cdot 1) \bmod 7 = 3$$

$$n_2 = (3 \cdot 3) \bmod 7 = 2$$

$$n_3 = (3 \cdot 2) \bmod 7 = 6$$

$$n_4 = (3 \cdot 6) \bmod 7 = 4$$

$$n_5 = (3 \cdot 4) \bmod 7 = 5$$

$$n_6 = (3 \cdot 5) \bmod 7 = 1 \quad \leftarrow \text{sequence repeats}$$

Choose a, m to obtain long period (maximum = $m - 1$); m usually close to the largest integer that can be represented in the computer.

Only use a subset of a single period of the sequence.

Random number generators (3)

$r_i = n_i/m$ are in $[0, 1]$ but are they ‘random’?

Choose a, m so that the r_i pass various tests of randomness:


uniform distribution in $[0, 1]$,

all values independent (no correlations between pairs),

e.g. L’Ecuyer, Commun. ACM 31 (1988) 742 suggests

$$a = 40692$$

$$m = 2147483399$$





Far better generators available, e.g. **TRandom3**, based on Mersenne twister algorithm, period = $2^{19937} - 1$ (a “Mersenne prime”).

See F. James, Comp. Phys. Comm. 60 (1990) 111; Brandt Ch. 4

The transformation method

Given r_1, r_2, \dots, r_n uniform in $[0, 1]$, find x_1, x_2, \dots, x_n that follow $f(x)$ by finding a suitable transformation $x(r)$.



Require: $P(r \leq r') = P(x \leq x(r'))$

i.e. $\int_{-\infty}^{r'} g(r) dr = r' = \int_{-\infty}^{x(r')} f(x') dx' = F(x(r'))$


That is, set $F(x) = r$ and solve for $x(r)$.

Example of the transformation method

Exponential pdf: $f(x; \xi) = \frac{1}{\xi} e^{-x/\xi}$ ($x \geq 0$)


Set $\int_0^x \frac{1}{\xi} e^{-x'/\xi} dx' = r$ and solve for $x(r)$.

→ $x(r) = -\xi \ln(1 - r)$ ($x(r) = -\xi \ln r$ works too.)



The acceptance-rejection method

Enclose the pdf in a box:




- (1) Generate a random number x , uniform in $[x_{\min}, x_{\max}]$, i.e.
$$x = x_{\min} + r_1(x_{\max} - x_{\min}) , \quad r_1 \text{ is uniform in } [0,1].$$
- (2) Generate a 2nd independent random number u uniformly distributed between 0 and f_{\max} , i.e. $u = r_2 f_{\max}$.
- (3) If $u < f(x)$, then accept x . If not, reject x and repeat.

Example with acceptance-rejection method

$$f(x) = \frac{3}{8}(1 + x^2)$$
$$(-1 \leq x \leq 1)$$

If dot below curve, use x value in histogram.




Improving efficiency of the acceptance-rejection method

The fraction of accepted points is equal to the fraction of the box's area under the curve.

For very peaked distributions, this may be very low and thus the algorithm may be slow.

Improve by enclosing the pdf $f(x)$ in a curve $C h(x)$ that conforms to $f(x)$ more closely, where $h(x)$ is a pdf from which we can generate random values and C is a constant.



Generate points uniformly over $C h(x)$.

If point is below $f(x)$, accept x .


Monte Carlo event generators

Simple example: $e^+e^- \rightarrow \mu^+\mu^-$

Generate $\cos\theta$ and ϕ :

$$f(\cos\theta; A_{FB}) \propto (1 + \frac{8}{3}A_{FB}\cos\theta + \cos^2\theta) ,$$

$$g(\phi) = \frac{1}{2\pi} \quad (0 \leq \phi \leq 2\pi)$$



Less simple: ‘event generators’ for a variety of reactions:

$e^+e^- \rightarrow \mu^+\mu^-$, hadrons, ...

$pp \rightarrow$ hadrons, D-Y, SUSY,...

e.g. PYTHIA, HERWIG, ISAJET...

Output = ‘events’, i.e., for each event we get a list of generated particles and their momentum vectors, types, etc.

A simulated event

Event listing (summary)								
I particle/jet KS	KF	orig	P_X	P_Y	P_Z	E	m	
1 !p+!	21	2212	0	0.000	0.000	7000.000	7000.000	0.938
2 !p+!	21	2212	0	0.000	0.000	-7000.000	7000.000	0.938
3 !g!	21	21	1	0.863	-0.323	1739.862	1739.862	0.000
4 !ubar!	21	-2	2	-0.621	-0.163	-777.415	777.415	0.000
5 !g!	21	21	3	-2.427	5.486	1487.857	1487.	
6 !g!	21	21	4	-62.910	63.357	-463.274	471.	
7 !^g!	21	1000021	0	314.363	544.843	498.897	979.	
8 !^g!	21	1000021	0	-379.700	-476.000	525.686	980.	
9 !~chi_1-!	21	-1000024	7	130.058	112.247	129.860	263.	
10 !sbar!	21	-3	7	259.400	187.468	83.100	330	
11 !c!	21	4	7	-79.403	242.409	283.026	381.	
12 !~chi_20!	21	1000023	8	-326.241	-80.971	113.712	385.	
13 !b!	21	5	8	-51.841	-294.077	389.853	491.	
14 !bbar!	21	-5	8	-0.597	-99.577	21.299	101.	
15 !~chi_10!	21	1000022	9	103.352	81.316	83.457	175.	
16 !s!	21	3	9	5.451	38.374	52.302	65.	
17 !cbar!	21	-4	9	20.839	-7.250	-5.938	22.	
18 !~chi_10!	21	1000022	12	-136.266	-72.961	53.246	181.	
19 !nu_mu!	21	14	12	-78.263	-24.757	21.719	84.	
20 !nu_mubar!	21	-14	12	-107.801	16.901	38.226	115.	
21 gamma	1	22	4	2.636	1.357	0.125	2.	
22 (~chi_1-)	11	-1000024	9	129.643	112.440	129.820	262.	
23 (~chi_20)	11	1000023	12	-322.330	-80.817	113.191	382.	
24 ~chi_10	1	1000022	15	97.944	77.819	80.917	169.	
25 ~chi_10	1	1000022	18	-136.266	-72.961	53.246	181.	
26 nu_mu	1	14	19	-78.263	-24.757	21.719	84.	
27 nu_mubar	1	-14	20	-107.801	16.901	38.226	115.	
28 (Delta++)	11	2224	2	0.222	0.012-2734.287	2734		

397 pi+	1	211	209	0.006	0.398	-308.296	308.297	0.140
398 gamma	1	22	211	0.407	0.087	-1695.458	1695.458	0.000
399 gamma	1	22	211	0.113	-0.029	-314.822	314.822	0.000
400 (pi0)	11	111	212	0.021	0.122	-103.709	103.709	0.135
401 (pi0)	11	111	212	0.084	-0.068	-94.276	94.276	0.135
402 (pi0)	11	111	212	0.267	-0.052	-144.673	144.674	0.135
403 gamma	1	22	215	-1.581	2.473	3.306	4.421	0.000
404 gamma	1	22	215	-1.494	2.143	3.051	4.016	0.000
405 pi-	1	-211	216	0.007	0.738	4.015	4.085	0.140
406 pi+	1	211	216	-0.024	0.293	0.486	0.585	0.140
407 K+	1	321	218	4.382	-1.412	-1.799	4.968	0.494
408 pi-	1	-211	218	1.183	-0.894	-0.176	1.500	0.140
409 (pi0)	11	111	218	0.955	-0.459	-0.590	1.221	0.135
410 (pi0)	11	111	218	2.349	-1.105	-1.181	2.855	0.135
411 (Kbar0)	11	-311	219	1.441	-0.247	-0.472	1.615	0.498
412 pi-	1	-211	219	2.232	-0.400	-0.249	2.285	0.140
413 K+	1	321	220	1.380	-0.652	-0.361	1.644	0.494
414 (pi0)	11	111	220	1.078	-0.265	0.175	1.132	0.135
415 (K_S0)	11	310	222	1.841	0.111	0.894	2.108	0.498
416 K+	1	321	223	0.307	0.107	0.252	0.642	0.494
417 pi-	1	-211	223	0.266	0.316	-0.201	0.480	0.140
418 nbar0	1	-2112	226	1.335	1.641	2.078	3.111	0.940
419 (pi0)	11	111	226	0.899	1.046	1.311	1.908	0.135
420 pi+	1	211	227	0.217	1.407	1.356	1.971	0.140
421 (pi0)	11	111	227	1.207	2.336	2.767	3.820	0.135
422 n0	1	2112	228	3.475	5.324	5.702	8.592	0.940
423 pi-	1	-211	228	1.856	2.606	2.808	4.259	0.140
424 gamma	1	22	229	-0.012	0.247	0.421	0.489	0.000
425 gamma	1	22	229	0.025	0.034	0.009	0.043	0.000
426 pi+	1	211	230	2.718	5.229	6.403	8.703	0.140
427 (pi0)	11	111	230	4.109	6.747	7.597	10.961	0.135
428 pi-	1	-211	231	0.551	1.233	1.945	2.372	0.140
429 (pi0)	11	111	231	0.645	1.141	0.922	1.608	0.135
430 gamma	1	22	232	-0.383	1.169	1.208	1.724	0.000
431 gamma	1	22	232	-0.201	0.070	0.060	0.221	0.000

PYTHIA Monte Carlo
 $pp \rightarrow \text{gluino-gluino}$

Monte Carlo detector simulation

Takes as input the particle list and momenta from generator.

Simulates detector response:

- multiple Coulomb scattering (generate scattering angle),
- particle decays (generate lifetime),
- ionization energy loss (generate Δ),
- electromagnetic, hadronic showers,
- production of signals, electronics response, ...

Output = simulated raw data → input to reconstruction software:
track finding, fitting, etc.

Predict what you should see at ‘detector level’ given a certain hypothesis for ‘generator level’. Compare with the real data.

Estimate ‘efficiencies’ = #events found / # events generated.

Programming package: **GEANT**