


# Measurement of the Longitudinal Single Spin Asymmetries for W Boson Production in Polarized Proton-Proton Collisions at STAR

Amani Kraishan  
Temple University  
for the STAR Collaboration


DIS 2019  
Torino, Italy  
8-12 April 2019



DOE NP contract: DE-SC0013405



# Spin Puzzle



Jaffe and Monahar showed in 1990 that the proton spin can be written as a sum of contributions from quark and gluon spin and orbital angular momentum.

$$\langle S_p \rangle = \frac{1}{2} = \frac{1}{2} \boxed{\Delta \Sigma} + \boxed{\Delta G} + L_q + L_g$$

$$\Delta \Sigma = \int (\Delta u + \Delta d + \Delta s + \Delta \bar{u} + \Delta \bar{d} + \Delta \bar{s}) dx$$


$$\Delta G = \int \Delta g(x) dx$$

*$\Delta f(x)$  Helicity Distribution:* Probability density for finding a parton in a longitudinally polarized nucleon with flavor  $f$  and momentum fraction  $x$  in a nucleon.

# Helicity Distribution

- DSSV08 global analysis.
- The total contribution of up and down quarks spin has been well constrained.
- The flavor separated contributions of the sea quarks, still have quite large uncertainties.
- The gluon polarization also shows a large uncertainty band.

Daniel de Florian, Rodolfo Sassot, Marco Stratmann and Werner Vogelsang, Phys. Rev. D80 (2009) 034030.




# Weak Boson Production

The production of W boson in polarized proton-proton collisions is an independent method of probing the polarization of the light quarks and antiquarks.


- W production provides **direct sensitivity** to the **u and d quark** and **antiquark** helicity distributions.
- Large-scale defined by ***W* mass** ( $\sim 80$  GeV).
- Simple **final state** of charged lepton: no dependency on **fragmentation functions**.
- V-A coupling of the weak interaction leads to perfect spin separation.

# Single Spin Asymmetry of W




$$A_L^{e^-} \approx \frac{\int_{\otimes(x_1, x_2)} [\bar{u}(x_1)d(x_2)(1 - \cos\theta)^2 - \bar{d}(x_1)\bar{u}(x_2)(1 + \cos\theta)^2]}{\int_{\otimes(x_1, x_2)} [\bar{u}(x_1)d(x_2)(1 - \cos\theta)^2 + \bar{d}(x_1)\bar{u}(x_2)(1 + \cos\theta)^2]}$$

$$A_L^{e^+} \approx \frac{\int_{\otimes(x_1, x_2)} [\bar{d}(x_1)u(x_2)(1 + \cos\theta)^2 - \bar{u}(x_1)\bar{d}(x_2)(1 - \cos\theta)^2]}{\int_{\otimes(x_1, x_2)} [\bar{d}(x_1)u(x_2)(1 + \cos\theta)^2 + \bar{u}(x_1)\bar{d}(x_2)(1 - \cos\theta)^2]}$$



# Relativistic Heavy-Ion Collider (RHIC)

RHIC is the world's first polarized proton collider




# STAR Detector

$$\eta = -\ln \tan \frac{\theta}{2}$$

- Calorimetry system with  $2\pi$  coverage:
  - BEMC ( $-1 < \eta < 1$ )
  - EEMC ( $1 < \eta < 2$ )
- TPC: Tracking and particle ID
  - ( $|\eta| < 1.3$ )

Approximate kinematic range at RHIC  
 $-2 < \eta < 2$        $0.06 < x < 0.4$




# Run 2013 Dataset

Production runs at  $\sqrt{s}=500/510\text{GeV}$  (long. polarization) in 2009, 2011, 2012 and 2013:


- W production (Quark polarization)
- Jet and Hadron production (Gluon polarization)

Run	$L (\text{pb}^{-1})$	P (%)	FOM ( $P^2 L$ ) ( $\text{pb}^{-1}$ )
Run 9	12	0.38	1.7
Run 11	9.4	0.49	2.3
Run 12	72	0.56	24
Run 13	200	0.56	63




# W Selection at STAR

The W selection algorithm is built based on the topological and kinematic differences between W events and QCD events



$$p + p \rightarrow W \rightarrow e + \nu$$

- Isolated track pointing to isolated cluster in the calorimeter.
- Missing energy in the opposite azimuthal direction.




$$p + p \rightarrow QCD \rightarrow jets$$


- Several tracks pointing to several towers.
- Vector  $p_T$  sum is balanced by opposite jet.

# Jacobian Peak

- $E_T^{2x2}/E_T^{4x4} > 0.95$




- $E_T^{2x2}/E_T^{\Delta R<0.7} > 0.88$



- Signed  $p_T$  balance  $> 14 \text{ GeV}/c$
- Away  $E_T < 11 \text{ GeV}$

**Signal of Jacobian Peak  
with  $E_T$  distribution after  
selection cuts**



# Background Estimation

## Electroweak Background:

This background arises from well-understood electroweak processes:

- $Z \rightarrow e^- + e^+$
- $W \rightarrow \tau + \nu$



## QCD Background:

### Second EEMC:

Background (di-jets) which counts as a W event by escaping detection through non-existing calorimeter coverage ( $-2 < \eta < -1$ ).

### Data-driven QCD:


Background which passes  $e^\pm$  isolation cuts.



# WA<sub>L</sub> from Run 2011+2012



- A<sub>L</sub> of W<sup>-</sup> shows indication that data are larger than the DSSV predictions.
- A<sub>L</sub> of W<sup>+</sup> is consistent with theoretical predictions with DSSV pdf.
- Indication of symmetry breaking of polarized sea.

PRL 113 (2014) 072301




# Impact of STAR 2011+2012 W A<sub>L</sub> Measurements (NNPDF)

R. Ball et al. (NNPDF Collaboration), Nucl. Phys. B887 (2014) 276.



E. Nocera, PoS DIS2014 (2014) 204.




- The flavor asymmetry of polarized antiquarks in the nucleon is positive
- It has almost the same absolute size as the flavor asymmetry of unpolarized antiquarks



# W A<sub>L</sub> from Run 2013

- Most precise W A<sub>L</sub> results from 2013 dataset.
- Consistent with 2011+2012 published results, with **40% uncertainty reduced**.
- Further confirmed the polarized sea asymmetry.


Phys. Rev. D 99 (2019) 51102



# Impact of STAR 2013 W A<sub>L</sub> Measurements (NNPDF)



Phys. Rev. D 99 (2019) 51102




- The data confirm the existence of a sizable, positive  $u$ -bar polarization in the range  $0.05 < x < 0.2$  .
- The data confirm the existence of a flavor asymmetry in the polarized quark sea.

# W A<sub>LL</sub> from Run 2013

Double spin asymmetry of W can also provide access to  
u-bar, d-bar polarization

$$A_{LL}^{W^+} \propto \frac{\Delta u}{u} \frac{\Delta \bar{d}}{\bar{d}}$$

$$A_{LL}^{W^-} \propto \frac{\Delta d}{d} \frac{\Delta \bar{u}}{\bar{u}}$$



# Summary

- Sea quark polarization plays an important role in understanding the nucleon spin structure.
- STAR W A<sub>L</sub> place unique and significant constraints on the polarized quark and antiquark distributions.
- Significant shift of the central value of  $\Delta\bar{u}$  by including the new STAR 2013 W A<sub>L</sub> results.
- First clear evidence of the flavor-asymmetry in the polarized quark sea.

**Phys. Rev. D 99 (2019) 51102**

