

Azimuthally Sensitive Buda-Lund Hydrodynamic Model and Fits to Spectra, Elliptic Flow and asHBT

András Ster¹, M.Csanád², T.Csörgő¹, B. Tomasik³

¹*MTA KFKI RMKI, Budapest, Hungary,*

²*ELTE University, Budapest, Hungary,*

³*Univ. Mat. Bela, Banská Bystrica, Slovakia*

- **Buda-Lund hydro model**
- **Observables**
- **Review of old central hydro results**
- **New non-central hydro results**
- **Conclusion**

The Buda-Lund hydro model

- General form of a particle emission function

$$S(x, p) d^4x = f(x, p) \ p_\mu \ d\sigma^\mu(x)$$

- With (Boltzmann) probability distribution for fluids

$$S(x, p) d^4x = \frac{g}{(2\pi)^3} \frac{p_\mu d\sigma^\mu(x)}{\exp\left(\frac{p_\mu u^\mu(x)}{T(x)} - \frac{\mu(x)}{T(x)}\right) + S_q}$$

The Buda-Lund hydro model

$\sigma(x)$	hypersurface
$u(x)$	flow field
$\mu(x)$	chemical
$T(x)$	potential temperature

must comply with the 5 differential equations of fluid dynamics:

- continuity
- momentum conservation
- energy conservation

The Buda-Lund hydro model

- Equations of non-relativistic inviscid fluid

$$\begin{aligned}\partial_t n + \nabla \cdot (n\mathbf{v}) &= 0, \\ \partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} &= -(\nabla p)/(mn), \\ \partial_t \epsilon + \nabla \cdot (\epsilon \mathbf{v}) &= -p \nabla \cdot \mathbf{v},\end{aligned}$$

- Not closed, EoS needed, for example

$$p = nT$$

$$\epsilon = \kappa p$$

The Buda-Lund hydro model

5 model principles

- 3D expansion with axial or ellipsoidal symmetry
- Local thermal equilibrium
- Analytic expressions for the observables
- Reproduction known exact hydro solutions (nonrelativistic, hubble and bjorken limits)
- Core-halo picture (long lived resonances)

M. Csanad, T.Cs rg , B. L rstad: Nucl.Phys.A742:80-94,2004; nucl-th/0310040

The Buda-Lund Hydro Model

Buda-Lund form of hydro fields:

in several cases parametric solutions of hydrodynamics

see M. Csanad's talk

$$d^4\sigma(x) = u(x)H(\tau)d^4x$$

$$u(x) = (\gamma, \sinh\eta_x, \sinh\eta_y, \sinh\eta_z)$$

$$\frac{\mu(x)}{T(x)} = \frac{\mu_0}{T_0} - s$$

$$\frac{1}{T(x)} = \frac{1}{T_0} \left(1 + \frac{T_0 - T_s}{T_s} s \right) \left(1 + \frac{(T_0 - T_e)(\tau - \tau_0)^2}{T_e 2\Delta\tau^2} \right)$$

The Buda-Lund Hydro Model (2)

- Where (in case of axial symmetry):

$$H(\tau) = \frac{1}{(2\pi\Delta\tau^2)^{1/2}} \exp\left(-\frac{(\tau - \tau_0)^2}{2\Delta\tau^2}\right)$$

$$S = \frac{r_t^2}{(2R_G^2)} + \frac{(\eta - y_0)^2}{2\Delta\eta^2}$$

$$\sinh(\eta_t) = \frac{\langle u_t \rangle r_t}{R_G} = H_t r_t$$

H_t: transverse Hubble constant

The Buda-Lund hydro model (3)

- **Observables**

$$\mathbf{N}_1(p) = \int d^4x S(x, p)$$

$$C_2(Q, p) = 1 + \lambda_* \exp(-Q_o^2 R_o^2 - Q_s^2 R_s^2 - Q_l^2 R_l^2)$$

$$S(x, p) = S_c(x, p) + S_h(x, p)$$

$$\mathbf{N}_1(p) = \frac{1}{\sqrt{\lambda_*}} \int d^4x S_c(x, p)$$

Buda-Lund Hydro: Observables

- Final form of the Invariant Momentum Distribution:

$$N(\mathbf{p}) = \frac{g}{(2\pi)^3} \overline{EV} \overline{C} \exp\left(-\frac{\mathbf{p} \cdot \mathbf{u}(\bar{x}) - \mu(\bar{x})}{T(\bar{x})} + s_q\right)$$

$$\overline{E} = m_t \cosh(\bar{\eta})$$

$$\overline{V} = 2\pi^{(3/2)} \overline{R}_{par} \overline{R}_t^2 \frac{\overline{\Delta\tau}}{\Delta\tau}$$

$$\overline{C} = \frac{1}{\sqrt{\lambda_*}} \exp\left(\frac{\overline{\Delta\eta}^2}{2}\right)$$

Buda-Lund Hydro Model: HBT radii

- Final form of the HBT radii:

$$\bar{R}_t^2 = \bar{R}_G^2 / [1 + (\langle u_t \rangle^2 + (T_0 - T_s)/T_s) \bar{E}/T_0]$$

$$\bar{R}_{par}^2 = \tau_0^2 / \Delta\bar{\eta}^2$$

$$\Delta\bar{\eta}^2 = \Delta\eta^2 / (1 + \Delta\eta^2 \bar{E}/T_0)$$



$$\Delta\bar{\tau}^2 = \Delta\tau^2 / [1 + ((T_0 - T_e)/T_e) \bar{E}/T_0]$$

$$\bar{\tau} = \tau_0, \quad \bar{r}_x = u_t R_G p_t / [T_0 + \bar{E}(u_t)(T_0 - T_s)/T_s]$$

$$\bar{r}_y = 0$$



$$\bar{\eta} = (y_0 - y) / [1 + \Delta\eta^2 m_t / T_0]$$

Buda-Lund fits to NA22 h+p





N. M. Agababyan et al, EHS/NA22 , PLB 422 (1998) 395
 T. Csörgő, hep-ph/001233, Heavy Ion Phys. 15 (2002) 1-80

Emission function from NA22 h+p





BudaLund fits to SPS Pb+Pb





A. Ster, T.Csörgő, B. Lörstad , Nucl.Phys. A661 (1999) 419-422, nucl-th/9907338

Emission function from SPS Pb+Pb





BudaLund fits to RHIC Au+Au



BudaLund hydro fits to 130 AGeV Au+Au



BudaLund hydro fits to 130 AGeV Au+Au




Emission function from RHIC Au+Au




BudaLund fits to RHIC Au+Au



BudaLund v1.5 fits to 200 AGeV Au+Au



BudaLund v1.5 fits to 200 AGeV Au+Au





<http://www.kfki.hu/~csorgo/budalund/budalund1.5.qm04.tar.gz>



M. Csand, et al., J.Phys.G30: S1079-S1082, 2004, nucl-th/0403074

Emission function from RHIC Au+Au



Buda-Lund fit results



BL v1.5 parameters	RHIC 200 GeV Au+Au	RHIC 130 GeV Au+Au	Pb+Pb SPS	h+p SPS
T_0 [MeV]	196 ± 13	214 ± 7	139 ± 6	140 ± 3
$\langle u_t \rangle$	1.6 ± 0.2	1.0 ± 0.1	0.55 ± 0.06	0.20 ± 0.07
R_G [fm]	13.5 ± 1.7	28.0 ± 5.5	7.1 ± 0.2	0.88 ± 0.13
R_s [fm]	12.4 ± 1.6	8.6 ± 0.4	28 ± 21	1.4 ± 0.3
T_{surf} [MeV]	0.5 T_0	0.5 T_0	0.5T_0	0.5T_0
τ_0 [fm/c]	5.8 ± 0.3	6.0 ± 0.2	5.9 ± 0.6	1.4 ± 0.1
$\Delta\tau$ [fm/c]	0.9 ± 1.2	0.3 ± 1.2	1.6 ± 1.5	1.3 ± 0.3
$\Delta\eta$	3.1 ± 0.1	2.4 ± 0.1	2.1 ± 0.4	1.36 ± 0.02
T_{evap} [MeV]	117 ± 12	102 ± 11	87 ± 24	-
μ_0^π [MeV]	-2 ± 14	63 ± 11		
μ_0^K [MeV]	16 ± 19	98 ± 19		
$\mu_0^{P^-}$ [MeV]	97 ± 28	315 ± 27		
μ_B [MeV]	61 ± 52	77 ± 38	0 fixed	0 fixed
χ^2/NDF	114/208=0.55	158/180=0.9	342/277=1.2	642/683=0.9
CL	100 %	88 %		

$$\beta_t = \frac{\langle u_t \rangle}{\sqrt{1 + \langle u_t \rangle^2}}$$



BudaLund fits to RHIC preliminary

p+p

BudaLund v1.5 fits - p+p data at 200 GeV





BudaLund v1.5 fits - p+p prel. at RHIC



T. Csörgő, et al., Heavy Ion Physics, hep-ph/0406042

Emission function from RHIC p+p





BudaLund fit results



BL v1.5 parameters	RHIC 200 GeV (prel) p+p	RHIC 130 GeV Au+Au	Pb+Pb SPS	h+p SPS
T_0 [MeV]	289 ± 8	214 ± 7	139 ± 6	140 ± 3
$\langle u_t \rangle$	0.04 ± 0.26	1.0 ± 0.1	0.55 ± 0.06	0.20 ± 0.07
R_G [fm]	1.2 ± 0.3	28.0 ± 5.5	7.1 ± 0.2	0.88 ± 0.13
R_s [fm]	1.13 ± 0.16	8.6 ± 0.4	28 ± 21	1.4 ± 0.3
T_{surf} [MeV]	0.5 T_0	0.5 T_0	0.5T_0	0.5T_0
τ_0 [fm/c]	1.1 ± 0.2	6.0 ± 0.2	5.9 ± 0.6	1.4 ± 0.1
$\Delta\tau$ [fm/c]	0.1 ± 0.5	0.3 ± 1.2	1.6 ± 1.5	1.3 ± 0.3
$\Delta\eta$	3.0 fixed	2.3 ± 0.4	2.1 ± 0.4	1.36 ± 0.02
T_{evap} [MeV]	90 ± 42	102 ± 11	87 ± 24	-
χ^2/NDF	89/71	158/180	342/277	642/683

BudaLund fits to NA49 data (preliminary HBT, work in progress)


BudaLund hydro fits to 158 AGeV Pb+Pb



BudaLund hydro fits to 158 AGeV Pb+Pb



BudaLund fits to NA49 data (preliminary HBT, work in progress)




BudaLund fit results of NA49 data

BudaLund parameters	158 AGeV	80 AGeV	40 AGeV	30 AGeV	20 AGeV
T_0 [MeV]	143 \pm 7	109 \pm 7	106 \pm 4	107 \pm 7	86 \pm 3
T_e [MeV]	141 \pm 5	105 \pm 3	103 \pm 2	101 \pm 3	86 \pm 2
H_t [c/fm]	0.099 \pm 0.003	0.104 \pm 0.004	0.111 \pm 0.003	0.110 \pm 0.005	0.106 \pm 0.004
H_i [c/fm]	0.155 \pm 0.005	0.129 \pm 0.005	0.123 \pm 0.003	0.120 \pm 0.004	0.107 \pm 0.002
R_G [fm]	7.1 \pm 0.2	7.9 \pm 0.4	9.8 \pm 0.7	9.8 \pm 0.9	12.1 \pm 1.7
R_s [fm]	18.7 \pm 2.0	24.5 \pm 3.8	21.5 \pm 1.8	20.7 \pm 3.0	25.5 \pm 2.7
$\Delta\tau$ [fm/c]	3.3 \pm 0.3	4.1 \pm 0.4	3.9 \pm 0.3	3.7 \pm 0.3	3.2 \pm 0.3
$\Delta\eta$	1.5 \pm 0.2	1.4 \pm 0.2	1.2 \pm 0.1	1.1 \pm 0.1	1.0 \pm 0.1
μ_0^π [MeV]	88 \pm 7	109 \pm 10	79 \pm 6	77 \pm 9	88 \pm 6
μ_0^K [MeV]	61 \pm 16	151 \pm 21	122 \pm 12	128 \pm 20	163 \pm 12
$\mu_0^{K^+}$ [MeV]	137 \pm 16	236 \pm 21	224 \pm 12	230 \pm 21	257 \pm 12
μ_0^P [MeV]	403 \pm 29	593 \pm 38	635 \pm 22	642 \pm 38	700 \pm 21
χ^2/NDF	198 /126 !	129 /128	200 /116 !	160 /116	128 /95

$$\langle u_t \rangle = H_t \cdot R_G$$

$$\tau_0 = 1/H_I$$

source parameter excitation functions



Some other hydro models

Models with acceptable results:

[nucl-th/0207016](#)

Buda-Lund / core-halo model.

T.Csörgő, A. Ster, Heavy Ion Phys. 17
(2003) 295-312.

[nucl-th/0204054](#)

Multiphase Transport model (AMPT).

Z. Lin, C. M. Ko, S. Pal.

[nucl-ex/0307026](#)

Blast wave model. F. Retière for STAR.

[nucl-th/0205053](#)

Hadron cascade model. T. Humanic.

Other interesting models:

[nucl-th/0208068](#)

3D hydro model. T. Hirano, & T.Tsuda.

[hep-ph/0209054](#)

Hadron model. W.Broniowski, A. Baran


W. Florkowski.

Femtoscopy of QGP

BudaLund fits indicate:

scaling of HBT radii

BudaLund prediction
(1995):
each $R^2 \sim 1 / m_t$



Conclusions on central collisions

- BudaLund model describes single particle distributions, rapidity distributions, HBT correlation function radii w/o puzzle in the following reactions:

$h+p$ @SPS, $p+p$ @RHIC, $Pb+Pb$ @SPS,
 $Au+Au$ @RHIC

- Rings of fire in $h+p$ @SPS and $p+p$ @RHIC
Fireballs in $Pb+Pb$ @SPS and $Au+Au$ @RHIC
- $T < T_c$ in $h+p$ and $Pb+Pb$ @SPS
 $T > T_c$ in $p+p$ and $Au+Au$ @RHIC; T_c

asBuda-Lund fits to non-central RHIC data

**Model extensions to ellipsoidal symmetry
for elliptic flow:**

calculate 2nd harmonic coefficient of anisotropy:

$$\frac{N(p)}{d\Phi} \propto 1 + 2v_2(p) \cos(2\Phi)$$

BudaLund fits to non-central RHIC data

Exact non-relativistic result for elliptic flow:

$$v_2 = \frac{I_1(w)}{I_0(w)} +$$

I_n : modified Bessel
functions

$$w = \frac{p_t^2}{4\bar{m}_t} \left(\frac{1}{T_{*,y}} - \frac{1}{T_{*,x}} \right)$$

$$\bar{m}_t = m_t \cosh(\eta_s - y)$$

$$T_{*,i} = T_0 + m_t \dot{X}_i^2 \frac{T_0}{T_0 + m_t a^2}$$

Effective temperatures in
reaction plane and in perp.

$$a^2 = \frac{T_0 - T_s}{T_s} = \left\langle \frac{\Delta T}{T} \right\rangle_r$$

For detailed calculations see: M. Csanad et al., hep-ph/0801.4434v2

BudaLund fits to non-central RHIC data

Model extensions to ellipsoidal symmetry

for asHBT radii:

$$R_o^2 = R_{*,x}^2 \cos^2 \varphi + R_{*,y}^2 \sin^2 \varphi + \beta_o^2 \Delta\tau_*^2 \quad (36a)$$

$$= \frac{R_{*,x}^2 + R_{*,y}^2}{2} + \beta_o^2 \Delta\tau_*^2 - \frac{R_{*,y}^2 - R_{*,x}^2}{2} \cos(2\varphi)$$

$$R_s^2 = R_{*,x}^2 \sin^2 \varphi + R_{*,y}^2 \cos^2 \varphi \quad (36b)$$

$$= \frac{R_{*,x}^2 + R_{*,y}^2}{2} + \frac{R_{*,y}^2 - R_{*,x}^2}{2} \cos(2\varphi),$$

$$R_{os}^2 = \frac{R_{*,y}^2 - R_{*,x}^2}{2} \sin(2\varphi), \quad (36c)$$

$$R_l^2 = R_{*,z}^2, \quad (36d)$$

X=R_X, Y=R_Y

We found that „a” is different in each direction

$$R_{*,x}^2 = X^2 \left(1 + \frac{m_t (a^2 + \dot{X}^2)}{T_0} \right)^{-1},$$

$$R_{*,y}^2 = Y^2 \left(1 + \frac{m_t (a^2 + \dot{Y}^2)}{T_0} \right)^{-1},$$

$$R_{*,z}^2 = Z^2 \left(1 + \frac{m_t (a^2 + \dot{Z}^2)}{T_0} \right)^{-1}.$$

asBuda-Lund fits to non-central RHIC data

**Model extensions to ellipsoidal symmetry
for azimuthally integrated spectra:**

Calculate the volume term for ellipsoids :


$$\bar{V} = 2\pi^{(3/2)} \bar{R}_{par} \bar{R}_t^2 \frac{\overline{\Delta\tau}}{\Delta\tau}$$




replaced \bar{R}_t^2 by $\bar{R}_x * \bar{R}_y$

asBudaLund fits to non-central RHIC data

BudaLund hydro fits to 200 GeV Au+Au



BudaLund hydro fits to 200 GeV Au+Au




asBuda-Lund fit results of non-central RHIC data

BudaLund source	RHIC 200 AGeV central (0-30%)	RHIC 200 AGeV non-central (20-30%)
parameters		
T_0 [MeV]	196 ± 13	179 ± 7
T_e [MeV]	117 ± 12	119 ± 7
H_x [c/fm]	0.118 ± 0.013	0.150 ± 0.002
H_y [c/fm]		0.111 ± 0.001
H_z [c/fm]	0.172 ± 0.008	0.187 ± 0.005
R_x [fm]	13.5 ± 1.7	7.8 ± 0.3
R_y [fm]		7.2 ± 0.2
R_{xS} [fm]	12.4 ± 1.6	12.2 ± 0.9
R_{yS} [fm]		12.5 ± 0.8
Δr [fm/c]	0.9 ± 1.2	2.7 ± 0.2
$\Delta\eta$	3.1 ± 0.1	2.5 ± 0.3
μ_0^π [MeV]	-2 ± 14	40 ± 8
μ_0^K [MeV]	16 ± 19	55 ± 13
μ_0^P [MeV]	97 ± 28	178 ± 22
χ^2/NDF	$114 / 208$	$261 / 152$
	(CL w/o radii ~10%)	


BudaLund geometrical picture of non-central RHIC reactions

Schematic view

20-30% $\sim b = 8-12$ fm



Universal v_2 scaling predicted by BudaLund model (2003)



Conclusion

- **asBuda-Lund model describes single particle distributions, elliptic flow and asHBT correlation function radii of 20-30% centrality data of 200 GeV Au+Au reactions at RHIC.**
- We find that T drops to 179 MeV from ~200 MeV of central collisions.
- The source at freeze-out was found to be slightly extended in the reaction plane ($R_x > R_y$), in agreement with M. Csanad's conclusion