First Result for a Full Two-Loop Five-Gluon Amplitude

Based on arXiv:1905.03733

Simon Badger
Durham University

Towards an Era of Precision Measurements at the LHC

A new era of precision measurements is dawning. CERN's Large Hadron Collider (LHC) will start Run 3 in 2021. To fully exploit the machine's potential, accurate theoretical predictions are required. These can be obtained through the computation of higher orders in perturbation theory.

All Feynman Integrals Computed Analytically in the Physical Region

Basis for all QCD amplitudes → pentagon functions

(*https://pentagonfunctions.hepforge.org/*)

Logarithms, dilogarithms, and generalizations thereof (Goncharov polylogarithms)

- Elegant analytic properties
- Fast and accurate numerical evaluation

Dmitry Chicherin
Max Planck Institute for Physics

Thomas Gehrmann
University of Zurich

For Many QCD Processes, Next-To-Leading-Order is Insufficient

E.g. strong coupling from 3-jet/2-jet ratio [1]:

$$\alpha_S(M_Z) = 0.1148 \pm 0.0014_{\text{(exp.)}} \pm 0.0018_{\text{(PDF)}} \pm 0.0050_{\text{(theory)}}$$

Large theoretical uncertainty!

At present, only observables involving up to four particles are available at next-to-next-to-leading order.

A Refined Amplitude Assembly



The code can be parallelized, and runs in a few hours using multi-threading on a modern computing node.

Gudrun Heinrich
Max Planck Institute for Physics

Johannes Henn
Max Planck Institute for Physics

Five-Particle Processes at the LHC

- Precision measurements of the strong coupling
- In-depth tests of the Standard Model
- Improved background to new physics searches

Bottleneck: two-loop five-particle amplitudes



Planar amplitudes [2,3]

Cutting-Edge Methods

- Generalized unitarity for Feynman integrands [4]
- Differential equations for Feynman integrals in the canonical form [2,5,6]
- Algorithmic construction of canonical basis using D-dimensional leading singularities [6]
- Computational algebraic geometry [7]
- Finite fields and functional reconstruction [8]

Tiziano Peraro
University of Zurich

Pascal Wasser
Johannes Gutenberg University Mainz

New Result: Full-Color All-Plus Helicity Amplitude

We find a remarkably simple expression for the infrared subtracted hard function:

Permutations of the external legs · Color SU(N_c) · Gluon spin dimension · Spinor-helicity variables

$$\mathcal{H}^{(2)}_{\substack{\text{double}\\\text{trace}}} = \sum_{S_5/\Sigma} \text{Tr}(12)[\text{Tr}(345) - \text{Tr}(543)] \sum_{\Sigma} \left\{ 6\,\kappa^2 \left[\frac{\langle 24\rangle[14][23]}{\langle 12\rangle\langle 23\rangle\langle 45\rangle^2} + 9\,\frac{\langle 24\rangle[12][23]}{\langle 12\rangle\langle 34\rangle\langle 45\rangle^2} \right] \right.$$

$$\left. + \kappa\,\frac{[15]^2}{\langle 23\rangle\langle 34\rangle\langle 42\rangle} \left[\text{box}^{4\,1\,5\,3}_{3\quad\;\;2\,4} + \text{box}^{1\;\;5\,4}_{\;\;2\;\;2} - 4\,\text{box}^{1\;\;5\,5}_{\;\;3\;\;4} - 4\,\text{box}^{1\;\;2\,4}_{\;\;3\;\;5} - 4\,\text{box}^{1\;\;2\,5}_{\;\;3\;\;3} \right] \right\}$$

Only logarithms and dilogarithms appear:

$$\text{box}^{3\;4\,5}_{2\quad\;1} = \text{Li}_2\left(1 - \frac{s_{12}}{s_{45}}\right) + \text{Li}_2\left(1 - \frac{s_{23}}{s_{45}}\right) + \log^2\left(\frac{s_{12}}{s_{23}}\right) + \frac{\pi^2}{6}$$

Valid in all physical scattering regions $s_{ij} \to s_{ij} + i0$

Surprise: coefficients of box functions are conformally invariant!



Yang Zhang
University of Science and Technology of China

Simone Zoia
Max Planck Institute for Physics

References and Acknowledgments

[1] CMS Collaboration, Eur.Phys.J. C73 (2013) no.10, 2604
[2] Gehrmann, Henn, Lo Presti, Phys.Rev.Lett. 116 (2016) no.6, 062001
[3] Badger, Brønnum-Hansen, Hartanto, Peraro, Phys.Rev.Lett. 120 (2018) no.9, 092001; Abreu, Febres Cordero, Ita, Page, Zeng, Phys.Rev. D97 (2018) no.11, 116014; Abreu, Febres Cordero, Ita, Page, Sotnikov, JHEP 1811 (2018) 116; Badger, Brønnum-Hansen, Hartanto, Peraro, JHEP 1901 (2019) 186; Abreu, Dormans, Febres Cordero, Ita, and Page, Phys.Rev.Lett. 122 (2019) no.8, 082002; Abreu, Dormans, Febres-Cordero, Ita, Page, Sotnikov, HEP 1905 (2019) 084;
[4] Badger, Mogull, Ochirov, O'Connell, JHEP 1608 (2016) 063
[5] Papadopoulos, Tommasini, Wever, JHEP 1604 (2016) 078; Gehrmann, Henn, Lo Presti, JHEP 1810 (2018) 103; Abreu, Page, Zeng, JHEP 1901 (2019) 006; Chicherin, Gehrmann, Henn, Lo Presti, Mitev, Wasser, JHEP 1903 (2019) 042; Abreu, Dixon, Herrmann, Page, Zeng, Phys.Rev.Lett. 122 (2019) no.12, 121603
[6] Chicherin, Gehrmann, Henn, Wasser, Zhang, Zoia, arXiv:1812.11160
[7] Zhang, JHEP 1209 (2012) 042; Mastrolia, Mirabella, Ossola, Peraro, Phys.Lett. B718 (2012) 173; Böhm, Georgoudis, Larsen, Schönemann, Zhang, JHEP 1809 (2018) 024
[8] von Manteuffel, Schabinger, Phys.Lett. B744 (2015) 101; Peraro, JHEP 1612 (2016) 030; Peraro, arXiv:1905.08019

This research received funding from the Swiss National Science Foundation (Ambizione grant PZ00P2 161341), the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme, Novel structures in scattering amplitudes (grant agreement No 725110), and High precision multi-jet dynamics at the LHC (grant agreement No 772009), under the Marie Skłodowska-Curie grant agreement 746223, from the COST Action CA16201 Particleface, and from the STFC Rutherford Fellowship ST/L004925/1.

European Research Council
Established by the European Commission

Do you want to learn more?

Listen to our podcast!