

# 10th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions



Contribution ID: 289

Type: Oral Presentation

## New PHENIX Results on Mid-Rapidity Bottom and Charm Production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV

Monday 1 June 2020 13:35 (20 minutes)

Energy loss of quarks in the hot and dense medium has been studied for decades. Both the experimental and theoretical efforts hinted that the energy loss is quark mass dependent (the yield of heavier quarks will be less suppressed). It was found that the electrons from heavy quarks (charm, and bottom) are less or similarly suppressed compared to that of light hadrons. The mass ordering of the suppression between charm and bottom was not clear by now due to large experimental uncertainties. We have fully exploited the events recorded at PHENIX from Au+Au collisions from RHIC Year-2014 run, and with the new charm and bottom  $p_T$  spectra from p+p collisions from Year-2015 run, we can obtain the new  $R_{AA}$  for charm and bottom quarks with smaller uncertainties. We will show the latest results on the  $R_{AA}$  of electrons and unfolded hadrons from charm and bottom quarks separately, and implications on the understanding of the quark mass dependence of the energy loss.

### Collaboration (if applicable)

PHENIX

### Track

Heavy Flavor and Quarkonia

### Contribution type

Contributed Talk

**Author:** DAVID, Gabor (Brookhaven National Laboratory)

**Presenter:** Dr KHATIWADA, Ajeeta (Purdue University (US))

**Session Classification:** Parallel

**Track Classification:** Heavy Flavor and Quarkonia