

10th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions



Contribution ID: 283

Type: Poster

Light-by-light scattering in ultra-peripheral Pb+Pb collisions in the ATLAS experiment

Tuesday, June 2, 2020 7:30 AM (1h 20m)

The ultra-peripheral Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV performed at the ATLAS experiment are used to study a rare light-by-light scattering process,

$\gamma\gamma \rightarrow \gamma\gamma$, allowed in Quantum Electrodynamics via a loop

diagram. The poster summarises recent light-by-light measurements conducted using a combination of 2015 and 2018 datasets recorded by the ATLAS experiment, corresponding to an integrated luminosity of 2.2 nb^{-1} . The light-by-light event candidates are required to consist of only two photons produced exclusively, each with transverse energy $E_T > 2.5$ GeV, pseudorapidity $|\eta| < 2.4$, diphoton invariant mass $m_{\gamma\gamma} > 5$ GeV, and with diphoton transverse momentum $p_{T\gamma\gamma} < 1$ GeV and acoplanarity below 0.01. The differential distributions, presented as functions of kinematic and angular variables of the final-state photons, are unfolded for detector effects. The fiducial and differential cross-sections are presented and compared with theoretical predictions. The diphoton invariant mass distribution is used to set limits on the production of axion-like particles.

Collaboration (if applicable)

ATLAS

Track

Initial State

Contribution type

Poster

Primary authors: COLLABORATION, ATLAS; ZIVKOVIC, Lidija (Institute of physics Belgrade (RS))

Presenter: MAJ, Klaudia (AGH University of Science and Technology (PL))

Session Classification: Poster session

Track Classification: Initial State