



Contribution ID: 50

Type: Poster

## Measurement of the azimuthal angle dependence of jet yields in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02\sim\text{TeV}$ with the ATLAS detector

It has been shown that high energy partons lose energy when traversing the hot, dense medium produced in heavy-ion collisions. However, the mechanism of the energy loss, including its dependence on the path-length of the shower in the medium, is not fully understood. This poster presents a measurement of single jet yields as a function of the azimuthal angle with respect to the event plane in Pb+Pb collisions at  $\sqrt{s_{NN}} = 5.02\sim\text{TeV}$ . Because jets at different angles with respect to the event plane traverse, on average, different path lengths of the medium, this measurement gives insight into the path-length dependence of parton energy loss. The azimuthal angle dependence of the yields is characterized by the parameter  $v_n^{jet}$ , which quantifies the magnitude of the modulation of the azimuthal angle distribution with respect to the  $n^{th}$  order event plane. While ATLAS has previously reported the  $v_2^{jet}$  in Pb+Pb at  $\sqrt{s_{NN}} = 2.76\sim\text{TeV}$ , this is the first ATLAS measurement of higher order  $v_n^{jet}$ .

### Collaboration (if applicable)

ATLAS

### Track

Jets and High Momentum Hadrons

### Contribution type

Poster

**Primary authors:** COLLABORATION, ATLAS; ZIVKOVIC, Lidija (Institute of physics Belgrade (RS))

**Presenter:** COLLABORATION, ATLAS

**Track Classification:** Jets and High Momentum Hadrons