

EW Baryogenesis through varying Yukawas in Composite Higgs models

Sebastian Bruggisser

In collaboration with:
T. Konstandin & O. Matsedonskyi
& G. Servant & B. von Harling

arXiv: 1706.08534,
1803.08546, 1804.07314



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386

Conditions for Baryogenesis

Sakharov '67

Conditions for Baryogenesis


Sakharov '67

- Baryon number violation

Conditions for Baryogenesis

Sakharov '67

- Baryon number violation
 - Electroweak Sphaleron



Conditions for Baryogenesis

Sakharov '67

- Baryon number violation
 - Electroweak Sphaleron
- CP violation

Conditions for Baryogenesis

Sakharov '67

- Baryon number violation

→ Electroweak Sphaleron


- CP violation

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

Farrar, Shaposhnikov '93

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_{\substack{i>j \\ d,s,b}} (m_i^2 - m_j^2) J_{CP}$$

Gavela, et al. '93
Huet, Sather '94

Jarlskog constant 

Conditions for Baryogenesis

Sakharov '67

- Baryon number violation

→ Electroweak Sphaleron

- CP violation

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_u (m_u^2 - M_W^2)$$

Jarlskog constant

Much too small!

Nikov '93

Gavela, et al. '93
Huet, Sather '94

Conditions for Baryogenesis

Sakharov '67

- Baryon number violation

→ Electroweak Sphaleron

- CP violation

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_u \sin \theta_u$$

Jarlskog constant

Much too small!

Nikov '93

Gavela, et al. '93
Huet, Sather '94

CP violation in the top sector:

- 2 Higgs doublet
- Composite Higgs

Cohen, Kaplan, Nelson '94

Espinosa, Gripaios, Konstandin, Riva '11

Conditions for Baryogenesis

Sakharov '67

- Baryon number violation

→ Electroweak Sphaleron

- CP violation

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_u (m_u^2 - m_t^2)$$

Jarlskog constant

Much too small!

Nikov '93

Gavela, et al. '93
Huet, Sather '94

CP violation in the top sector:

- 2 Higgs doublet
- Composite Higgs

Cohen, Kaplan, Nelson '94

Espinosa, Gripaios, Konstandin, Riva '11

Supersymmetric CP violation:

- Chargino/Neutralino sector
- Sfermion sector

Carena et al. '97, '00, '02

Konstandin, Prokopek, Schmidt, Seco '05
Cirigliano, Li, Profumo, Ramsey-Musolf '09
Chung, Garbrecht, Ramsey-Musolf '09
Kozaczuk, Profumo, Ramsey-Musolf '12

Conditions for Baryogenesis

Sakharov '67

- Baryon number violation
 - Electroweak Sphaleron
- CP violation
 - Dynamical CP violation from varying Yukawas

Conditions for Baryogenesis

Sakharov '67

- Baryon number violation
 - Electroweak Sphaleron
- CP violation
 - Dynamical CP violation from varying Yukawas
- Out-of-equilibrium

Conditions for Baryogenesis


Sakharov '67

- Baryon number violation
 - Electroweak Sphaleron
- CP violation
 - Dynamical CP violation from varying Yukawas
- Out-of-equilibrium
 - Strong first-order electroweak phase transition

Electroweak baryogenesis

$\langle h \rangle \neq 0$

$\langle h \rangle = 0$




Electroweak baryogenesis

$\langle h \rangle \neq 0$

$\langle h \rangle = 0$

$$\Gamma_{ws} = 10^{-6} T \exp(-ah(z)/T)$$




Electroweak baryogenesis

$\langle h \rangle \neq 0$

Sphaleron suppressed


Sphaleron active $\langle h \rangle = 0$

$$\Gamma_{ws} = 10^{-6} T \exp(-ah(z)/T)$$




z


Electroweak baryogenesis




Electroweak baryogenesis



Electroweak baryogenesis



Electroweak baryogenesis




Electroweak baryogenesis

$$\langle h \rangle \neq 0$$

Sphaleron suppressed


Sphaleron active $\langle h \rangle = 0$



$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)'' m V \right]_{ii}$$

where V s.t. $V^\dagger m^\dagger m V$ is diagonal

Electroweak baryogenesis




$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)'' m V \right]_{ii}$$

where V s.t. $V^\dagger m^\dagger m V$ is diagonal

- Zero for constant Yukawas

Electroweak baryogenesis




$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)^{''} m V \right]_{ii}$$

where V s.t. $V^\dagger m^\dagger m V$ is diagonal

- Zero for constant Yukawas
- Only active during the variation of Yukawas

Electroweak baryogenesis



$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)^{''} m V \right]_{ii}$$

where V s.t. $V^\dagger m^\dagger m V$ is diagonal

- Zero for constant Yukawas
- Only active during the variation of Yukawas
- Connection between flavour and Higgs physics: Froggat-Nielsen, Randall-Sundrum, Composite Higgs, ...

Composite Higgs

Composite Higgs

- Higgs: composite object of new nearly conformal confining sector $H \sim \bar{\psi}\psi$

Composite Higgs

- Higgs: composite object of new nearly conformal confining sector $H \sim \bar{\psi}\psi$
- Masses of remaining resonances: $m_\star \sim g_\star f$

Composite Higgs

- Higgs: composite object of new nearly conformal confining sector $H \sim \bar{\psi}\psi$
 - Masses of remaining resonances: $m_\star \sim g_\star f$
- 
- Confinement scale

Composite Higgs


- Higgs: composite object of new nearly conformal confining sector $H \sim \bar{\psi}\psi$
- Masses of remaining resonances: $m_\star \sim g_\star f$  typical strong coupling $\propto N^{-1/2}$

Confinement scale

Composite Higgs

- Higgs: composite object of new nearly conformal confining sector $H \sim \bar{\psi}\psi$
- Masses of remaining resonances: $m_\star \sim g_\star f$  typical strong coupling $\propto N^{-1/2}$ Confinement scale
- Confinement scale set by the vev of the dilaton $\langle\chi\rangle \sim f$ breaking conformal invariance

Composite Higgs

- Higgs: composite object of new nearly conformal confining sector $H \sim \bar{\psi}\psi$
- Masses of remaining resonances: $m_\star \sim g_\star f$  Confinement scale
typical strong coupling $\propto N^{-1/2}$
- Confinement scale set by the vev of the dilaton $\langle\chi\rangle \sim f$ breaking conformal invariance
- Mass gap: Higgs is a PNGB of symmetry breaking \mathcal{G}/\mathcal{H}

Composite Higgs


Composite Higgs

- What we need:
 - Variation of Yukawa couplings for CP-violation
 - Strong first-order electroweak phase transition


Composite Higgs

- What we need:
 - Variation of Yukawa couplings for CP-violation
 - Strong first-order electroweak phase transition
- What CH naturally provides:
 - Varying Yukawas through partial compositeness
 - Joint confinement-electroweak phase transition


Studying the phase transition in the Dilaton-Higgs potential




Studying the phase transition in the Dilaton-Higgs potential



Studying the phase transition in the Dilaton-Higgs potential




Studying the phase transition in the Dilaton-Higgs potential




Espinosa, Gripaios,
Konstandin, Riva '11

Studying the phase transition in the Dilaton-Higgs potential




Espinosa, Gripaios,
Konstandin, Riva '11

Studying the phase transition in the Dilaton-Higgs potential




Espinosa, Gripaios,
Konstandin, Riva '11

Studying the phase transition in the Dilaton-Higgs potential




Studying the phase transition in the Dilaton-Higgs potential



Espinosa, Gripaios,
Konstandin, Riva '11

- What are the conditions for trajectories b & c?


Studying the phase transition in the Dilaton-Higgs potential



Espinosa, Gripaios,
Konstandin, Riva '11

- What are the conditions for trajectories b & c ?
- Is the confinement PT strongly first-order?

Studying the phase transition in the Dilaton-Higgs potential



- What are the conditions for trajectories b & c?
- Is the confinement PT strongly first-order?
- Is there enough CP violation along the trajectories b & c?


Partial compositeness and Higgs potential

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R$


Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R$



Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$



Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops
 - $V_h \sim \alpha \sin^2 \frac{h}{f} + \beta \sin^4 \frac{h}{f}$

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops
 - $V_h \sim \alpha \sin^2 \frac{h}{f} + \beta \sin^4 \frac{h}{f}$ with $\alpha \sim \beta \sim y^2$

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops
 - $V_h \sim \alpha \sin^2 \frac{h}{f} + \beta \sin^4 \frac{h}{f}$ with $\alpha \sim \beta \sim y^2$ and $f \sim \langle \chi \rangle$

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops
 - $V_h \sim \alpha \sin^2 \frac{h}{f} + \beta \sin^4 \frac{h}{f}$ with $\alpha \sim \beta \sim y^2$ and $f \sim \langle \chi \rangle$
 - However, $v \ll f \Rightarrow |\alpha^0/\beta^0| \sim v^2/f^2$ with $f \gtrsim 0.8 \text{ TeV}$

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops
 - $V_h \sim \alpha \sin^2 \frac{h}{f} + \beta \sin^4 \frac{h}{f}$ with $\alpha \sim \beta \sim y^2$ and $f \sim \langle \chi \rangle$
 - However, $v \ll f \Rightarrow |\alpha^0/\beta^0| \sim v^2/f^2$ with $f \gtrsim 0.8 \text{ TeV}$
 - Potential quickly detunes for $\chi \neq \chi_0$

Partial compositeness and Higgs potential


- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops
 - $V_h \sim \alpha \sin^2 \frac{h}{f} + \beta \sin^4 \frac{h}{f}$ with $\alpha \sim \beta \sim y^2$ and $f \sim \langle \chi \rangle$
 - However, $v \ll f \Rightarrow |\alpha^0/\beta^0| \sim v^2/f^2$ with $f \gtrsim 0.8 \text{ TeV}$
 - Potential quickly detunes for $\chi \neq \chi_0$
 $\alpha, \beta > 0 \Rightarrow \mathbf{valley \ along} \quad h/\chi = 0 \quad \mathbf{Trajectory \ a}$

Partial compositeness and Higgs potential

- SM Yukawas from linear mixing $y_L \bar{q}_L \mathcal{O}_L + y_R \bar{q}_R \mathcal{O}_R \Rightarrow \lambda_q \sim \frac{y_L[\chi] y_R[\chi]}{g_\star}$
- Mixings explicitly break Goldstone shift symmetry
 - Higgs potential generated by SM loops
 - $V_h \sim \alpha \sin^2 \frac{h}{f} + \beta \sin^4 \frac{h}{f}$ with $\alpha \sim \beta \sim y^2$ and $f \sim \langle \chi \rangle$
 - However, $v \ll f \Rightarrow |\alpha^0/\beta^0| \sim v^2/f^2$ with $f \gtrsim 0.8 \text{ TeV}$
 - Potential quickly detunes for $\chi \neq \chi_0$

$\alpha, \beta > 0$	\Rightarrow	valley along	$h/\chi = 0$	Trajectory a
$\alpha, \beta < 0$	\Rightarrow	valley along	$h/\chi \sim \pi/2$	Trajectory c


Studying the phase transition in the Dilaton-Higgs potential



- What are the conditions for trajectories b & c ? 
- Is the confinement PT strongly first-order?
- Is there enough CP violation along the trajectories b & c ?

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:



Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4$$




Typical coupling of the dilaton:

Meson-like dilaton: $g_\chi \sim N^{-1/2}$

Glueball-like dilaton: $g_\chi \sim N^{-1}$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:



Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$




Typical coupling of the dilaton:

Meson-like dilaton: $g_\chi \sim N^{-1/2}$

Glueball-like dilaton: $g_\chi \sim N^{-1}$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential
- From dimensional analysis and large-N counting: $F_{CFT}[\chi = 0] \simeq -cN^2T^4$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential
- From dimensional analysis and large-N counting: $F_{CFT}[\chi = 0] \simeq -cN^2T^4$
- Use $\frac{45N^2}{4}$ strong sector bosons to reproduce $\mathcal{N} = 4$ SYM result for c .

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential
- From dimensional analysis and large-N counting: $F_{CFT}[\chi = 0] \simeq -cN^2 T^4$
- Use $\frac{45N^2}{4}$ strong sector bosons to reproduce $\mathcal{N} = 4$ SYM result for c .

Studying the phase transition in the Dilaton-Higgs potential



- What are the conditions for trajectories b & c ?
- Is the confinement PT strongly first-order?
- Is there enough CP violation along the trajectories b & c ?

CP violation

$$m = |m|e^{i\theta}$$

$$S \propto \text{Im} \left[V^\dagger m^{\dagger''} m V \right] = (|m|^2 \theta')'$$

1 Flavour

Multi-Flavour

CP violation

$$m = |m|e^{i\theta}$$

$$S \propto \text{Im} [V^\dagger m^{\dagger''} m V] = (|m|^2 \theta')'$$

1 Flavour

Agrees with semi-classical treatment

Multi-Flavour

CP violation

$$m = |m|e^{i\theta}$$

$$S \propto \text{Im} [V^\dagger m^{\dagger''} m V] = (|m|^2 \theta')'$$

1 Flavour

θ has to be space dependent!

Agrees with semi-classical treatment

$$\lambda(z) = e^{i\theta_1} f(z) + e^{i\theta_2} g(z)$$

Multi-Flavour

CP violation

$$m = |m|e^{i\theta}$$

$$S \propto \text{Im} [V^\dagger m^{\dagger''} m V] = (|m|^2 \theta')'$$

1 Flavour

θ has to be space dependent!

Agrees with semi-classical treatment

$$\lambda(z) = e^{i\theta_1} f(z) + e^{i\theta_2} g(z)$$

- No varying phase needed

Multi-Flavour

CP violation

$$m = |m|e^{i\theta}$$

$$S \propto \text{Im} [V^\dagger m^{\dagger''} m V] = (|m|^2 \theta')'$$

1 Flavour

θ has to be space dependent!

Agrees with semi-classical treatment

$$\lambda(z) = e^{i\theta_1} f(z) + e^{i\theta_2} g(z)$$

- No varying phase needed
- Need sizeable off-diagonal Yukawas

Multi-Flavour

CP violation

$$m = |m|e^{i\theta}$$

$$S \propto \text{Im} [V^\dagger m^{\dagger''} m V] = (|m|^2 \theta')'$$

1 Flavour


θ has to be space dependent!

Agrees with semi-classical treatment

$$\lambda(z) = e^{i\theta_1} f(z) + e^{i\theta_2} g(z)$$

- No varying phase needed
- Need sizeable off-diagonal Yukawas
- Flavour symmetries can significantly influence the final result

Studying the phase transition in the Dilaton-Higgs potential




- What are the conditions for trajectories b & c ?
- Is the confinement PT strongly first-order?
- Is there enough CP violation along the trajectories b & c ?

Two field tunneling

One field

$$S_E \sim 140$$

$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$




Two field tunneling

One field

$$S_E \sim 140$$

$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$

$$\alpha = \begin{cases} 3 & O(4) \\ 2 & O(3) \end{cases}$$



Two field tunneling


One field

$$S_E \sim 140$$

$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$

$$\alpha = \begin{cases} 3 & O(4) \\ 2 & O(3) \end{cases}$$

$r \sim \text{"time"}$



Two field tunneling

One field


$$S_E \sim 140$$

$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$

$$\phi(r \rightarrow \infty) = 0$$

$$\alpha = \begin{cases} 3 & O(4) \\ 2 & O(3) \end{cases}$$

$r \sim$ “time”



Two field tunneling

One field

$$S_E \sim 140$$


$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$

$$\phi(r \rightarrow \infty) = 0$$

$$\left. \frac{d\phi}{dr} \right|_{r=0} = 0$$

$$\alpha = \begin{cases} 3 & O(4) \\ 2 & O(3) \end{cases}$$

$r \sim$ “time”



Two field tunneling

One field

$$S_E \sim 140$$


$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$

$$\phi(r \rightarrow \infty) = 0$$

$$\left. \frac{d\phi}{dr} \right|_{r=0} = 0$$

$$\alpha = \begin{cases} 3 & O(4) \\ 2 & O(3) \end{cases}$$

$r \sim$ “time”



Two field tunneling

One field

$$S_E \sim 140$$


$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$

$$\phi(r \rightarrow \infty) = 0$$

$$\left. \frac{d\phi}{dr} \right|_{r=0} = 0$$

$$\alpha = \begin{cases} 3 & O(4) \\ 2 & O(3) \end{cases}$$

$r \sim$ “time”



Two field tunneling

One field

$$S_E \sim 140$$


$$\frac{d^2\phi}{dr^2} + \frac{\alpha}{r} \frac{d\phi}{dr} = V'(\phi)$$

$$\phi(r \rightarrow \infty) = 0$$

$$\left. \frac{d\phi}{dr} \right|_{r=0} = 0$$

$$\alpha = \begin{cases} 3 & O(4) \\ 2 & O(3) \end{cases}$$

$r \sim$ “time”



Two field tunneling

Two fields

$$S_E \sim 140$$


$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Two field tunneling

Two fields

$$S_E \sim 140$$

$$\frac{d^2 \vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$




Two field tunneling

Two fields

$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$




Two field tunneling

Two fields

$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$




Two field tunneling

Two fields

$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$




Two field tunneling

Two fields

$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Guess $\vec{\phi}_g(x)$




Two field tunneling

Two fields

$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Guess $\vec{\phi}_g(x)$



Two field tunneling

Two fields


$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Guess $\vec{\phi}_g(x)$

$$\frac{d^2x}{dr^2} + \frac{\alpha}{r} \frac{dx}{dr} = \partial_x V(\vec{\phi}_g(x))$$

$$\frac{d^2\vec{\phi}_g}{dx^2} \left(\frac{dx}{dr} \right)^2 - \nabla_{\perp} V(\vec{\phi}_g(x)) = 0$$



Two field tunneling

Two fields


$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Guess $\vec{\phi}_g(x)$

$$\frac{d^2x}{dr^2} + \frac{\alpha}{r} \frac{dx}{dr} = \partial_x V(\vec{\phi}_g(x))$$

$$\underbrace{\frac{d^2\vec{\phi}_g}{dx^2} \left(\frac{dx}{dr} \right)^2 - \nabla_{\perp} V(\vec{\phi}_g(x))}_{\equiv \vec{N}} = 0$$



Two field tunneling

Two fields


$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Guess $\vec{\phi}_g(x)$

$$\frac{d^2x}{dr^2} + \frac{\alpha}{r} \frac{dx}{dr} = \partial_x V(\vec{\phi}_g(x))$$

$$\underbrace{\frac{d^2\vec{\phi}_g}{dx^2} \left(\frac{dx}{dr} \right)^2 - \nabla_{\perp} V(\vec{\phi}_g(x))}_{\equiv \vec{N}} = 0$$



Two field tunneling

Two fields


$$S_E \sim 140$$

$$\frac{d^2 \vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Guess $\vec{\phi}_g(x)$

$$\frac{d^2 x}{dr^2} + \frac{\alpha}{r} \frac{dx}{dr} = \partial_x V(\vec{\phi}_g(x))$$

$$\underbrace{\frac{d^2 \vec{\phi}_g}{dx^2} \left(\frac{dx}{dr} \right)^2 - \nabla_{\perp} V(\vec{\phi}_g(x))}_{\equiv \vec{N}} = 0$$



Two field tunneling

Two fields


$$S_E \sim 140$$

$$\frac{d^2\vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$

Guess $\vec{\phi}_g(x)$

$$\frac{d^2x}{dr^2} + \frac{\alpha}{r} \frac{dx}{dr} = \partial_x V(\vec{\phi}_g(x))$$

$$\underbrace{\frac{d^2\vec{\phi}_g}{dx^2} \left(\frac{dx}{dr} \right)^2 - \nabla_{\perp} V(\vec{\phi}_g(x))}_{\equiv \vec{N}} = 0$$



Two field tunneling

Two fields


$$S_E \sim 140$$

$$\frac{d^2 \vec{\phi}}{dr^2} + \frac{\alpha}{r} \frac{d\vec{\phi}}{dr} = \nabla V(\vec{\phi})$$



Guess $\vec{\phi}_g(x)$

$$\frac{d^2 x}{dr^2} + \frac{\alpha}{r} \frac{dx}{dr} = \partial_x V(\vec{\phi}_g(x))$$



$$\underbrace{\frac{d^2 \vec{\phi}_g}{dx^2} \left(\frac{dx}{dr} \right)^2 - \nabla_{\perp} V(\vec{\phi}_g(x))}_{\equiv \vec{N}} = 0$$





The phase transition



The phase transition




The phase transition




colours in
CFT, used in
large N expansion

The phase transition




colours in
CFT, used in
large N expansion




Generic mass
of composite
resonances
 $m_* = g_* f$


The phase transition



No viable EW
Minimum found




The phase transition





No viable EW
Minimum found

Reheating temperature
s.t. $h/T_{\text{reheat}} < 1$





The phase transition



The phase transition

Meson-like
dilaton



$$g_\chi = \frac{4\pi}{\sqrt{N}}$$



The phase transition

Glueball-like
dilaton




$$g_\chi = \frac{4\pi}{N}$$



The phase transition

Meson-like
dilaton




$$g_\chi = \frac{4\pi}{\sqrt{N}}$$



The phase transition

Glueball-like
dilaton

$$g_\chi = \frac{4\pi}{N}$$



Experimental signatures

Experimental signatures

- Light dilaton

Experimental signatures

- Light dilaton
- Modification of Yukawa coupling

Experimental signatures

- Light dilaton
- Modification of Yukawa coupling
 - Real part: Measurable at LHC and future linear colliders

Experimental signatures

- Light dilaton
- Modification of Yukawa coupling
 - Real part: Measurable at LHC and future linear colliders
 - Imaginary part: electron and neutron EDMs

Experimental signatures

- Light dilaton
- Modification of Yukawa coupling
 - Real part: Measurable at LHC and future linear colliders
 - Imaginary part: electron and neutron EDMs
- Gravitational waves in LISA spectrum

Summary

- Framework for CP-violation and diffusion for z-dependent Yukawas.
- Fully consistent and general formalism (diffusion and CP-violation from first principle).
- Application possible to low-scale flavour physics (Froggatt-Nielsen, Randall-Sundrum, Composite Higgs etc.)

Baldes, Konstandin,
Servant '16
1604.04526 & 1608.03254


Von Harling, Servant '16
1612.02447

SB, Matsedonskyi,
Von Harling, Servant
1803.08546, 1804.07314


Summary

- Composite Higgs is a prime model for EWBG with varying Yukawas
 - Strong first-order PT from combined confinement-EWPT
 - Varying Yukawas from partial compositeness
 - Detectable signal in gravitational waves
 - Detection possible in electron/neutron EDM

Phase transitions




First-order phase transition




Higher order phase transition

Phase transitions




First-order phase transition




Higher order phase transition

In the SM: not first order

Phase transitions



First-order phase transition




Higher order phase transition

In the SM: not first order


Yukawa variation:

Baldes, Konstandin, Servant '16

Phase transitions



First-order phase transition



Higher order phase transition

In the SM: not first order


Yukawa variation:

Singlet extensions:


Baldes, Konstandin, Servant '16

Espinosa, Gripaios,
Konstandin, Riva '11

Phase transitions



First-order phase transition



Higher order phase transition

In the SM: not first order

Yukawa variation:

Singlet extensions:


Higher order operators:

Baldes, Konstandin, Servant '16


Espinosa, Gripaios,
Konstandin, Riva '11

Delaunay, Grojean, Wells '07
Grinstein, Trott '08

Phase transitions



First-order phase transition



Higher order phase transition

In the SM: not first order

Yukawa variation:

Singlet extensions:

Higher order operators:

Joint confinement-electroweak phase transition: stay tuned


Baldes, Konstandin, Servant '16

Espinosa, Gripaios,
Konstandin, Riva '11

Delaunay, Grojean, Wells '07
Grinstein, Trott '08


Baryon asymmetry

$$\eta_B = \frac{n_B(-\infty)}{s} = \frac{135 N_c}{4\pi^2 v_w g_*} \int_{-\infty}^{+\infty} dz \Gamma_{ws} \mu_L \text{Exp} \left[-\frac{3}{2} A \frac{1}{v_w} \int_{-\infty}^z dz_0 \Gamma_{ws} \right]$$



1. Sakharov condition (B-violation)
2. Sakharov condition (CP-violation)
3. Sakharov condition (out of equilibrium)


Baryon asymmetry



1. Sakharov condition (B-violation)
2. Sakharov condition (CP-violation)
3. Sakharov condition (out of equilibrium)

Baryon asymmetry


$$\eta_B = \frac{n_B(-\infty)}{s} = \frac{135 N_c}{4\pi^2 v_w g_*} \int_{-\infty}^{+\infty} dz \Gamma_{ws} \mu_L \text{Exp} \left[-\frac{3}{2} A \frac{1}{v_w} \int_{-\infty}^z dz_0 \Gamma_{ws} \right]$$



1. Sakharov condition (B-violation)
2. Sakharov condition (CP-violation)
3. Sakharov condition (out of equilibrium)

Baryon asymmetry

$$\eta_B = \frac{n_B(-\infty)}{s} = \frac{135 N_c}{4\pi^2 v_w g_*} \int_{-\infty}^{+\infty} dz \Gamma_{ws} \mu_L \text{Exp} \left[-\frac{3}{2} A \frac{1}{v_w} \int_{-\infty}^z dz_0 \Gamma_{ws} \right]$$




1. Sakharov condition (B-violation)

2. Sakharov condition (CP-violation)

3. Sakharov condition (out of equilibrium)

Baryon asymmetry

$$\eta_B = \frac{n_B(-\infty)}{s} = \frac{135 N_c}{4\pi^2 v_w g_*} \int_{-\infty}^{+\infty} dz \Gamma_{ws} \mu_L \text{Exp} \left[-\frac{3}{2} A \frac{1}{v_w} \int_{-\infty}^z dz_0 \Gamma_{ws} \right]$$



1. Sakharov condition (B-violation)
2. Sakharov condition (CP-violation)
3. Sakharov condition (out of equilibrium)

System and Kernel

$$A(z) v'(z) + B(z) v(z) = S(z)$$

Unknowns



$$(\mu_{t_{R/L}}, \mu_{b_{R/L}}, \mu_{s_{R/L}}, \mu_{c_{R/L}}, \mu_h \\ u_{t_{R/L}}, u_{b_{R/L}}, u_{s_{R/L}}, u_{c_{R/L}}, u_h)$$

System and Kernel

$$A(z) v'(z) + B(z) v(z) = S(z)$$

Unknowns

$$\begin{aligned} &(\mu_{t_{R/L}}, \mu_{b_{R/L}}, \mu_{s_{R/L}}, \mu_{c_{R/L}}, \mu_h \\ &u_{t_{R/L}}, u_{b_{R/L}}, u_{s_{R/L}}, u_{c_{R/L}}, u_h) \end{aligned}$$



$$\mu(z) = \int_{-\infty}^{+\infty} dz_0 G(z, z_0) S(z_0)$$


System and Kernel

$$A(z) v'(z) + B(z) v(z) = S(z)$$

Unknowns

$$(\mu_{t_{R/L}}, \mu_{b_{R/L}}, \mu_{s_{R/L}}, \mu_{c_{R/L}}, \mu_h \\ u_{t_{R/L}}, u_{b_{R/L}}, u_{s_{R/L}}, u_{c_{R/L}}, u_h)$$

$$\eta_B = \int_{-\infty}^{+\infty} dz \# \Gamma_{ws}(z) e^{-\# z} \mu_L(z)$$



$$\mu(z) = \int_{-\infty}^{+\infty} dz_0 G(z, z_0) S(z_0)$$

System and Kernel

$$A(z) v'(z) + B(z) v(z) = S(z)$$

Unknowns


$$(\mu_{t_{R/L}}, \mu_{b_{R/L}}, \mu_{s_{R/L}}, \mu_{c_{R/L}}, \mu_h \\ u_{t_{R/L}}, u_{b_{R/L}}, u_{s_{R/L}}, u_{c_{R/L}}, u_h)$$

$$\eta_B = \int_{-\infty}^{+\infty} dz \# \Gamma_{ws}(z) e^{-\# z} \mu_L(z)$$

$$\eta_B = \sum_i \int_{-\infty}^{+\infty} dz_0 K_i(z_0) S_i(z_0)$$

Kernel

- Weak Sphaleron
- Interactions (GF)
- Numerical factors



Source

- CP-violation

Varying Yukawas across the wall

Effective description following from Flavon-Higgs coupling

Broken phase

$$y(y_0, y_1, \phi, n) = (y_0 - y_1) \left[1 - \left(\frac{\phi}{v} \right)^n \right] + y_1$$


↑
Symmetric phase
Higgs vev

Free parameter

Higgs vev in
broken phase


Baldes, Konstandin,
Servant '16

1608.03254 & 1604.04526



Yukawas

$$Y_{tc}(z, n) = \begin{pmatrix} e^i y(1, 0.008, \phi(z), n) & y(1, 0.04, \phi(z), n) \\ y(1, 0.2, \phi(z), n) & y(1, 1, \phi(z), n) \end{pmatrix}$$




← Broken phase

Symmetric phase →

Yukawas

$$Y_{tc}(z, n) = \begin{pmatrix} e^i y(1, 0.008, \phi(z), n) & y(1, 0.04, \phi(z), n) \\ y(1, 0.2, \phi(z), n) & y(1, 1, \phi(z), n) \end{pmatrix}$$



← Broken phase


Symmetric phase →

$$\eta_B = \sum_i \int_{-\infty}^{+\infty} dz_0 K_i(z_0) S_i(z_0)$$


$$\eta_B \approx 5.5 \cdot 10^{-10}$$

$$\eta_B \approx 2.1 \cdot 10^{-12}$$


Kernel for n=10




Source for n=10



Kernel for n=0.1



Source for n=0.1




$$\eta_B = \sum_i \int_{-\infty}^{+\infty} dz_0 K_i(z_0) S_i(z_0)$$


$$\eta_B \approx 5.5 \cdot 10^{-10}$$

$$\eta_B \approx 2.1 \cdot 10^{-12}$$


Kernel for n=10




Source for n=10



Kernel for n=0.1



Source for n=0.1




$$\eta_B = \sum_i \int_{-\infty}^{+\infty} dz_0 K_i(z_0) S_i(z_0)$$

$$\eta_B \approx 5.5 \cdot 10^{-10}$$


Kernel for n=10

K(z)



Source for n=10


S(z)



$$\eta_B \approx 2.1 \cdot 10^{-12}$$


Kernel for n=0.1

K(z)



Source for n=0.1

S(z)




$$\eta_B = \sum_i \int_{-\infty}^{+\infty} dz_0 K_i(z_0) S_i(z_0)$$


$$\eta_B \approx 5.5 \cdot 10^{-10}$$

$$\eta_B \approx 2.1 \cdot 10^{-12}$$


Kernel for n=10




Source for n=10



Kernel for n=0.1





Source for n=0.1



Experimental signatures

$$\Gamma_{qqh} \sim -\beta_{\lambda_q} s_\delta \sin[v/f] + \lambda_q (c_\delta \cos[v/f] - s_\delta \sin[v/f])$$





Experimental signatures

Beta function of Yukawa
coupling of quark q



$$\Gamma_{qqh} \sim -\beta_{\lambda_q} s_\delta \sin[v/f] + \lambda_q (c_\delta \cos[v/f] - s_\delta \sin[v/f])$$





Experimental signatures

Beta function of Yukawa
coupling of quark q




$$\Gamma_{qqh} \sim -\beta_{\lambda_q} s_\delta \sin[v/f] + \lambda_q c_\delta \cos[v/f] - s_\delta \sin[v/f])$$




Experimental signatures

$$\Gamma_{qqh} \sim -\beta_{\lambda_q} s_\delta \sin[v/f] + \lambda_q c_\delta \cos[v/f] - s_\delta \sin[v/f])$$



CLIC sensitivity on $\text{Re} [\delta\lambda_t] \sim 4\%$ at 1σ

Abramowicz et al. '13




$\text{Im} [\delta\lambda_t] \lesssim 0.018$ @ 90% CL


Cirigliano, Dekens,
de Vries, Mereghetti '16

Experimental signatures

$$\alpha = \frac{\epsilon}{\rho_{\text{rad}}} \simeq \frac{(V_{\text{tot}}[0,0] - V_{\text{tot}}[v, \chi_0])_{T_n}}{3\pi^2 N^2 T_n^4 / 8}, \quad \left. \frac{\beta}{H} \simeq T_n \frac{dS_E}{dT} \right|_{T_n}$$



Meson





LISA: $\alpha \gtrsim 0.1$ & $1 \lesssim \beta/H \lesssim 10^4$

Grojean, Servant '06
Caprini et al. '15

Experimental signatures

$$\alpha = \frac{\epsilon}{\rho_{\text{rad}}} \simeq \frac{(V_{\text{tot}}[0,0] - V_{\text{tot}}[v, \chi_0])_{T_n}}{3\pi^2 N^2 T_n^4 / 8}, \quad \frac{\beta}{H} \simeq T_n \frac{dS_E}{dT} \Big|_{T_n}$$



LISA: $\alpha \gtrsim 0.1$ & $1 \lesssim \beta/H \lesssim 10^4$

Grojean, Servant '06
Caprini et al. '15

CP Violation

$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)^{''} m V \right]_{ii}$$

CP Violation

$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)^{''} m V \right]_{ii}$$

- Top-only: varying complex phase; couple top right to two operators

CP Violation

$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)^{''} m V \right]_{ii}$$

- Top-only: varying complex phase; couple top right to two operators
- Multi-flavour: $y \bar{q} \mathcal{O} \rightarrow y_{ij} \bar{q}_i \mathcal{O}_j \rightarrow \mathcal{I}_{ij} y_{jj} \bar{q}_i \mathcal{O}_j$

CP Violation

$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)^{''} m V \right]_{ii}$$

- Top-only: varying complex phase; couple top right to two operators
- Multi-flavour: $y \bar{q} \mathcal{O} \rightarrow y_{ij} \bar{q}_i \mathcal{O}_j \rightarrow \mathcal{I}_{ij} y_{jj} \bar{q}_i \mathcal{O}_j$

$$m_{ij} \sim (\mathcal{I}^L)_{ik} (y^L)_{kk} (g_\star^{-1})_{kl} (y^R)_{ll} (\mathcal{I}^R)_{lj} h$$

CP Violation

$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)'' m V \right]_{ii}$$

- Top-only: varying complex phase; couple top right to two operators
- Multi-flavour: $y \bar{q} \mathcal{O} \rightarrow y_{ij} \bar{q}_i \mathcal{O}_j \rightarrow \mathcal{I}_{ij} y_{jj} \bar{q}_i \mathcal{O}_j$


Partial compositeness


$$m_{ij} \sim (\mathcal{I}^L)_{ik} (y^L)_{kk} (g_\star^{-1})_{kl} (y^R)_{ll} (\mathcal{I}^R)_{lj} h$$

CP Violation

$$S_i^{CPV} \sim \text{Im} \left[V^\dagger (m^\dagger)'' m V \right]_{ii}$$

- Top-only: varying complex phase; couple top right to two operators
- Multi-flavour: $y \bar{q} \mathcal{O} \rightarrow y_{ij} \bar{q}_i \mathcal{O}_j \rightarrow \mathcal{I}_{ij} y_{jj} \bar{q}_i \mathcal{O}_j$




CP-Violation

$$m_{ij} \sim (\mathcal{I}^L)_{ik} (y^L)_{kk} (g_\star^{-1})_{kl} (y^R)_{ll} (\mathcal{I}^R)_{lj} h$$

\mathbb{I}_L	y_L	g_\star^{-1}	y_R	\mathbb{I}_R	condition on d_{ij} for $S_{\text{CPV}} = 0$
d	2	d	2	d	always
	1		1		always
d	2	\emptyset	2	d	$d_{11} + d_{22} = d_{12} + d_{21}$
	1		2		always
	2		1		always
	1		1		always
\emptyset	2	d	2	d	always
	1		1		always
\emptyset	2	\emptyset	2	d	$d_{11} = d_{12}, d_{22} = d_{21}$
	1		2		always
	2		1		$d_{11} = d_{21}$
	1		1		always
\emptyset	2	d	2	\emptyset	$d_{11} = d_{12} = d_{21} = d_{22}$
	1		1		always
\emptyset	2	\emptyset	2	\emptyset	$d_{11} = d_{12} = d_{21} = d_{22}$
	2		1		$d_{11} = d_{21}$
	1		2		$d_{11} = d_{12}$
	1		1		always

Baryon asymmetry: multi-flavour case with U(1) flavour symmetries

$$m_{ij} \sim (y^L)_{ii} (g_\star^{-1})_{ij} (y^R)_{jj} \sin\left(\frac{h}{f}\right) \cos\left(\frac{h}{f}\right) f \quad \text{Csaki, Falkowski, Weiler '08}$$




Baryon asymmetry: multi-flavour case with U(1) flavour symmetries

$$m_{ij} \sim (y^L)_{ii} (g_\star^{-1})_{ij} (y^R)_{jj} \sin\left(\frac{h}{f}\right) \cos\left(\frac{h}{f}\right) f$$

Csaki, Falkowski,
Weiler '08

Anarchic




Baryon asymmetry: multi-flavour case with U(1) flavour symmetries

$$m_{ij} \sim (y^L)_{ii} (g_\star^{-1})_{ij} (y^R)_{jj} \sin\left(\frac{h}{f}\right) \cos\left(\frac{h}{f}\right) f$$


Csaki, Falkowski,
Weiler '08

Anarchic
Two large entries




Baryon asymmetry: top-only case

$$\mathcal{L} \supset y_L \bar{t}_L \mathcal{O}_L + y_R^1 \bar{t}_R^1 \mathcal{O}_R^1 + y_R^2 \bar{t}_R^2 \mathcal{O}_R^2$$



$$\lambda_t [\chi] \sim \frac{y_L [\chi] (y_R^1 [\chi] + y_R^2 [\chi])}{g_\star}$$


Baryogenesis



Dimensional Analysis and large N

Idea: restore dimensions of \hbar

$$[V] = \frac{[\hbar]}{L^4}, \quad [\phi] = [f] = \frac{\sqrt{[\hbar]}}{L}, \quad [g] = \frac{1}{\sqrt{[\hbar]}}, \quad [\partial_\mu] = \frac{1}{L}$$




$$V \sim g^2 \phi^4 \quad \text{or} \quad V \sim g^2 f^4$$

Dimensional Analysis and large N

Idea: restore dimensions of \hbar

$$[V] = \frac{[\hbar]}{L^4}, \quad [\phi] = [f] = \frac{\sqrt{[\hbar]}}{L}, \quad [g] = \frac{1}{\sqrt{[\hbar]}}, \quad [\partial_\mu] = \frac{1}{L}$$



$$V \sim g^2 \phi^4 \quad \text{or} \quad V \sim g^2 f^4$$

Large N expansion: for a meson like Higgs $g_\star \sim \frac{4\pi}{\sqrt{N}}$

Canonically normalised fields

$$\frac{1}{2}(\partial_\mu h)^2$$

$h \rightarrow h + 2\pi f k$ ($k \in \mathbb{Z}$) for constant f

$h \rightarrow h + 2\pi \chi k$ ($k \in \mathbb{Z}$) for dynamical χ

$$\theta = h/\chi : \quad \theta \rightarrow \theta + 2\pi k \quad \mathcal{L} = \frac{1}{2}\chi^2(\partial_\mu \theta)^2 + \frac{1}{2}(\partial_\mu \chi)^2$$

We need canonically normalised fields for vacuum tunneling:

$$\chi_1 = \chi \sin \theta, \quad \chi_2 = \chi \cos \theta$$

Canonically normalised fields

$$\frac{1}{2}(\partial_\mu h)^2$$

$h \rightarrow h + 2\pi f k$ ($k \in \mathbb{Z}$) for constant f



$h \rightarrow h + 2\pi \chi k$ ($k \in \mathbb{Z}$) for dynamical χ

$$\theta = h/\chi : \quad \theta \rightarrow \theta + 2\pi k \quad \mathcal{L} = \frac{1}{2}\chi^2(\partial_\mu \theta)^2 + \frac{1}{2}(\partial_\mu \chi)^2$$

We need canonically normalised fields for vacuum tunneling:

$$\chi_1 = \chi \sin \theta, \quad \chi_2 = \chi \cos \theta$$

Canonically normalised fields

$$\frac{1}{2}(\partial_\mu h)^2$$

$h \rightarrow h + 2\pi f k$ ($k \in \mathbb{Z}$) for constant f



$h \rightarrow h + 2\pi \chi k$ ($k \in \mathbb{Z}$) for dynamical χ



$$\theta = h/\chi : \quad \theta \rightarrow \theta + 2\pi k \quad \mathcal{L} = \frac{1}{2}\chi^2(\partial_\mu \theta)^2 + \frac{1}{2}(\partial_\mu \chi)^2$$

We need canonically normalised fields for vacuum tunneling:

$$\chi_1 = \chi \sin \theta, \quad \chi_2 = \chi \cos \theta$$

Canonically normalised fields

$$\frac{1}{2}(\partial_\mu h)^2$$

$h \rightarrow h + 2\pi f k$ ($k \in \mathbb{Z}$) for constant f



$h \rightarrow h + 2\pi \chi k$ ($k \in \mathbb{Z}$) for dynamical χ



$$\theta = h/\chi : \quad \theta \rightarrow \theta + 2\pi k \quad \mathcal{L} = \frac{1}{2}\chi^2(\partial_\mu \theta)^2 + \frac{1}{2}(\partial_\mu \chi)^2$$




We need canonically normalised fields for vacuum tunneling:

$$\chi_1 = \chi \sin \theta, \quad \chi_2 = \chi \cos \theta$$

Temperature Corrections

- Hard to compute in the symmetric phase
- From dimensional analysis and large-N counting:
 $F_{CFT}[\chi = 0] \simeq -cN^2T^4$ where c depends on the number of d.o.f. per color in the strong sector
- For definiteness we use a 5-D estimate to fix c:

$$F_{CFT}[\chi = 0] \simeq -\frac{\pi^2 N^2}{8} T^4$$



Temperature Corrections

Temperature Corrections

- Use standard one loop result to interpolate between the symmetric phase and the broken phase.

Temperature Corrections

- Use standard one loop result to interpolate between the symmetric phase and the broken phase.
- We use standard one loop result for SM fermions

Temperature Corrections


- Use standard one loop result to interpolate between the symmetric phase and the broken phase.
- We use standard one loop result for SM fermions

$$\Delta V_{1-loop}^T = \sum_{\text{bosons}} \frac{n T^4}{2\pi^2} J_b \left[\frac{m^2}{T^2} \right] - \sum_{\text{fermions}} \frac{n T^4}{2\pi^2} J_f \left[\frac{m^2}{T^2} \right]$$

$$J_b[x] = \int_0^\infty dk \ k^2 \log \left[1 - e^{-\sqrt{k^2+x}} \right] \quad \text{and} \quad J_f[x] = \int_0^\infty dk \ k^2 \log \left[1 + e^{-\sqrt{k^2+x}} \right]$$

$$\sum_{\text{CFT bosons}} n + \sum_{\text{CFT fermions}} n = \frac{45N^2}{8\pi^2}$$

Temperature Evolution



Collider phenomenology

Collider phenomenology

- Detect Dilaton

Collider phenomenology

- Detect Dilaton
- Fit parameters of EFT to data

Collider phenomenology

- Detect Dilaton
- Fit parameters of EFT to data
- E.g.: Dilaton quark coupling

Collider phenomenology

- Detect Dilaton
- Fit parameters of EFT to data
- E.g.: Dilaton quark coupling

$$V_{qq\chi} \sim c_\delta \beta_{\lambda_q} \frac{v}{f} + c_\delta \lambda_q \frac{v}{f} + s_\delta \lambda_q$$

Collider phenomenology

- Detect Dilaton
- Fit parameters of EFT to data
- E.g.: Dilaton quark coupling


Beta function of Yukawa
coupling of quark q

$$V_{qq\chi} \sim c_\delta \beta_{\lambda_q} \frac{v}{f} + c_\delta \lambda_q \frac{v}{f} + s_\delta \lambda_q$$


Collider phenomenology

- Detect Dilaton
- Fit parameters of EFT to data
- E.g.: Dilaton quark coupling

Beta function of Yukawa
coupling of quark q


$$V_{q\bar{q}\chi} \sim c_\delta \beta_{\lambda_q} \frac{v}{f} + c_\delta \lambda_q \frac{v}{f} + s_\delta \lambda_q$$


Higgs-Dilaton mass mixing.
Depend on parameters of potential.

Collider phenomenology

- Detect Dilaton
- Fit parameters of EFT to data
- E.g.: Higgs quark coupling

Beta function of Yukawa
coupling of quark q

$$\Gamma_{qqh} \sim -\beta_{\lambda_q} s_\delta \sin[v/f] + \lambda_q (c_\delta \cos[v/f] - s_\delta \sin[v/f])$$


Higgs-Dilaton mass mixing.
Depend on parameters of potential.

Froggat-Nielsen

$U(1)_{FN}$ with two FN fields χ, σ

Yukawa type interactions after SSB

$$\mathcal{L} \supset \tilde{y}_{ij} \left(\frac{\langle \chi \rangle}{\Lambda_\chi} \right)^{\tilde{n}_{ij}} \bar{Q}_i \tilde{\phi} U_j + y_{ij} \left(\frac{\langle \chi \rangle}{\Lambda_\chi} \right)^{n_{ij}} \bar{Q}_i \phi D_j$$

$$+ \tilde{Y}_{ij} \left(\frac{\langle \sigma \rangle}{\Lambda_\sigma} \right)^{\tilde{n}_{ij}} \bar{Q}_i \tilde{\phi} U_j + Y_{ij} \left(\frac{\langle \sigma \rangle}{\Lambda_\sigma} \right)^{n_{ij}} \bar{Q}_i \phi D_j$$

VEVs during EWSB


$$\phi : 0 \rightarrow v_\phi \quad \sigma : \Lambda_\sigma/5 \rightarrow \Lambda_\sigma/5 \quad \chi : \Lambda_\chi \rightarrow 0$$

$\chi - \phi$ -Potential

Charge assignment

$$Q_{FN}(\sigma) = Q_{FN}(\chi) = -1$$

\bar{Q}_3 (0)	\bar{Q}_2 (+2)	\bar{Q}_1 (+3)
U_3 (0)	U_2 (+1)	U_1 (+4)
D_3 (+2)	D_2 (+2)	D_1 (+3)



Froggat-Nielsen


$$Y_{t-c} = \begin{pmatrix} \tilde{y}_{cc}\epsilon_\chi^3 & \tilde{y}_{ct}\epsilon_\chi^2 \\ \tilde{y}_{tc}\epsilon_\chi^1 & \tilde{y}_{tt}\epsilon_\chi^0 \end{pmatrix} + \begin{pmatrix} \tilde{Y}_{cc}\epsilon_\sigma^3 & \tilde{Y}_{ct}\epsilon_\sigma^2 \\ \tilde{Y}_{tc}\epsilon_\sigma^1 & \tilde{Y}_{tt}\epsilon_\sigma^0 \end{pmatrix}$$

$$\tilde{y}_{i \neq j} = \tilde{Y}_{i \neq j} = 1$$


$$\tilde{y}_{tt} = \tilde{Y}_{tt} = 1/2$$

$$\tilde{y}_{cc} = \tilde{Y}_{cc} = e^{i\theta}$$

$$S(z) \cdot 10^6$$




$$K(z) \cdot 10^6$$





$$\eta_B = \sum_i \int_{-\infty}^{+\infty} dz_0 K_i(z_0) S_i(z_0) \simeq 1.15 \cdot 10^{-10}$$

Randall-Sundrum



Randall-Sundrum

$$\eta_B = \sum_i \int_{-\infty}^{+\infty} dz_0 K_i(z_0) S_i(z_0) \simeq 9.81 \cdot 10^{-11}$$



Deriving the equations

- Hermitian part of the Kadanoff-Baym equations
- Expand to second order in gradients (smooth background) and at tree level
- Neglect off-diagonals (fast flavour oscillations)
- Fluid type Ansatz for particle densities:

$$f_i = \frac{1}{e^{\beta(\omega_i + v_w k_z - \mu_i)} \pm 1} + \delta f_i$$


- Take different momenta and average over energy and momentum

Kinetic equations

$$\left(k_z \partial_z - \frac{1}{2} \left([V^\dagger (m^\dagger m)' V] \right)_{ii} \partial_{k_z} \right) f_{L,i} \approx \mathbf{C} + \mathcal{S}$$

$$\left(k_z \partial_z - \frac{1}{2} \left([V^\dagger (m^\dagger m)' V] \right)_{ii} \partial_{k_z} \right) f_{R,i} \approx \mathbf{C} - \mathcal{S}$$

Collision term



Source depends on m



link to

Yukawa couplings

CP-violating source term:

$$\mathcal{S} \equiv \frac{\text{sign}[k_z]}{2\tilde{k}} \text{Im} \left[V^\dagger m^{\dagger''} m V \right]_{ii} \partial_{k_z} f_{L/R,i}$$

(V are the Eigenvectors of $m^\dagger m$)

Network equations

Fluide type Ansatz:

$$f_i = \frac{1}{e^{\beta(\omega_i + v_w k_z - \mu_i)} \pm 1} + \delta f_i$$

$$u \equiv \left\langle \frac{k_z}{\omega_0} \delta f \right\rangle$$

CP-odd Energy-Momentum average, linear in: μ_i, u_i and v_w :

$$v_w K_1 \mu' + v_w (m^2)' K_2 \mu + u' - \langle \mathbf{C} \rangle = 0$$

$$-K_4 \mu' + v_w \tilde{K}_5 u' + v_w (m^2)' \tilde{K}_6 u - \left\langle \frac{k_z}{\omega_{0i}} \mathbf{C} \right\rangle = \pm v_w K_8 \text{Im} [V^\dagger m^{\dagger''} m V]$$

Interactions

Source

Network equations

Fluide type Ansatz:

$$f_i = \frac{1}{e^{\beta(\omega_i + v_w k_z - \mu_i)} \pm 1} + \delta f_i$$

$$u \equiv \left\langle \frac{k_z}{\omega_0} \delta f \right\rangle$$

CP-odd Energy-Momentum average, linear in: μ_i, u_i and v_w :

$$v_w K_1 \mu' + v_w (m^2)' K_2 \mu + u' - \langle \mathbf{C} \rangle = 0$$

$$-K_4 \mu' + v_w \tilde{K}_5 u' + v_w (m^2)' \tilde{K}_6 u - \left\langle \frac{k_z}{\omega_{0i}} \mathbf{C} \right\rangle = -v_w K_8 \text{Im} [V^\dagger m^{\dagger''} m V]$$

Interactions


Source

$$v_w\,K_1\,\mu' + v_w(m^2)'\,K_2\,\mu + u' - \Gamma^{\rm inel}\sum_i\mu_i = 0$$

$$-K_4\,\mu' + v_w\,\tilde{K}_5\,u' + v_w(m^2)'\,\tilde{K}_6\,u + \Gamma^{\rm tot}u = \pm v_w K_8\,{\rm Im}\left[V^\dagger {m^\dagger}'' m V\right]$$

$$v_w K_1 \mu' + v_w (m^2)' K_2 \mu + u' - \Gamma^{\text{inel}} \sum_i \mu_i = 0$$

$$-K_4 \mu' + v_w \tilde{K}_5 u' + v_w (m^2)' \tilde{K}_6 u + \Gamma^{\text{tot}} u = \pm v_w K_8 \text{Im} [V^\dagger m^{\dagger''} m V]$$



e.g. Fromme, Huber '06


hep-ph/0604159

Interactions:

Couple different particle species together

$$v_w K_1 \mu' + v_w (m^2)' K_2 \mu + u' - \Gamma^{\text{inel}} \sum_i \mu_i = 0$$

$$-K_4 \mu' + v_w \tilde{K}_5 u' + v_w (m^2)' \tilde{K}_6 u + \Gamma^{\text{tot}} u = \pm v_w K_8 \text{Im} [V^\dagger m^{\dagger''} m V]$$



e.g. Fromme, Huber '06

hep-ph/0604159 Interactions:

Couple different particle species together


- Yukawa interactions e.g. $t_L \leftrightarrow t_R + h$ $\Gamma_{y,q} = 4.2 \times 10^{-3} y_q^2 T$
- Helicity flip e.g. $t_L \leftrightarrow t_R$ $\Gamma_{m,q} = \frac{m_q^2}{63T}$
- W-scattering e.g. $t_L \leftrightarrow b_L$ $\Gamma_W = \frac{T}{60}$
- Higgs number violation $h \leftrightarrow 0$ $\Gamma_h = \frac{m_W^2}{50T}$
- Strong sphaleron all L \leftrightarrow all R $\Gamma_{ss} = 4.9 \times 10^{-4} T$

Collision term

$$\langle \mathbf{C} \rangle = \Gamma^{\text{inel}} \sum_i \mu_i, \quad \left\langle \frac{k_z}{\omega_{0i}} \mathbf{C} \right\rangle = -\Gamma^{\text{tot}} u$$

- Yukawa interactions e.g. $t_L \leftrightarrow t_R + h$ $\Gamma_{y,q} = 4.2 \times 10^{-3} y_q^2 T$
- Helicity flip e.g. $t_L \leftrightarrow t_R$ $\Gamma_{m,q} = \frac{m_q^2}{63T}$
- W-scattering e.g. $t_L \leftrightarrow b_L$ $\Gamma_W = \frac{T}{60}$
- Higgs number violation $h \leftrightarrow 0$ $\Gamma_h = \frac{m_W^2}{50T}$
- Strong sphaleron all L \leftrightarrow all R $\Gamma_{ss} = 4.9 \times 10^{-4} T$


$$\phi(z) = \frac{1}{2} \left(1 - \text{Tanh} \frac{z}{L_w} \right)$$



$$\Gamma_{ws} = 10^{-6} T \exp(-a\phi(z)/T)$$


$a \approx 40$

NEW




EW Baryogenesis

Cohen, Kaplan, Nelson '91




EW Baryogenesis

Cohen, Kaplan, Nelson '91




EW Baryogenesis

Cohen, Kaplan, Nelson '91



EW Baryogenesis

Cohen, Kaplan, Nelson '91




$\langle H \rangle = 0$
Sphaleron
active

Chiral asymmetry
converted to
Baryon asymmetry

CP-violation
at
phase interface

EW Baryogenesis


Cohen, Kaplan, Nelson '91



EW Baryogenesis

Cohen, Kaplan, Nelson '91

$\langle H \rangle \neq 0$
Sphaleron
inactive
↓
Baryon number
frozen



CP-violation
at
phase interface

$\langle H \rangle = 0$
Sphaleron
active
↓
Chiral asymmetry
converted to
Baryon asymmetry

CP-violation in the SM and beyond

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

Farrar, Shaposhnikov '93

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_{\substack{i>j \\ d,s,b}} (m_i^2 - m_j^2) J_{CP}$$

Gavela, *et al.* '93
Huet, Sather '94

Jarlskog constant



Based solely on
reflection coefficients

CP-violation in the SM and beyond

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_{\substack{i>j \\ u,c,t}} (m_i^2 + m_j^2)$$

Jarlskog constant

Much too small!
(measured: $\eta_B \sim 8.9 \cdot 10^{-11}$)

Popovshnikov '93

Gavela, et al. '93
Huet, Sather '94

Based solely on
reflection coefficients

CP-violation in the SM and beyond

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_{i>j} (m_i^2 + m_j^2)$$

Jarlskog constant

Much too small!
(measured: $\eta_B \sim 8.9 \cdot 10^{-11}$)

Popovshnikov '93

Gavela, *et al.* '93
Huet, Sather '94

Based solely on
reflection coefficients

Models with diffusion

2 Higgs doublet Cohen, Kaplan, Nelson '94

MSSM

Cline, Joyce, Kainulainen '97

Calculate CP-violating source,
inject to Boltzmann equation

CP-violation in the SM and beyond

In the SM: $\eta_B \lesssim 10^{-2} \Delta_{CP}$

$$\Delta_{CP} \sim (M_W^6 T_c^6)^{-1} \prod_{\substack{i>j \\ u,c,t}} (m_i^2 - m_j^2) \prod_{\substack{i>j \\ u,c,t}} (m_i^2 + m_j^2)$$

Jarlskog constant

Much too small!
(measured: $\eta_B \sim 8.9 \cdot 10^{-11}$)

Popovshnikov '93

Gavela, *et al.* '93
Huet, Sather '94

Based solely on
reflection coefficients

Models with diffusion

2 Higgs doublet Cohen, Kaplan, Nelson '94

MSSM

Cline, Joyce, Kainulainen '97

Calculate CP-
violating source,
inject to Boltzmann
equation

CP-violation and diffusion equation
from first principles (Kadanoff-Baym)

Prokopec, Schmidt,
Weinstock '03

Source for μ in SM

$$S \sim \Im [V^\dagger m^{\dagger''} m V]$$

$$m = y(z) \cdot \frac{\phi(z)}{\sqrt{2}}$$

Source for μ in SM

$$S \sim \Im [V^\dagger m^{\dagger''} m V]$$

For constant y :

$$S \sim \underbrace{\Im [V^\dagger y^\dagger y V]}_{=0} \phi'' \phi$$

$$m = y(z) \cdot \frac{\phi(z)}{\sqrt{2}}$$

Source for μ in SM

$$S \sim \Im [V^\dagger m^\dagger'' m V]$$

For constant y :

$$S \sim \underbrace{\Im [V^\dagger y^\dagger y V]}_{=0} \phi'' \phi$$



$\Rightarrow z$ dependent Yukawas

$$m = y(z) \cdot \frac{\phi(z)}{\sqrt{2}}$$



EW scale flavour physics

{ Composite Higgs
Froggatt-Nielsen
Randall-Sundrum
...

The phase transition in the SM and beyond





The phase transition in the SM and beyond



In the SM: not first order



The phase transition in the SM and beyond



In the SM: not first order
Yukawa variation:

Baldes, Konstandin, Servant '16

The phase transition in the SM and beyond



In the SM: not first order



Yukawa variation:

Singlet extensions:

Baldes, Konstandin, Servant '16

Espinosa, Gripaios,
Konstandin, Riva '11

The phase transition in the SM and beyond



In the SM: not first order

Yukawa variation:

Singlet extensions:


Minimal Composite Higgs: Up next

Baldes, Konstandin, Servant '16

Espinosa, Gripaios,
Konstandin, Riva '11

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:



Need small explicit breaking of conformal symmetry


$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

All scales in the strong sector (including f) are determined by χ

\Rightarrow Yukawas vary as χ varies

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:



Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4$$




Typical coupling of the dilaton:

Meson-like dilaton: $g_\chi \sim N^{-1/2}$

Glueball-like dilaton: $g_\chi \sim N^{-1}$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:



Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$




Typical coupling of the dilaton:

Meson-like dilaton: $g_\chi \sim N^{-1/2}$

Glueball-like dilaton: $g_\chi \sim N^{-1}$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential
- From dimensional analysis and large-N counting: $F_{CFT}[\chi = 0] \simeq -cN^2T^4$

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry

$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential
- From dimensional analysis and large-N counting: $F_{CFT}[\chi = 0] \simeq -cN^2 T^4$
- Use $\frac{45N^2}{4}$ strong sector bosons to reproduce $\mathcal{N} = 4$ SYM result for c .

Conformal symmetry breaking

The vev of the dilaton sets the scale of the strong sector:




Need small explicit breaking of conformal symmetry


$$V_\chi = c_\chi g_\chi^2 \chi^4 - \epsilon[\chi] \chi^4 + \dots$$

- For $T \ll \chi$ use 0-Temperature potential
- From dimensional analysis and large-N counting: $F_{CFT}[\chi = 0] \simeq -cN^2 T^4$
- Use $\frac{45N^2}{4}$ strong sector bosons to reproduce $\mathcal{N} = 4$ SYM result for c .


Quark masses in Composite Higgs through partial compositeness




Quark masses in Composite Higgs through partial compositeness



Quark masses in Composite Higgs through partial compositeness



Quark masses in Composite Higgs through partial compositeness




$$\mathcal{L} \supset y_i \bar{q}_i \mathcal{O}_i$$

$$\frac{\partial y_i}{\partial \log \mu} = \gamma_i y_i + c_i \frac{y_i^3}{g_\star^2} + \dots$$

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$


Fermionic operators
of the strong sector, e.g:

$$\mathcal{O}_i \sim \psi \bar{\psi} \psi$$




Goldstone matrix
containing the Higgs
 $U \sim \exp[ih/f]$


Quark masses in Composite Higgs through partial compositeness



Quark masses in Composite Higgs through partial compositeness



Quark masses in Composite Higgs through partial compositeness



The Higgs potential

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

Higgs potential generated at one loop with SM-fermions

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

Higgs potential generated at one loop with SM-fermions

$$V_h = g_\star^2 f^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{f}\right)$$

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

Higgs potential generated at one loop with SM-fermions

Dimensional
Analysis

$$V_h = g_\star^2 f^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{f}\right)$$

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

Higgs potential generated at one loop with SM-fermions

Dimensional Analysis

Loop factor

$$V_h = g_\star^2 f^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{f}\right)$$

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

Higgs potential generated at one loop with SM-fermions

Dimensional Analysis

Loop factor

Expansion in small explicit breaking

$$V_h = g_\star^2 f^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{f}\right)$$

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

Higgs potential generated at one loop with SM-fermions

Dimensional Analysis

Loop factor

Expansion in small explicit breaking

Trigonometric functions (Goldstone nature of Higgs)

$$V_h = g_\star^2 f^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{f}\right)$$

The Higgs potential

Explicit breaking of G through elementary-composite couplings:

$$\mathcal{L} \supset y_i f \bar{q}_i U \psi_i$$



SM-fermions don't
fill complete multiplets of G

Higgs potential generated at one loop with SM-fermions

Dimensional Analysis

Loop factor

Expansion in small explicit breaking

Trigonometric functions (Goldstone nature of Higgs)

$f \sim \langle \chi \rangle$

$$V_h = g_\star^2 f^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{f}\right)$$

Tuning of the Higgs potential

$$V_h = g_\star^2 \chi^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{\chi}\right)$$

Tuning of the Higgs potential

$$V_h = g_\star^2 \chi^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{\chi}\right)$$

Chose characteristic $\mathcal{I}_i \rightarrow \sin^2, \sin^4$

$$V_h^0 = \alpha^0 \sin^2 \left(\frac{h}{\chi}\right) + \beta^0 \sin^4 \left(\frac{h}{\chi}\right)$$

Tuning of the Higgs potential

$$V_h = g_\star^2 \chi^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{\chi}\right)$$

Chose characteristic $\mathcal{I}_i \rightarrow \sin^2, \sin^4$

$$V_h^0 = \alpha^0 \sin^2 \left(\frac{h}{\chi}\right) + \beta^0 \sin^4 \left(\frac{h}{\chi}\right)$$

Reproduce Higgs
mass and vev at

$$\chi = f$$

Tuning of the Higgs potential

$$V_h = g_\star^2 \chi^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{\chi}\right)$$

Chose characteristic $\mathcal{I}_i \rightarrow \sin^2, \sin^4$

$$V_h^0 = \alpha^0 \sin^2 \left(\frac{h}{\chi}\right) + \beta^0 \sin^4 \left(\frac{h}{\chi}\right)$$

Reproduce Higgs mass and vev at $\chi = f$ $\Rightarrow \alpha^0 = -2\beta^0 \sin^2(v/f), \quad \beta^0 = \frac{1}{8} m_h^2 f^2 / \sin^2(v/f)$

Tuning of the Higgs potential

$$V_h = g_\star^2 \chi^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{\chi}\right)$$

Chose characteristic $\mathcal{I}_i \rightarrow \sin^2, \sin^4$

$$V_h^0 = \alpha^0 \sin^2 \left(\frac{h}{\chi}\right) + \beta^0 \sin^4 \left(\frac{h}{\chi}\right)$$

Reproduce Higgs mass and vev at $\chi = f$ $\Rightarrow \alpha^0 = -2\beta^0 \sin^2(v/f), \quad \beta^0 = \frac{1}{8} m_h^2 f^2 / \sin^2(v/f)$

$$|\alpha^0/\beta^0| = 2 \sin^2(v/f)$$

Tuning of the Higgs potential

$$V_h = g_\star^2 \chi^4 \frac{y^2}{(4\pi)^2} \sum_i c_i \left(\frac{y}{g_\star}\right)^{p_i} \mathcal{I}_i \left(\frac{h}{\chi}\right)$$

Chose characteristic $\mathcal{I}_i \rightarrow \sin^2, \sin^4$

$$V_h^0 = \alpha^0 \sin^2 \left(\frac{h}{\chi}\right) + \beta^0 \sin^4 \left(\frac{h}{\chi}\right)$$

Reproduce Higgs mass and vev at $\Rightarrow \alpha^0 = -2\beta^0 \sin^2(v/f), \quad \beta^0 = \frac{1}{8}m_h^2 f^2 / \sin^2(v/f)$

$$\chi = f$$

$$|\alpha^0/\beta^0| = 2 \sin^2(v/f) \quad \sin^2 \left(\frac{v}{f}\right) \lesssim 0.1 \dots 0.2$$

Full potential

$$V = V_\chi + (\chi/f)^4 V_h^0 + (V_h^{NDA}[y] - V_h^{NDA}[y_0])$$

Full potential

$$V = V_\chi + (\chi/f)^4 V_h^0 + (V_h^{NDA}[y] - V_h^{NDA}[y_0])$$



Dilaton potential

Full potential

$$V = V_\chi + (\chi/f)^4 V_h^0 + (V_h^{NDA}[y] - V_h^{NDA}[y_0])$$



Dilaton potential



Tuned Higgs
potential

Full potential

$$V = V_\chi + (\chi/f)^4 V_h^0 + (V_h^{NDA}[y] - V_h^{NDA}[y_0])$$

Dilaton potential Tuned Higgs
potential

$$V_h^{NDA}[y] = \underbrace{c_\alpha \sum_{n_f} g_\star^2 \frac{N_c y^2[\chi]}{(4\pi)^2} \chi^4 \sin^2 \frac{h}{\chi}}_{\alpha} + \underbrace{c_\beta \sum_{n_f} g_\star^2 \frac{N_c y^2[\chi]}{(4\pi)^2} \left(\frac{y}{g_\star}\right)^{p_\beta} \chi^4 \sin^4 \frac{h}{\chi}}_{\beta}$$

Full potential

$$V = V_\chi + (\chi/f)^4 V_h^0 + (V_h^{NDA}[y] - V_h^{NDA}[y_0])$$

Dilaton potential Tuned Higgs
potential

$$V_h^{NDA}[y] = \underbrace{c_\alpha \sum_{n_f} g_\star^2 \frac{N_c y^2[\chi]}{(4\pi)^2} \chi^4 \sin^2 \frac{h}{\chi}}_{\alpha} + \underbrace{c_\beta \sum_{n_f} g_\star^2 \frac{N_c y^2[\chi]}{(4\pi)^2} \left(\frac{y}{g_\star}\right)^{p_\beta} \chi^4 \sin^4 \frac{h}{\chi}}_{\beta}$$

$\alpha, \beta > 0 \Rightarrow$ **valley along** $h/\chi = 0$ **Trajectory a**

Full potential

$$V = V_\chi + (\chi/f)^4 V_h^0 + (V_h^{NDA}[y] - V_h^{NDA}[y_0])$$



$$V_h^{NDA}[y] = \underbrace{c_\alpha \sum_{n_f} g_\star^2 \frac{N_c y^2[\chi]}{(4\pi)^2} \chi^4 \sin^2 \frac{h}{\chi}}_{\alpha} + \underbrace{c_\beta \sum_{n_f} g_\star^2 \frac{N_c y^2[\chi]}{(4\pi)^2} \left(\frac{y}{g_\star}\right)^{p_\beta} \chi^4 \sin^4 \frac{h}{\chi}}_{\beta}$$

$\alpha, \beta > 0$	\Rightarrow	valley along	$h/\chi = 0$	Trajectory a
$\alpha, \beta < 0$	\Rightarrow	valley along	$h/\chi \sim \pi/2$	Trajectory c