



Contribution ID: 253

Type: **Oral Presentation**

## Overview of recent STAR open heavy flavor measurements

*Tuesday, 27 August 2019 13:00 (30 minutes)*

The experiments at RHIC and LHC have produced convincing evidence that strongly interacting partonic matter, Quark Gluon Plasma (QGP), is created in central ultrarelativistic collisions of heavy ions. Charm quarks are ideal probe of the QGP since they are dominantly produced in hard gluon-gluon interactions in early stages of nuclear collisions at RHIC energies. Therefore, they are sensitive to the whole evolution of the hot and dense matter. Thanks to the excellent vertex resolution provided by the Heavy Flavor Tracker detector, STAR is able to measure charm quark production in multiple channels via reconstruction of hadronic decays of  $D^0$ ,  $D^{+/-}$ ,  $D_s$  and  $\Lambda_c$  hadrons. In this talk we will present recent STAR measurements of nuclear modification factors of  $D^0$  and  $D^{+/-}$  mesons,  $D^0$  directed, elliptic and triangular flow and  $D_s/D^0$  and  $\Lambda_c/D^0$  yield ratios. These measurements will be discussed in context of charm quark energy loss, charm quark transport in the QGP and final hadronization.

**Primary author:** BIELCIK, Jaroslav (Czech Technical University (CZ))

**Presenter:** BIELCIK, Jaroslav (Czech Technical University (CZ))

**Session Classification:** Workshop on Heavy Ion Physics