Strangeness in Quark Matter 2019



Contribution ID: 186

Type: Contributed talk

Directed flow, vorticity and Lambda-Polarisation in heavy-ion collisions at FAIR and NICA energies

Tuesday 11 June 2019 16:50 (20 minutes)

The directed flow, vorticity and polarisation of Lambda were calculated within the UrQMD model for intermediate-centrality A+A collisions at FAIR and NICA energies. Also, we investigated vorticity and helicity development of protons and pions in these reactions. The slope of the directed flow dv_1/dy at midrapidity in calculations with and without mean-field potentials is obtained as a function of the beam energy for $p, p^-, \Lambda, \Lambda^-, K^+, K^-, \pi^+, \pi^-$. Results of the calculations are compared to the STAR experimental data.

Collaboration name

Track

Hydrodynamics, chirality and vorticity

Primary authors: ZABRODIN, Evgeny (University of Oslo (NO)); BRAVINA, Larisa (University of Oslo (NO)); VITIUK, Oleksandr (Taras Shevchenko National University of Kyiv); SORIN, Alexander (Joint Institute for Nuclear Research (RU)); TERYAEV, Oleg (JINR)

Presenter: VITIUK, Oleksandr (Taras Shevchenko National University of Kyiv)

Session Classification: Hydrodynamics, Chirality and Vorticity